

**WARNING:** Harmful or fatal if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of reach of children. Keep children and pets off treated area until spray is washed and dried. Do not use on newly seeded lawns.

This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning. (Keep out of smoke).

**DIRECTIONS FOR USE**

**HOUSEHOLD PESTS:** For control of Roaches, Ants, Silverfish, Waterbugs, Crickets, Spiders, Centipedes and Box Elder Bugs add 3 to 4 tablespoons in 2 quarts of water. Thoroughly spray infested cracks and crevices, baseboards, "behind shelving and bookcases for silverfish", storage area and water drains in basements. For Ants spray ant trails. For Ants and Box Elder Bugs treat window frames, door sills, openings around water pipes, electrical outlets and other places where they may enter. For Flies, Mosquitoes and Wasps mix at same rate and spray screens, doors, door sills, window frames and areas frequented by these pests such as porches, and refuse disposal area. Repeat application as needed for control of any insect listed.

**LAWN INSECTS:** For following insects add the designated number of tablespoons to 2 gallons for 1,000 square feet. For Ants, Armyworms, Crickets, Grasshoppers and Spiders use ¼ tablespoon. Spray lawn area and ornamentals as needed. For Ants treat hills and trails. For Chiggers, Earwigs and Ticks use 1½ tablespoons, and thoroughly treat turf and ornamentals. For Chinch Bugs and

ACCEPTED  
APR 20 1973

**72% CHLORDANE  
EMULSIFIABLE  
CONCENTRATE**

**ACTIVE INGREDIENTS:**

Chlordane Technical \* ..... 72.0%  
Petroleum Hydrocarbons ..... 23.5%

**INERT INGREDIENTS** ..... 4.5%  
Total ..... 100.0%

\*Equivalent to 43.0% octachloro-4,7 methanotetrahydroindane and 29.0% of related compounds.

**For Controlling Soil Insects — Grasshoppers**

**WARNING:**

**KEEP OUT OF REACH OF CHILDREN**

See Side Panel For Additional Cautions

**COLORADO INTERNATIONAL CORP., COMMERCE CITY, COLO. 80022**

Webworms use 2¼ tablespoons and treat a  
For White Grubs and Wireworms use 4  
Wash into soil immediately after application  
on newly seeded lawns.

**CRABGRASS SUPPRESSION:** Mix 4 oz water to cover 160 square feet (1½ pint water per 1,000 square feet) of lawn or to crabgrass seedlings as germination or sprout (Around April 15th). Will not injure established lawn. This spray will also control lawn insect "For Lawn". Keep children and pets off until the spray has been washed into the soil and has completely dried. Do not use on newly seeded lawns.

**VEGETABLE GARDENS:** For the following crops use the designated number of tablespoons of concentrate per 1,000 square feet of water and spray to cover 1,000 square feet.

**BEANS:** For Flea Beetles, Green Stink Beetles use 1½ tablespoons. Do not apply to green or snap beans until pods begin to form.

**CABBAGE:** For Imported Cabbage Worms, Cabbage Moth Larvae, Flea Beetles, Green Leafhoppers, Harlequin Bugs, Mole Crickets use 1½ tablespoons. Apply when injury is imminent. Do not use until heads begin to form.

**POTATOES:** Broadcast soil application prior to planting use 10 lbs. actual Chlordane per acre (5 lbs. actual Chlordane per acre). Harrow or disk in the soil prior to planting. For Flea Beetles and Flea Beetles, use 1 tablespoon per 1,000 square feet. They appear and as needed.

**ONIONS:** For Thrips use 1½ tablespoons per 1,000 square feet. They appear. Do not use on Green or Spring Onions.

**CUTWORMS:** use 1½ tablespoons per 1,000 square feet. base of newly set plants such as Lettuce, Spinach, and Okra. Do not apply after edible portion has formed.

**GRASSHOPPERS:** To protect garden vegetation adjacent to garden; use 1½ tablespoons per 1,000 square feet. Spray directly affected flower and ornamentals.

**NOTICE:** Seller warrants that this product conforms to the ingredient statement on the label. Since use, such as weather, compatibility with other products and condition of application equipment varies, we make no claims other than those stated on the label. EPA Reg. No. 4715-301 Net Content

ACCEPTED  
APR 20 1973  
U.S. DEPARTMENT OF AGRICULTURE  
OFFICE OF PEST CONTROL AND PLANT QUARANTINE  
WASHINGTON, D.C.

Harmful or fatal if swallowed. Contact with  
 eye toxic symptoms. Avoid breathing spray  
 of contact with skin, wash with soap and  
 contamination of feed and foodstuffs. Keep  
 of children. Keep children and pets off  
 until spray is washed and dried. Do not use  
 on lawns.

is toxic to fish and wildlife. Keep out of any  
 water. Birds feeding on treated areas may be  
 killed. Do not apply where runoff is likely to occur. Do  
 not apply when weather conditions favor drift from  
 treated areas. Do not contaminate water by cleaning of  
 disposal of wastes. Apply this product only  
 in accordance with this label.

is highly toxic to bees exposed to direct  
 residues on crops. Protective information  
 printed from your Cooperative Agricultural  
 Extension Office.

Do not use in or near water and containers and dispose of wastes by  
 in-crop lands away from water supplies.  
 Wastes should be disposed of by punching holes in them  
 and burying with wastes, or by burning. (Keep out of

**DIRECTIONS FOR USE**

**PESTS:** For control of Roaches, Ants,  
 Fleas, Earwigs, Crickets, Spiders, Centipedes and  
 other insects add 3 to 4 tablespoons in 2 quarts of  
 water and spray infested cracks and crevices,  
 behind shelving and bookcases for silver-  
 leaf area and water drains in basements. For  
 trails. For Ants and Box Elder Bugs treat  
 door sills, openings around water pipes,  
 basements and other places where they may enter.  
 Mosquitoes and Wasps mix at same rate and  
 apply to door sills, window frames and areas  
 where these pests such as porches, and refuse  
 repeat application as needed for control of  
 these pests.

**INSECTS:** For following insects add the designated  
 amount to 2 gallons for 1,000 square feet.  
 Fleas, Wireworms, Crickets, Grasshoppers and  
 Beetles. Spray lawn area and ornamentals.  
 Ants treat hills and trails. For Chiggers,  
 Fleas use 1 1/2 tablespoons, and thoroughly  
 spray ornamentals. For Chinch Bugs and

ACCEPTED  
 APR 20 1973

**72% CHLORDANE  
 EMULSIFIABLE  
 CONCENTRATE**

**ACTIVE INGREDIENTS:**

Chlordane Technical .....	72.0%
Petroleum Hydrocarbons .....	23.5%
INERT INGREDIENTS .....	4.5%
Total .....	100.0%

\*Equivalent to 43.0% octachloro-4,7 methanotetra-  
 hydroindane and 29.0% of related compounds.

**For Controlling Soil Insects – Grasshoppers**

**WARNING:**

**KEEP OUT OF REACH OF CHILDREN**

See Side Panel For Additional Cautions

**COLORADO INTERNATIONAL CORP., COMMERCE CITY, COLO. 80022**

Webworms use 2 1/4 tablespoons and treat as for Chiggers.  
 For White Grubs and Wireworms use 4 tablespoons.  
 Wash into soil immediately after application. Do not use  
 on newly seeded lawns.

**CRABGRASS SUPPRESSION:** Mix 4 oz. in 1 gallon  
 water to cover 160 square feet (1 1/2 pints in 6 gallons  
 water per 1,000 square feet) of lawn or turf to suppress  
 crabgrass seedlings as germination or sprouting occurs.  
 (Around April 15th). Will not injure established lawns.  
 This spray will also control lawn insects listed under  
 "For Lawn". Keep children and pets off treated areas  
 until the spray has been washed into the soil and grass  
 has completely dried. Do not use on newly seeded lawns.

**VEGETABLE GARDENS:** For the following insects add  
 the designated number of tablespoons to 2 gallons of  
 water and spray to cover 1,000 square feet.

**BEANS:** For Flea Beetles, Green Stink Bugs use 1 1/2  
 tablespoons. Do not apply to green or snap beans after the  
 pods begin to form.

**CABBAGE:** For Imported Cabbage Worm, Diamond-  
 Back Moth Larvae, Flea Beetles, Green Stink Bugs,  
 Harlequin Bugs, Mole Crickets use 1 1/2 tablespoons.  
 Apply when injury is imminent. Do not make application  
 after heads begin to form.

**POTATOES:** Broadcast soil application prior to planting,  
 use 10 lbs. actual Chlordane per acre (5 qts. per acre).  
 Harrow or disk in the soil prior to planting. For  
 Flea Beetles and Flea Beetles, use 1 tablespoon. Apply when  
 they appear and as needed.

**ONIONS:** For Thrips use 1 1/2 tablespoons. Apply when  
 they appear. Do not use on Green or Spring Onions.

For **CUTWORMS** use 1 1/2 tablespoons. Apply around  
 base of newly set plants such as Lettuce, Peppers, Beets  
 and Okra. Do not apply after edible portions begin to  
 form.

For **GRASSHOPPERS:** To protect gardens spray waste  
 vegetation adjacent to garden; use 1 1/2 tablespoons diluted  
 as above. Spray directly affected flowering plants and  
 ornamentals.

**NOTICE:** Seller warrants that this product conforms to  
 the ingredient statement on the label. Since conditions of  
 use, such as weather, compatibility with other chemicals,  
 and condition of application equipment will vary, Seller  
 makes no claims other than those stated on this label.

EPA Reg. No. 4715-301 Net Contents \_\_\_\_\_

ACCEPTED  
 APR 20 1973