

WARNING: KEEP OUT OF REACH OF CHILDREN SEE BACK PANEL FOR ADDITIONAL WARNINGS

NET WEIGHT\_\_\_\_LBS.

U.S.D.A. REG. NO. 5185-90

# ACTIVE INGREDIENTS:

Trichloro-s-trizinetrione
(Available Chlorine 60%)

### INERT INGREDIENTS:

# TOTAL INGREDIENTS

When used as directed, this product is electing agent.

The filter system should be backwashed should be between 7.4 and 7.6 adjusting

It is recommended that initial users of the water with 30 ppm Trihydroxy triazine. If the dissipation effect of sunlight on child

If there is no initial stabilization with To 3-4 weeks' treatment with BIO-GUARD Co 66 will gradually develop the desired level both chlorine and Trihydroxy triazine are

In addition to the normally recommend pools require extra dosages periodicall

ANTIDOTE: Call a physician.

EXTERNAL: For skin and eyes,

large amounts of

medical attention.

INTERNAL: Take large amount

olive oil or cooki

ACCEPTED

#### ACTIVE INGREDIENTS:

Trichloro-s-trizinetrione 66.52%
(Available Chlorine 60%)

INERT INGREDIENTS: 33.48%

TOTAL INGREDIENTS 100.00%

When used as directed, this product is effective as a pool water disinfecting agent.

The filter system should be backwashed or properly cleaned. The pH should be between 7.4 and 7.6 adjusting if necessary.

It is recommended that initial users of his product first stabilize pool water with 30 ppm Trihydroxy triazine. This will stabilize water against the dissipation effect of sunlight on chlorine.

If there is no initial stabilization with Trihydroxy triazine, a period of 3-4 weeks' treatment with BIO-GUARD CHLORINE CONCENTRATE NO. 66 will gradually develop the desired level. When using this product, both chlorine and Trihydroxy triazine are being added to the water.

In addition to the normally recommended amounts of chlorine most pools require extra dosages periodically, which are referred to as

superchlorination. This is recommended weekly and the amount is approximately four times the daily dosage. Application should be made after sunlight hours and when the pool is not in use. Superchlorination should not exceed 8 ppm available chlorine.

#### DIRECTIONS

If there is an initial stabilization with Trihydroxy triazine, use 3 ounces BIO-GUARD CONCENTRATE NO. 66 per 10,000 gallons of water. Through regular use of a test kit, adjust amount added to maintain pool water at 1.0-1.5 ppms of residual chlorine. There will be a rapid reduction in the required amounts of chlorine, if the pool has been properly prestabilized.

If there is no initial stabiliation, use 3-4 ounces of BIO-GUARD CHLO-RINE CONCENTRATE NO. 66 daly per 10,000 gallons water. In 3-4 weeks the pool will become stabilized. Through regular use of a test kit, adjustment should be made in the amount added to maintain the pool at 1.0-1.5 ppms of residual chlorine.

Using either of the above methods, super-chlorination is recommended weekly. The use of a quartenary ammonium type algicide will complement this total program of swimming pool maintenance. Maintain pool water total alkalinity between 80 and 120 ppms.



ANTIDOTE: Call a physician.

EXTERNAL: For skin and eyes, immediately flood with

large amounts of water. For eyes get

medical attention.

INTERNAL: Take large amounts of milk, followed by

olive oil or cooking oil.

WARNING! Oxidizing material. Contact with heat, moisture or foreign materials may result in fire. Will burn with the evolution of irritation and hazardous gases.

Store in a cool dry place. Keep away from heat or open flame. Avoid contact with skin and eyes. Avoid breathing dust. DO NOT TAKE INTERNALLYI Harmful if swallowed. Keep container closed when not in use.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Cameraman's Note

U.S.D.A. REG. NO. 5185-90
NET WEIGHT\_\_\_\_LBS.

Poor Copy

ACCEPTED

185-90

R