ACCEPTED

FEB 25 1974

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO.

BIO-QUAT 20

BIO-QUAT 20 is an efficient disinfectant of high germicidal and bacteriostatic potency. When used as directed, BIO-QUAT 20 is odorless, and non-staining.

DIRECTIONS

BIO-QUAT 20 is a concentrate; dilute as directed for use.

For Use in Poultry Houses and Equipment:

- 1. Remove all poultry and feeds from premises, coops and crates.
- 2. Remove all litter and droppings from floors, walls and surfaces or facilities occupied or traversed by poultry. Empty all troughs, racks and other feeding and watering appliances.
- 3. Thoroughly clean all surfaces with a compatible nonionic detergent. Rinse away loosened soils.
- 4. To disinfect non-porous surfaces or equipment (glass, plastic, smooth metal), thoroughly wet with a solution containing 1 ounce of BIO-QUAT 20 per 4 gallons of water.
- 5. To disinfect porous or highly contaminated surfaces (floors, walls, ceilings, rafters, curtains, joists), apply a wetting concentration of a solution containing 1 ounce of BIO-QUAT 20 per 2 gallons of water.
- 6. Ventilate buildings, coops and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set or dried.
- 7. Treated feeders and waterers must be thoroughly rinsed with potable water prior to reuse.

MOLD CONTROL OF WALLS, CEILING IN FOOD PLANTS: Clean thoroughly, rinse, then spray or sponge with solution containing one sunce of BIO-QUAT 20 to one gallon of water. Spray or sponge surfaces until moist and repeat applications weekly or when new growth appears.

BIO-QUAT 20

ACTIVE INGREDIENTS:

Alkyl (C12, 70%; C14, 30%) Dimethyl	
Benzyl Ammonium Chloride	20%
INERT INGREDIENTS:	80%
TOTAL	100%

DANGER

KEEP OUT OF REACH OF CHILDREN

See First Aid statement and other precautions on side panel.

A Product Of

BIO-LAB, INC.

DECATUR, GEORGIA

EPA Reg. No. 5185-110

NET CONTENTS:

BIO-QUAT 20

SANITATION OF EQUIPMENT IN FOOD PROCESSING PLANTS, IN-CLUDING MEAT PLANTS: Clean and rinse equipment thoroughly. Then apply a solution of one ounce of BIO-QUAT 20 to 4 gallons of water. Flush equipment with a potable water rinse before reusing.

MOLD CONTROL OF WALLS, CEILING IN FOOD PLANTS: Clean thoroughly, rinse, then spray or sponge with solution containing one ounce of BIO-QUAT 20 to one gallon of water. Spray or sponge surfaces until moist and repeat applications weekly or when new growth appears. IMPORTANT: Soaps reduce the germicidal efficiency of BIO-QUAT 20. Care should be taken to rinse objects thoroughly before placing in BIO-QUAT solution. Thoroughly clean all objects and surfaces before disinfecting with BIO-QUAT solution.

DANGER:

Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard it.