

Mock's Cura-Lawn will effectively prevent turf-grass damage by many diseases, and will limit the damage done by these infections if treatment is started promptly at the first sign of symptoms. It cannot, however, bring dead grass back to life, and it is important that the damaged area be reseeded and fertilized with the proper Super-turf product as soon as the disease is arrested.

Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with eyes, skin or clothing. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuff. Keep children and pets off treated area until material is washed into soil and grass is dry.

To protect fish and wildlife, do not contaminate streams, lakes or ponds with this material.

Internal: Emetic of tablespoon of mustard in Tumbler of water. Call a physician.

ACCEPTED
Jun 22-1967

A **Superturf** LAWN PRODUCT

MOCK SEED COMPANY, PITTSBURGH, PA. 15230

5738-15

CAUTION — KEEP OUT OF REACH OF CHILDREN

See Rear Panel For Additional Cautions

Active Ingredients

2,4-Dichloro-6-(o-chloroanilino)-s-triazine _____	3.90%
A co-ordination product of zinc ion and manganese ethylene bisdithiocarbamate _____	3.90%
In which the ingredients are:	
Manganese ⁺⁺ _____	0.78%
Zinc ⁺⁺ _____	0.10%
Ethylene bisdithiocarbamate ion (C ₄ H ₆ N ₂ S ₄) _____	3.02%
Inert Ingredients _____	92.20%
TOTAL _____	100.00%

USDA Reg. No. 5238-15

MOCK SEED COMPANY, PITTSBURGH, PA. 15230

10 LBS. NET WT.

**Double
Action!**

**2,500
FOOT SIZE**

MOCK'S

CURA-LAWN

For Control of Many Turfgrass Diseases

A Superturf
LAWN PRODUCT

CAUTION — KEEP OUT OF REACH OF CHILDREN

See Back Panel for Additional Caution

Active Ingredients

2,4-Dichloro-4-(4-chlorophenoxy)phenol	2.0%
A co-crystalline product of the isomers and respective ethylene dimethacrylates	2.0%
In which the ingredients are:	
Mecoprop-P	0.5%
Diquat	0.5%
Phytol 2,4-Dichlorophenoxy	1.0%
Mecoprop-P	1.0%
Diquat	1.0%

Net Weight 2.500 lbs. (1134 kg)

A Superturf

LAWN PRODUCT

CAUTION — KEEP OUT OF REACH OF CHILDREN

See Rear Panel For Additional Cautions

Active Ingredients

2,4-Dichloro-6-o-chloroanilino-s-triazine	3.90%
A co-ordination product of zinc ion and manganese ethylene bisdithiocarbamate	3.90%
In which the ingredients are:	
Manganese ⁺⁺	0.70%
Zinc ⁺⁺	0.10%
Ethylene bisdithiocarbamate ion (C ₂ H ₂ N ₂ S ₄)	3.02%
Inert Ingredients	92.20%
TOTAL	100.00%

USA Reg. No. 2512-15
MCKEE SEED COMPANY, PITTSBURGH, PA. 15220
10 LBS. NET WT.

REGISTERED
Jun-29-1967
52287
5228-15

10 LBS. NET WEIGHT

MOCK'S CURA-LAWN

Contains DYRENE* and DITHANE M-45**

TREATS 2,500 SQUARE FEET OF LAWN AREA

*Reg. U.S. Pat. Off. by Chemagro Corp.

**Reg. U.S. Pat. Off. by Rohm & Haas Co.

Cura-Lawn controls many fungus diseases of lawns, golf greens, and other fine turf areas. By combining the fungicidal properties of two of the most effective materials for prevention and control of turf diseases, Mock's have developed a product that has an unusually "wide spectrum" of disease control. Furthermore, since Cura-Lawn is a dry, granular material, it can be applied with a spreader at times when a sprayer is not practical, such as in freezing weather.

Diseases controlled by Cura-Lawn are: Snow Mold (Typhula Blight and Fusarium Patch), Sclerotinia Dollar Spot, Copper Spot, Red Thread, Helminthosporium Blight (Leaf Spot and Melting Out), Rhizoctinia Brown Patch, Fusarium Roseum Blight, and Rust.

	Preventive	Curative
Mock's	8	9
Scott's	6	7
Lawn Beauty	7	8

NOTE: The above spreader settings are approximate. For absolute accuracy, measure out 100 square feet, and adjust your spreader to apply exactly 6½ ounces of Cura-Lawn on this measured area (or 9½ ounces, if you are using the curative rate). Rate of walk, moisture content, and spreader condition can vary the setting necessary to apply the correct amount.

With a calibrated spreader apply at the rate of 4 pounds per 1000 square feet over the lawn area for prevention of the diseases named on this package. Treatments should be repeated every two weeks for season long protection.

DO NOT WATER FOR 24 HOURS AFTER APPLYING. Application when grass is moist will often improve results.

If disease has already done considerable damage, apply at the rate of 6 pounds per 1,000 square feet (Curative Rate) at ten day intervals until improvement is noted. Protect against Snow Mold by applying to lawn in November at the Curative Rate. Repeat, if possible, during a mid-winter thaw.