

20-10-5

TEXACO

Quality Line

TEXACO

Nu-turf



Controls Weeds
as PLANTAIN-DANDELION-BUCKWORN

FEED & WEED



5

NET WEIGHT 22 LBS.

Jan 16, 1973

52885



Nu-turf
20-10-5
FEED & WEED

MINIMUM GUARANTEED ANALYSIS

LAWN FOOD

Total Nitrogen (N)	20.00%
4.0% Ammoniacal Nitrogen	
16.0% Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	10.00%
From Phosphoric Acid	
Soluble Potash (K ₂ O)	5.00%
From Muriate of Potash	
Potential acidity 900 lbs. calcium carbonate equivalent per ton.	

HERBICIDAL

ACTIVE INGREDIENTS

Dimethylamine Salts of 2,4-Dichlorophenoxyacetic Acid*	0.94%
Isooctyl Ester of Silvex (2-(2,4,5-Trichlorophenoxy) Propionic Acid)**	0.58%
INERT INGREDIENTS	98.48%
Total	100.00%

*Equivalent to 78% 2,4-Dichlorophenoxyacetic Acid
**Equivalent to 41% Silvex

A Product of
FEDERAL CHEMICAL COMPANY
LOUISVILLE, KENTUCKY 40202
A Subsidiary of TEXACO INC.

INSTRUCTIONS

This product is a 20-10-5 formula lawn food for feeding grasses while controlling broad leaf weeds such as dandelion, plantain, buckhorn, pig-weed, ragweed, sourgrass and many others. Can be used in Spring, Summer and Fall. Best results will be obtained when weeds are growing vigorously.

1. Spread the contents of this bag over 5000 sq. ft.
2. Apply when grass is wet from dew or sprinkling.
3. Do not apply when rain is predicted and do not water the lawn for 24 hours after application.
4. Do not treat new lawns until after third mowing.
5. Do not apply on windy days to avoid drift to surrounding plants.
6. Avoid application around or near evergreens, shrubs and other flowering plants.

20-10-5

TEXACO

TEXACO

Nu-turf



COVERS 5,000 SQ. FT. (50'x100')

NET WEIGHT 22 LBS.

TEXACO

20-10-5

NET WEIGHT 22 LBS



Nu-turf
20-10-5
FEED & WEED

20% Nitrogen	10.0%
10% Phosphorus	5.0%
5% Potassium	2.5%
10% Iron	5.0%
5% Manganese	2.5%
5% Zinc	2.5%
5% Boron	2.5%
5% Copper	2.5%
5% Sulfur	2.5%
5% Magnesium	2.5%
5% Calcium	2.5%
5% Silicon	2.5%
5% Selenium	2.5%
5% Vanadium	2.5%
5% Nickel	2.5%
5% Molybdenum	2.5%
5% Cobalt	2.5%
5% Fluorine	2.5%
5% Chlorine	2.5%
5% Iodine	2.5%
5% Bromine	2.5%
5% Barium	2.5%
5% Strontium	2.5%
5% Zirconium	2.5%
5% Niobium	2.5%
5% Manganese	2.5%
5% Silicon	2.5%
5% Selenium	2.5%
5% Vanadium	2.5%
5% Nickel	2.5%
5% Molybdenum	2.5%
5% Cobalt	2.5%
5% Fluorine	2.5%
5% Chlorine	2.5%
5% Iodine	2.5%
5% Bromine	2.5%
5% Barium	2.5%
5% Strontium	2.5%
5% Zirconium	2.5%
5% Niobium	2.5%

FEDERAL CHEMICAL COMPANY
10000 LEE ROAD, PHOENIX, ARIZONA 85022
A DIVISION OF TEXACO INC.

INSTRUCTIONS

1. This 20-10-5 Nu-turf is a lawn feed for feeding grasses when applied at the rate of 1/2 lb. per 1,000 sq. ft. Apply at 1/2 lb. per 1,000 sq. ft. or as directed on the label. Can be used in Spring, Summer or Fall. Best results will be obtained when weeds are growing vigorously.
2. Spread the contents of this bag over 5,000 sq. ft.
3. Apply when grass is wet from dew or watering.
4. Do not apply when rain is predicted and do not water the lawn for 24 hours after application.
5. Do not feed new lawns until after third mowing.
6. Do not feed on windy days to avoid drift to surrounding plants.
7. Avoid application around or near evergreens, shrubs and other flowering plants.

CAUTION

Avoid contact with eyes or clothing. Avoid absorption and contamination of feedstuffs. Wash hands after use. Do not apply to flowers, vegetables or ornamentals. Do not apply to St. Augustine grass, bermudagrass, or Bahia grass. Stay away from water holes, irrigation canals and ditches. Avoid contamination of irrigation ditches or other water for domestic purposes. Vapor from this product may be flammable in the immediate vicinity. This product is for use on lawns. Keep out of holes, ditches or ponds. Do not reuse container. Destroy by burning in a safe place away from other buildings. Do not burn.

SUGGESTED SPREADER SETTINGS

Rate	1/2 lb. per 1,000 sq. ft.
Setting	1/2 lb. per 1,000 sq. ft.
Direction	1/2 lb. per 1,000 sq. ft.
Speed	1/2 lb. per 1,000 sq. ft.
Height	1/2 lb. per 1,000 sq. ft.

These specific settings are suggested as a guide to application. They are not intended as a guarantee of results. Results will vary with soil conditions and type of spreader used.