

©  
**CHACON**

ACCEPTED

FEB 6 1987

U.S. Department of Agriculture  
Fertilizer Act, as amended, for the pesticide  
registered under EPA Reg. No. 5719-110

# MOLE CRICKET CONTROL

ALSO CONTROLS  
GRUB WORMS CHINCH BUGS



AND OTHER TURF GRASS INSECTS

ACTIVE INGREDIENT  
 1-Methyl-2-(2-chlorophenyl)-2-methyl-1-propanol phosphorothioic acid, benzylate  
 INERT INGREDIENTS

U.S. Patent No. 3,671,087  
 EPA REG. NO.

EPA EST. NO. 44616 MO 1

1.5%  
 55.5%



READ THE LABEL BEFORE USE

**STORAGE AND DISPOSAL**

Household: Do not reuse empty container. Wrap and put in trash collection or pesticide disposal (except household). Pesticide that cannot be used according to label directions must be disposed of according to Federal, State, or local procedures under the Resource Conservation and Recovery Act or other disposal (except household). Do not use container in connection with food, feed, or drinking water. Dispose of in a sanitary way or by incineration if permitted by State and local authorities.  
 Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store original container and out of the reach of children.  
 Rinse and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Do not walk through spilled material. Dispose of pesticide as directed above.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

**PRECAUTIONARY STATEMENTS**

SEE STATEMENTS OF PRACTICAL TREATMENT AND ADDITIONAL PRECAUTIONARY STATEMENTS.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION**

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Keep out of reach of children.

**STATEMENT OF PRACTICAL TREATMENT**

If in eyes: Flush with plenty of water. If on skin: Wash with soap and hot water.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and wildlife. Do not apply directly to water or wetlands. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read these entire Directions for Use before using. This insecticide is recommended for use on lawns and other ornamental and recreational turf grass areas.

**APPLICATION EQUIPMENT**

Apply over the turf area with a suitable granular turf application equipment. Both drop type or rotary type spreaders may be used. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly.

**APPLICATION TECHNIQUE**

Application using a pull type drop spreader can be aided by a splash board mounted under the hopper for more uniform granule distribution. A more uniform application can be made by spreading half the required amount of Ontario Granules over the area, and then immediately apply the remaining half at right angles or crosswise. The turf recommended rate must be used from maximum weight. When applying, use appropriate marking systems and make application when the wind is calm to prevent missed areas.

**RESTRICTIONS:**

Do not mow treated turf area before rainfall or irrigation. Do not graze treated areas or use clippings from treated areas for feed or forage.

**SPREADER SETTING**

Set spreader for an application rate of 3 lbs. per 1000 sq. ft. of area.

MODEL	SETTING
Cyclone, model B-1	3.5
Scotts PF-1/011B	6.0

**RECOMMENDED APPLICATIONS FOR TURF GRASSES**

CROP	PEST	DOSAGE	REMARKS
Turf Grasses	SOL INSECTS White grub larvae (such as Japanese beetle, Black turfgrass anthracnose, Crickets, Phyllocnistis coccana, Green June beetle, Oriental beetle, Asiatic garden beetle)	3 lbs. per 1000 sq. ft.	Be Sure to Read "Application Equipment" Section of the label.  Do not apply more than 6 lbs. per 1000 sq. ft. per year or an equivalent amount of 1.45 ounces active ingredients per 1000 sq. ft. regardless of formulation used. Minimum interval between applications must be 30 days.  For White Grub and Boring Larval Control: Water turf area thoroughly within 24 hours of application, using sufficient water to wet soil to 1 1/2 inches deep.
	Boring (larval stage)		
	Hyperodes weevil (larval stage)		For Hyperodes Weevil Control: Apply between full bloom periods of forsythia and flowering dogwood, usually April 15 to May 15, and water in as for white grub larval control.
Mole Crickets			For Mole Cricket Control: Water immediately after application using 1/2 to 1 inch of water. Consult your local turf, state Agriculture Experiment Station, or state Extension Service Specialists for more specific information regarding timing of application.
SURFACE INSECTS Sod Webworm larvae Chinch bug		1.5 or 3 lbs. per 1000 sq. ft.	Apply 1/2 to 1 inch of water within 24 hours of treatment. Reapply as necessary for pest re-infestations. Do not apply more than 6 lbs. per 1000 sq. ft. per year or an equivalent amount of 1.45 ounces active ingredients per 1000 sq. ft. regardless of formulation used. Minimum interval between applications is 15 days for 1.5 lb. rate; 30 days for 3 lb. rate.  Consult your local turf, state Agriculture Experiment Station, or state Extension Service Specialists for more specific information regarding timing of application.

Ontario is a Reg. TM of the Parent Company of Farbwerkebrunnen Bayer GmbH, Leverkusen

NET WEIGHT 9 LBS.

Manufactured for  
**CHACON CHEMICAL CORPORATION**  
 2600 Yates Avenue • City of Commerce, CA 90040  
 A GAFAMERICA COMPANY

**BEST AVAILABLE COPY**