pm 23 5815-30 FEB 1 1995 Ms. Michelle Bell The Andersons, Lawn Fertilizer Division d/b/a Pree Flow Fertilizer P.O. Box 119 Maumee, Oil 43537 Dear Ms. Bell: Subject: Ine Andersons Graden Weed Preventer EPA Redistration No. 5815-35 Application Dated December 30, 1994, Request To Amend Labeling by Adding WPS Exclusionary Statement and Using Split Labeling The proposed labeling submitted with your application dated march 25, 1994 has been reviewed and tound acceptable under the Federal Insecticide, Fungicide and Podenticide Act (FIFRA) as amended, provided that: And the same exclusionary statement to the "nomeowner use Only" laceling as the one you added to the "Commercial Use Only" labeling submitted with this application. Revise the labeling as described in this Agency's letter of october 17, 1994, in regards to when to apply this product. 3. Subbit one (1) copy of the final grinted labels before releasing the product for shipment under the subject labelinu. if these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, Section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copies of the labeling are enclosed for your records. Sincerely yours, Joanne I. Miller Product Manager (23) Pundicide-Herbicide Branch Registration Division (7505C) Enclosure E.milson:Diskette ABC17:02-01-95 BEST AVAILABLE COPY

5815-35C Page 1 of 5

(FRONT PANEL)

COMMERCIAL USE ONLY

with Committee in EPA Letter Design

FEB 1 (395

THE ANDERSONS GARDEN WEED PREVENTER

Unite the Fiction to readle, Fortunate, and Ross to the period to regardered under FPA Rep. No. 5815-35

ACTIVE INGREDIENT

TOTAL 100.00%

KEEF OUT OF REACH OF CHILDREN

CAUTION

If in Eyes: Flush with plenty of water. Call a physician if irritation persists.

If on Skir: Whom with glenty of scup and water. Call a physician if irritation persists.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 5815-35

EPA Est. 9198-OH-1

10 POUNDS NET WT.

COVERS 2,500 SQ. FT.

5815-350 Page 2 of 5

(BACK PANEL)

COMMERCIAL USE ONLY

THE ANDERSONS GARDEN WEED PREVENTER

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

CAUTION

Hazards to Humans and Domestic Animals

Harmil of evallowed. Caused moderate eye orbitation. Avoid contain with eyes or plotning. Wash thoroughly with soap and water after handling. Avoid breathing dust. Do not contaminate food and feedstuffs. Do not store hear seed, fertilizer or other herbicides.

ENVIRONMENTAL HAZARDS

Collect and incorporate granules spilled on the soil surface. Do not apply directly to water, or to areas where surface water is present or to intertigal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is in violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

DO NOT APPLY IN SUFFOLK COUNTY, LONG ISLAND, NEW YORK.

FOR USE ON: Beans (dry, field, snap), broccoli, Brussels sprouts, cappage, cantaloupe, cauliflower, collards, corn (field), corn (sweet), cotton, dress (upland), cucumber, eggplant, garlic, honeydew melon, horseradish, kale, leeks, lettuce (head), lettuce (leaf), mung beans, mustard greens, onlons, peppers, potato (whole or cut pieces), Southern beans (blackeyed peas), radish (grown for seed), soybeans, squash, strawberry, sweet potatoes, tomato, turnip (greens), turnip (roots), watermelon, yams (Ipomoea spp.).

5815-35C Page 3 of 5

(BACK PANEL)

COMMERCIAL USE ONLY

Also For Use On The Following Ornamentals and Trees: abelia, African violet, ageratum, ajuga, alyssum, arborvitae, ash, aster, athletic field turf, Austrian pine, azalea, baby's breath, barberry, bellflower, birch, bleeding heart, bloodleaf, blue bedder salvia, blue spruce, border forsythia, Boston ivy, boxweed, bugloss, California privet, camellia, candle larkspur, candy tuft, Cape marigold, chestnut, chrysanthemum, cinquefoil, coleus, columbine, coneflower, coral bells, coreopsis, cotoneaster, cottonwood, creeping juniper, creeping thyme, cuphea, cypress, dahlias, daisies, deutzia, dogweed, dwarf forsythia, elaeagnus, elm, English ivy, English ivy (ground cover), euonymus, evening primrose, fernleaf, yarrow, feverfew, fir, flowering crabapple, forget-me-not, forsythia, four-o-clock, foxglove, gaillardia, geranium, gladiolus, gold dust, golden marguerite, goldtuft, golf fairways/golf greens, ground-hemlock, gum, hawthorn, heath, holly, honeysuckle, hyrid tea roses, hydrangea, iris, ivy (ground cover), Japanese andromeda, Japanese parberry, Japanese privet, Japaese yew, juniper, lantana, larkspur, lavender cotton, Liquistrum, lilas, lilies, locicily pine, locust, lupines, magnolia, muple, marigold, mentor parcerry, mock-brange, morning-glory, mountain-laurel, multiflora rose, hasturtium, oak, orpine, pachysandra, pachysandra (ground tower, paxistima, peonies, petunias, pink, pittosporum, ponderosa pine, popular, podocarpus, purple coneflower, sed bine, redbud, rhododendron, rose, rode mous, Fischer blive, solver, abariosa, doarlet sage, Scotch pine, enore pine, snapdragon, spiderwort, spirea, sprude, stonedrop, strawflower, sundrops, sunflowers, sweet pea, sycamore, torch lily, tree peony, tulip tree, variegated privet, verbena, violets, wallflower, walnut, weigela, white spruce, willow, wormwood, yarrow, yellow pine, yew and zinnia.

The Andersons Garden Weed Preventer is a granular formulation for use as a pre-emergence application for control of crabgrass and other annual grasses and certain broadleaved weeds on mineral soils.

Application should be made to a uniform weed-free soil surface within two to three days maximum following cultivation to prevent subsurface germinating weeds from becoming established. Disturbing the soil surface following application will reduce effectiveness.

If there is no rain within three or four days after application, sprinkler irrigation (1/2 to 1 inch of water) should be used for best results. ANY APPLICATION OF TRIPLE X GARDEN WEED PREVENTER SHOULD BE MADE PRIOR TO WEED SEED GERMINATION.

Apply uniformly at the rate of 4 pounds per 1,000 sq. ft. in the early spring before crabgrass and other annual grasses and broadleaved weed seeds germinate. Crabgrass seed germination usually coincides with the time Forsythia blossoms start to fall. One application applied before weed seed germination in early spring usually provides season-long control. Where spring seeding is necessary, there should be a delay of approximately 60 days after the application.

(BACK PANEL)

COMMERCIAL USE ONLY

USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT DURING MIXING/LOADING AND APPLYING, REPAIR AND CLEANING OF MIXING, LOADING, AND APPLICATIONS EQUIPMENT, DISPOSAL OF THE PESTICIDE, AND EARLY REENTRY INTO TREATED AREAS: Wear a long-sleeved shirt and long-legged pants and chemical resistant gloves.

IMPORTANT! If pesticide comes in contact with skin, wash off with soap and water. Always wash hands, face and arms with soap and water before smoking, eating, drinking, or toileting.

AFTER WORK: Before removing gloves, wash them with scap and water. Take off all work clothes and snoes. Enower using scap and water. Do not reuse contaminated slothing. Clothing work during work must be laundered separately from household articles. Clothing that becomes heavily contaminated or usenance must be destroyed apporating to State and local regulations. HEAVILY CONTAMINATEL OF OPENCHED CLOTHING CANNOT BE ACEQUATELY DECONTAMINATED.

Do not enter treated areas for 24 hours after application unless wearing long-sleeved shirt and long pants.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

(BACK PANEL)

COMMERCIAL USE ONLY

WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions For Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and turf conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer.

Manufactured by:

THE ANDERSONS
P.O. BOX 119
MAUMEE, OH 43537

Date Printed: 11/02/94

Date Approved: Supersedes: Label Change:

4 10

5815-35H Page 1 of 4

ACCEPTED
with COMMENTS
in EPA Letter Traced:

FEB 1 1095

(FRONT PANEL)

HOMEOWNER USE CNLY

THE ANDERSONS GARDEN WEED PREVENTER

Under the Franciscoss side. Fundamic, and Long to the Act as amended, for the provide registered under EPA ROL No. 5815-35

ACTIVE INGREDIENT

DFCA (dimethyl tetrachloroterephthalate).... 5.00%

INERT INGREDIENTS 95.00%

TOTAL 100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

of in Eyes: Flush with plenty of Water. Call a physician if irritation persists.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 5815:35

EPA Est. 9198-CH-1

. . .

10 FOUNDS NET WT.

COVERS 2,500 SQ. FT.

5815-35H Page 2 of 4

(BACK PANEL)

HOMEOWNER USE ONLY

THE ANDERSONS GARDEN WEED PREVENTER

KEE, OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

CAUTION

Harards to Humans and Domestic Animals

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or slotning. Wash thoroughly with soap and water after handling. Avoid breathing dust. To not contaminate food and feedstuffs. Do not store hear seed, fertilizer or other herbicides.

ENVIPONMENTAL HAZARDS

Collect and incorporate granules spilled on the soil surface. Do not apply to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is in violation of Federal Law to use this product in a manner inconsistent with its labeling.

DO NOT APPLY IN SUFFOLK COUNTY, LONG ISLAND, NEW YORK.

FOR USE ON: Beans (dry, field, snap), brodcoli, Brussels sprouts, cubbage, cantaloupe, cauliflower, collards, corn (field), corn (sweet), cotton, cress (upland), cucumber, eggplant, garlic, honeydew melon, horseradish, kile, leeks, lettuce (head), lettuce (leaf), mung beans, mustard greens, onions, peppers, potato (whole or cut pieces), Southern beans (blackeyed peas), radish (grown for seed), soybeans, squash, strawberry, sweet potatoes, tomato, turnip (greens), turnip (roots), watermelon, yams (Ipomoea spp.).

Also For Use On The Following Crnamentals and Trees: abelia, African violet, ageratum, aguga, alyssum, arborvitae, ash, aster, athletic field turf, Austrian pine, azalea, baby's breath, barberry, beliflower, birch, bleeding heart, bloodleaf, blue bedder salvia, blue spruce, border forsythia, Boston ivy, boxweed, bugloss, California privet, camellia, candle larkspur, vandy tuft, Cape marigold, chestnit, corysanthemum, singuefoil, coleus, columbine,

• • •

(BACK PANEL)

HOMEOWNER USE ONLY

coneflower, coral bells, coreopsis, cotoneaster, cottonwood, creeping juniper, creeping thyme, cuphea, cypress, dahlias, daisies, deutzia, dogweed, dwarf forsythia, elaeagnus, elm, English ivy, English ivy (ground cover), euonymus, evening primrose, fernleaf, yarrow, feverfew, fir, flowering crabapple, forget-me-not, forsythia, four-o-clock, foxglove, gaillardia, geranium, gladiolus, gold dust, golden marguerite, goldtuft, golf fairways/golf greens, ground-hemlock, gum, hawthorn, heath, holly, honeysuckle, hyrid tea roses, hydrangea, iris, ivy (ground cover), Japanese andromeda, Japanese barberry, Japanese privet, Japaese yew, juniper, lantana, larkspur, lavender cotton, liqustrum, lilac, lilies, loblolly pine, locust, lupines, magnolia, maple, marigold, mentor barberry, mock-orange, morning-glory, mountain-laurel multiflora rose, nasturtium, oak, orpine, pachysandra, pachysandra (ground tover:, paxistima, peonies, petunias, pine, pittosporum, ponderosa pine, poplar, podocarpus, purple coneflower, red pine, redbud, rhododendron, rose, rose moss, Pussian blive, salvia, scapiosa, boarlet sage, Scotch pine, shore pine, snaparagon, 'plierwort, spirea, sprude, stonedrop, strawflower, sundrops, sunflowers, sweet pea, sycamore, toron lily, tree peony, tulip tree, variegated privet, versena, violets, wallflower, walnut, weigela, white oprude, willow, wormwood, yarrow, yellow pine, yew and zinnia.

The Andersons Garden Weed Freventer is a granular formulation for use as a pre-emergence application for control of trabgrass and other annual grasses and certain broadleaved weeds on mineral soils.

Application should be made to a uniform weed-free soil surface within two to three days maximum following cultivation to prevent subsurface germinating weeds from becoming established. Disturbing the soil surface following application will reduce effectiveness.

If there is no rain within three or four days after application, sprinkler irrigation (1/2 to 1 inch of water) should be used for best results. ANY APPLICATION OF TRIPLE X GARDEN WEED PREVENTER SHOULD BE MADE PRIOR TO WEED SEED GERMINATION.

Apply uniformly at the rate of 4 pounds per 1,000 sq. ft. in the early spring before crabgrass and other annual grasses and broadleaved weed seeds germinate. Crabgrass seed germination usually coincides with the 'time Forsythia blossoms start to fall. One application applied before weed seed germination in early spring usually provides season-long control. . Where spring seeding is necessary, there should be a delay of approximately 60 days after the application.

USE ONLY WHEN WEAPING THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT DURING **** MIXING/LOADING AND APPLYING, REPAIR AND CLEANING OF MIXING, LOADING, AND APPLICATIONS EQUIPMENT, DISPOSAL OF THE PESTICIDE, AND EARLY REENTRY INTO ***** TREATED APEAS: Were a long-sleeved shirt and long-legged pants and chemical ***** resistant gloves.

(BACK PANEL)

HOMEOWNER USE ONLY

IMPORTANT' If pesticide comes in contact with skin, wash off with soap and water. Always wash hands, face and arms with soap and water before smoking, eating, drinking, or toileting.

AFTER WORK: Before removing gloves, wash them with soap and water. Take off all work clothes and shoes. Shower using soap and water. Do not reuse contaminated clothing. Clothing worn during work must be laundered separately from household articles. Clothing that becomes heavily contaminated or drenched must be destroyed according to State and local regulations. HEAVILY CONTAMINATED OR DRENCHED CLOTHING CANNOT BE ADEQUATELY DECONTAMINATED.

Do not enter treated areas for 24 hours after application unless wearing long-sleeved shirt and long pants.

STORAGE AND DISPOSAL

To not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and dispard in trash. CONTAINER DISPOSAL: Do not reuse bag. Dispard bag in trash.

WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions For Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OF IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and turf conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer.

Manufactured by:

THE ANDERSONS
P.O. BOX 119
MAUMEE, OH 43537

Date Printed: 11/02/94

Date Approved: Supersedes: Label Change: