(Front Pasel)

Black(R) Leaf Since 1879


Tomato and Vegetable Dust

Easy to use Garden dust

Fungicide-insecticide. Controls listed insects and diseases on tomatoes, as well as other listed vegetables.

KEEP OUT OF REACHEPA Reg. No. 5887-40OF CHILDREN.EPA Est. No. 5887-IL-1C A U T I O NSee additional precautionaryStatements on back panel.EPA Est. No. 5887-IL-1

NET 10 0Z (283 g)



| • | |
|-----------|-----------|
| | • • • |
| | |
| • • • • | • • |
| | |
| | |
| c | • • • |
| | 4.4 |
| | |
| | |
| | |
| | 6 L I I S |
| · · · | |
| 1 I | ۴. |
| < c < 4 § | |
| | |
| L. L. | |
| | |
| | |
| | |
| • • • | • • |
| | |
| | |
| | |
| `.; | |
| | |

•

{

(

(Back panel)

۰., ۱

TOMATO & VEGETABLE GARDEN CUST is a convenient and ready-to-use garden dust; being an effective combination of an insecticide and a fungicide, which will control diseases and insects on vegetables as shown on chart below.

STATEMENTS PRECAUTIONARY HAZARDS TO HUMANS AND DOMESTIC ARIMALS

CAUTION: Harmful if inhaled or swallowed. Do not breathe dust or spray mist. Do not take internally. Avoid contact with skin. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. Food utensils, such as tablescourds, should not be used for food purposes after use with pesticides Keep out of reach of children and animals.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product, or allow it to drift, to blooming crops or weeds if bees are visiting the treatment area. Protection information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TOMATO & VEGETABLE GARDEN DUST may be used in the convenient plastic spray duster, or also as a liquid spray. Best results will be obtained when spraying in early morning or late evening during periods of low temperature and air movement. TOMATO & VEGETABLE GARDEN DUST should be applied to all exposed parts of the plant, including the lower leaf surfaces.

| CROPS | PESTS & DISEASES | AMOUNT | MAY BE |
|---|---|--|--------------------------------------|
| | CONTROLLED | TO USE | APPLIED |
| Potatoes and Tomatoes | PESTS: Lact Bugs, Leafhoppers, Stink Bugs, Tarnished Plant Bugs, Tomato Fruitworms, and Tomato Hornworms, DISEASES: Anthracnose, early and late Blight, Gray Leaf Spot, and Septoria Leaf Spot | 11 tbsp (3 ozl per gal (24 g/L) | Up to 5 days before harvest |
| Dry Beans, Lima Beans, and String Beans | PESTS: Aphids, Armyworms, Bean Leaf Beetles, Japanese Beetles, Mexican Bean Beetles, Stink Bugs, Tarnished Plant Bugs, and Yelvet Bean Caterpillars DISEASES: Downy Kildew | 11 tbsp (3 oz) per gal (74 g/L) | Up to 4 days before harvest |
| Cucumbers, | PESIS: Cucumber Beetles, Flea Beetles, | | Up tu |
| Pumpkins, and | Leafhoopers, Helonworms, Pickleworms, | | 5 days |

| Pumpkins, and Squashes | Leafhopper and Squash | | Pickleworms, | per gal | 5 days before |
|---------------------------|--------------------------|----------------|-----------------|----------|------------------|
| | DISEASES: | Anthracnose an | id Downy Hildew | (24 ġ/L) | harvest |

SPRAYING: Mix 11 tablespoonfuls (3 ounces) in about 1 pint of water until all of the dust is moist or pasty, and then dilute to 1 gallon with water (or mix 24 grams in about 100 mL of water until all of the dust is moist or pasty, and then dilute to 1 liter with water). During the spraying, please continue to shake or agitate. Avoid storing this diluted spray,

TOMATO & VEGETABLE GARDEN DUST should be used as soon as bugs begin to appear. The spray or dusting should be repeated at 5 to 10 day intervals.

FUNGUS CONTROL: To control the listed diseases that can affect your vegetables, a dusting program, using TOMATO & VEGETABLE GARDEN DUST, should corrence as soon as foliage appears; repeat every 1 or 2 weeks.

STORAGE AND DISPOSAL

GENERAL: Do not contaminate water, food, feed, fertilizer, or other pesticides by use, storage, or disposel. If damaged, leaking, or spilled, sweep up and wrap waste, then dispose of as below.

STORAGE: Keep in upright, tightly closed original container in a cool, dry, locked area away from flames, sparks, sunlight, other chemicals, and other sources of ignition, heat, degradation, or contamination.

, DISPOSAL: Do not reuse empty original container. Securely wrap in several · layers of newspaper and discard in trash.

. This container sold by weight and not by volume. If it does not appear full when opened, it is because contents have settled during shipping and handling.

| | | | | 1 | | | |
|---|---|---|---|---|---|-----------------------------|--|
| • | | ٠ | | | ٠ | Manufactured by | |
| 1 | • | | 1 | • | | BLACK LEAF PRODUCTS COMPANY | |
| | | ٠ | | | | CHICAGO, ILLINOIS 60609 | |

0687 .

. .

625

itade in U.S.A.

•••••

. .

÷

(UPC CODE HERE) 76842 00625 1



(

€

۵.