Reg # 5887-51

ł

ACCEPTED

OCT 1 3 1992

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide

EPA Reg. No. 5887-

(Front Panel)

Black Leaf (R)

Warf (R) Pellets

Contains Warfarin

KILLS NORWAY RATS, ROOF RATS AND HOUSE MICE

KEEP OUT OF REACH OF CHILDREN CAUTION See back panel for additional precautionary statements

EPA REG. NO. 5887-51 EPA EST. NO. 70-KY-1 NET WT 1 1b (16 oz) 453 g

(Back Panel)

)

1

WARF is a special mixture of palatable bait materials and an effective rodent-killing ingredient.

WARF must be taken daily over a period of several days. Rats and mice generally do not associate WARF with its cumulative weakening effects and continue to feed on it as long as they are able to do so.

WARF contains an articoagulant chemical that reduces the clotting ability of the blood and upon repeated feedings, causes fatal hemorrhaging in rats and mice, and other warmblooded animals. Use this effective product with care.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin Ki intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat at necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS

This product is toxic to mammals and birds. Do not apply this product to water, swamps, bogs, or potholes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



Black Leaf Warf Pellets (bleaf\wpellrev) Page 2

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

<u>IMPORTANT:</u> Do not expose children, pets, or other nontarget animals to rodenticides. To help prevent accidents:

- 1. Store product not in use in a location out of reach of children and pets.
- 2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamperresistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
- 3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

<u>USE RESTRICTIONS</u> - For control of Norway rats, roof rats, and house mice in and around homes and similar man-made structures. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

<u>SELECTION OF TREATMENT AREAS:</u> Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signed have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

)

RATS: Open bag and apply 4 to 16 oz of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Open bag and apply 1/4 to 1/2 oz. (1-2 level tablespoons) of bait at 8- to 12- foot intervals. Larger placements (up to 2 oz.) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS/MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation,



Black Leaf Warf Pellets (bleaf\wpellrev) Page 3

)

1

limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of intestation is present, establish permanent bait stations and replenish as needed.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets. DISPOSAL: Do not reuse empty container. Securely wrap in newspaper and discard in trash.

DISTRIBUTED BY BLACK LEAF PRODUCTS UPC CODE #76842 00566 A Division of Wilbur-Ellis Company BUCKNER, KENTUCKY 40010 B-992 550 Made in U.S.A.