

CAUTION: Harmful if swallowed.

Avoid breathing of dust or spray. Wear regular long sleeved work clothing. Do not take internally. Skin contact may be harmful. Avoid contact. Wash hands and face before eating. Avoid contamination of food and keep out of reach of children.

This product is highly toxic to pollinating bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

CONTAINER DISPOSAL: Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place.

NOTICE: Payer assumes all risks of use. Storage or handling of this material not in strict accordance with directions given hereafter. Protect from temperatures below 32 F. Keep cover on container to prevent drying.

See also Trade Mark of the Union Carbide Corp. for the active ingredient, Carbaryl. N-methylcarbamate, U.S. Pat. No. 2,405,475. Carbaryl is a registered trademark of Union Carbide Corp.

DISTRIBUTED BY
BLACK LEAF PRODUCTS CO.
ELGIN, ILLINOIS 60120

00-641


Black Leaf

SINCE 1879

For healthier tomatoes and vegetables

ACTIVE INGREDIENTS:	By Weight
Carbaryl (1-naphthyl-N-methylcarbamate)	22.5%
INERT INGREDIENTS	77.5%
Total	100.0%

CAUTION
KEEP OUT OF THE REACH OF CHILDREN
See side panel for other cautions



EPA REG. NO. 5887-108 AA
Net Contents 16 fl. oz. (1 pint)

ACCEPTED
AUG 22 1975
Under the Federal Insecticide, Fungicide, and Plant Disease Act, as amended, in accordance with the requirements of the Federal Food, Drug, and Cosmetic Act, as amended.
EPA Reg. No. 5887-108 AA

EPA EST. NO. 7-1115

DIRECTIONS FOR USE OF BLACK LEAF TOMATO

BLACK LEAF TOMATO & VEGETABLE SPRAY is an extremely potent mixture of material into nearly full spray tank or pre-mixture into spray tank. Add balance of water. Agitate thoroughly cover all plant surfaces.

PRECAUTIONS: Avoid spray drifting to adjoining food crops. Do not use during periods when honey bees are very active. Bees hives well in advance to locate hives at a safe distance. Consult Federal Agricultural Authorities for more information on the use of insecticides, fungicides and miticides, but not with lime.

FOR USE ON	PESTS CONTROLLED
Tomatoes, egg plant, peppers, potatoes	Colorado potato beetle, flea beetle, leafhoppers
Cucumbers, melons, squash, pumpkins	European corn borer, fall armyworm, lace bugs, stink bugs
Peas, beans, peas, beans	Melonworm, pickleworm

5887-108

Black Leaf

DE 1879

EPA EST

DIRECTIONS FOR USE OF BLACK LEAF TOMATO & VEGETABLE SPRAY SHAKE WELL BEFORE USING

BLACK LEAF TOMATO & VEGETABLE SPRAY is an extremely fine dispersion of Sevin in water. Pour recommended amount of material into nearly full spray tank or pre-mix material in another container half full of water and pour mixture into spray tank. Add balance of water. Agitate while mixing and using. Apply as a coarse wet spray to thoroughly cover all plant surfaces.


PRECAUTIONS: Avoid spray drifting to adjoining food or storage crops in bloom. For protection of honey bees, avoid unnecessary use during periods when honey bees are visiting crops. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application. Refer to State or Federal Agricultural Authorities for more information on the control of the pests listed. Sevin is compatible with common insecticides, fungicides and miticides, but not with lime or other highly alkaline materials.

ACCEPT
AUG 22 1975
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended.
 EPA REG. NO. 5887-108

FOR USE ON	PESTS CONTROLLED	SPRAY DIRECTIONS
Tomatoes, egg-plant, peppers, potatoes	Colorado potato beetle, flea beetles, leathoppers	Mix 2 oz. in 3 gal. of water (2 qts. in sufficient water for 1 acre). Spray when insects or damage appear and repeat at 7 to 10 day intervals when necessary.
	European corn borer, fall armyworm, lace bugs, stink bugs	Mix 4 oz. in 3 gal. of water. Spray when insects or damage appear and repeat at 7 to 10 day intervals when necessary.
Cucumbers, pumpkins, squash	Melonworm, pickleworm	Mix 2 oz. in 3 gal. of water. Spray when insects or damage appear and repeat at 7 to 10 day intervals when necessary.

for healthier tomatoes and vegetables

ACTIVE INGREDIENTS: By Weight
 Carbaryl (1-naphthyl-N-methylcarbamate) 22.5%
INERT INGREDIENTS 77.5%
 Total 100.0%



EPA REG. NO. 5887-108 AA
 Net Contents 16 fl. oz. (1 pint)

CAUTION
 KEEP OUT OF THE REACH OF CHILDREN
 See side panel for other cautions

As soon as you see the first signs of black leaf, spray your tomatoes, eggplants, and other vegetables with Black Leaf. You'll see the leaves turn black again. This is the first sign of black leaf. It's caused by a fungus that attacks the leaves of your plants. Black Leaf kills the fungus before it can get to the leaves. So your plants stay healthy and you get a better harvest.

Black Leaf

For healthier tomatoes and vegetables

EPA EST. 7-1175

ACCEPTED

AUG 28 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act.

ACTIVE INGREDIENTS 6.43%
 Carbaryl (1-naphthol) 22.05%
 Methylcarbamate 33.52%
INERT INGREDIENTS 39.99%

CAUTION
 KEEP OUT OF THE REACH OF CHILDREN



EPA REG. NO. 136-1508-A
 EPA REG. NO. 136-1508-A

1k

ACCEPTED

AUG 24 1975

U.S. DEPARTMENT OF AGRICULTURE

FOOD AND NUTRITION ASSISTANCE ACT

HEALTHIER TOMATOES AND VEGETABLES

INGREDIENTS	by weight
1-naphthyl N	
ibamate	22.5%
INGREDIENTS	17.5%
Total	500.0



IN
 T OF THE
 F CHILDREN
 and for other purposes

EPA REG. NO. 300-100-AA
 1-1-1975