

ACCEPTED
MAY 19 1981
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under,
 EPA Reg. No.

Environmental Hazards

Do not allow this material to drift. Avoid contamination of feed and foodstuff.

Physical or Chemical Hazards

Do not use during or immediately prior to periods of extreme high temperatures (90 or higher) as foliage damage may result. Do not apply within 3 weeks of an oil spray or injury may result. Use only in accordance with local recommendations.

Certain crops and other plants are susceptible to injury from Sulfur. Do not use Sulfur on any crop unless local conditions have proven its use to be safe.

Do not breathe dust or mist. Avoid inhaling dust or mist. Do not get it into eyes, nose or throat. Avoid contamination of clothing.

In California do not permit workers to enter the treated area where substantial contact will be made with treated foliage until following indicated days have lapsed after application: Grapes 1, Peaches and Nectarines 1, Citrus 1.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS PRODUCT IS AN AGRICULTURAL FUNGICIDE FOR THE BELOW INDICATED USES:

Storage and Disposal

Prohibitions: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide Disposal: Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

Container Disposal: Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

General: Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

**BEN SUI
 SPRAY**

ACTIVE INGREDIENTS
 Sulfur ..
 INERT INGREDIENTS
 TOTAL.....

(CONTAINS BENTONITE)

Keep Out of Reach of Children

CAUTION:

If in eyes, wash with copious quantities of water minutes. If irritation persists, get medical assistance.

MANUFACTURED FOR
MOYER CHEMICAL COMPANY
 1310 BAYSHORE HIGHWAY P.O. BOX 945 SAN JOSE, CA
 EPA EST. NO. 279-CA-2
EPA REG. NO. 5967-101 AA
NET WEIGHT 50 LBS.

BEN SUL SPRAY

INGREDIENTS	By Weight
Sulfur	81.25%
INGREDIENTS	18.75%
TOTAL	100.00%

(CONTAINS BENTONITE)

Keep Out of Reach of Children

CAUTION:

Eyes, wash with copious quantities of water for 15 minutes. If irritation persists, get medical assistance.

3 81

MANUFACTURED FOR

MOYER CHEMICAL COMPANY

SHORE HIGHWAY P.O. BOX 945 SAN JOSE, CALIFORNIA

EPA EST. NO. 279-CA-2

EPA REG. NO. 5967-101 AA

NET WEIGHT 50 LBS.

31



ALMONDS: Scab, Brown Rot—6 lbs. per 100 gals. Apply in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals.

APPLES, PEARS: Scab, Powdery Mildew—5 lbs. or 4 lbs. plus 1 gal. liquid Lime-Sulfur per 100 gals. Apply as a full coverage spray at pre-bloom, calyx and early cover spray. Under extreme disease conditions, 7 to 8 lbs. may be needed.

GRAPES: Powdery Mildew—5 lbs. per 100 gals. Apply as needed to maintain control. Always follow local recommendations for best results.

NECTARINES, PEACHES: Powdery Mildew, Brown Rot—5 lbs. per 100 gals. Apply at pink bud for Brown Rot control. For Mildew, apply at petal fall and as needed to maintain control until pit hardening. Check local recommendations.

CITRUS (Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangelos, Tangerines): For Citrus Flat Mite, Citrus Rust Mite, Citrus Thrips, Yuma Spider Mites—Apply at least 60 lbs. in 400 to 1500 gals. of water. Apply when pests first appear and repeat as necessary. Do not apply during or preceding high temperatures. Do not apply within two months of an oil application.

STRAWBERRIES: For Mildew, use 5 to 10 lbs. per acre. **By Air:** Apply to 10 to 20 gals. of water. **By Ground:** Apply in 200 to 500 gals. of water.

STRAWBERRIES are sensitive to sulfur, especially during periods of high temperature. Do not apply during hot weather, or when it is anticipated that hot weather will follow application. It is suggested that you contact your local Cooperative Agricultural Extension Service for further advice.

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS OF WATER PER ACRE UNLESS OTHERWISE NOTED:

GROUND APPLICATION— Deciduous Fruit Crops, Grapes and Strawberries—60 to 800 gals. Citrus—40 to 3,000 gals. (On young fruit planting—Use minimum of 15 gals. water per acre.) Where recommended for AIR APPLICATION—Strawberries—10 to 20 gals. Orchard and Vineyard Crops—10 to 20 gals.

32