

## DIRECTIONS

**FOR FOGGING:** When fogging out of doors, use 5 pints per acre of Vapona 1% for 0.05 pounds per acre of actual DDVP. For indoor fogging use 1 gal. of Vapona 1% per 64000 cu. ft. (or 1 pint per 8,000 cu. ft.) for control of flies, mosquitoes, gnats and other small flying insects. Remove or cover aquaria during treatment. When fogging indoors, eliminate all sources of ignition (pilot-lights, fires, open flames) and disconnect electric power. Apply dense fog when wind is calm 1 to 5 seconds per 1000 cu. ft. Before applying, reduce air movement as much as possible by closing doors, windows and other openings. DO NOT contaminate water, feed or foodstuffs, milk or milking utensils. DO NOT use this solution for fogging purposes in areas where animals have received direct application within 8 hours.

The use of 1% Vapona in households, restaurants, warehouses, food processing plants and industrial plants is limited to pest control operators only.

Occupants should vacate premises before indoor treatments and should not reoccupy the premises until it has been ventilated. After application open several windows and doors to create air currents to ensure reduction of Insecticide vapors. Food should be removed and food handling equipment covered with wrapping paper during application, or washed after application is completed. During fogging or mist spraying indoors, use mask or respirator of a type passed by the U.S. Department of Agriculture for DDVP protection. DO NOT FOG INTO OR NEAR AN OPEN FLAME.

**WARNING:** May be fatal if swallowed, inhaled, or absorbed through the skin or eyes. Wash thoroughly with soap and water after handling and before eating or smoking. NO NOT contaminate feed, water, or foodstuffs. Keep out of reach of children. If swallowed, induce vomiting immediately and get medical attention. Do not get in eyes, on skin, or on clothing. Rapidly absorbed through skin or eyes. Do not breath vapor spray mist. In case of contact, immediately remove contaminated clothing and flush skin and eyes with plenty of water; for eyes get medical attention. Wear clean rubber gloves, protective clothing and goggles. Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for protection against DDVP. Wash contaminated clothing with soap and hot water before reuse. Do not reuse shoes or boots until free of all contamination. ATROPINE is antidotal. Do not contaminate any body of water by the cleaning of equipment or disposal of wastes and containers. Toxic to aquatic organisms and fish.

Net Contents 1 Gallon

VAPONA



INSEC-TI-FOG

## FOR FOGGING OR SPRAYING

For use in Foggers (against flies, mosquitoes, gnats), Mist Sprayers and Hand Sprayers. Ready to use coarse spray solution for control of Fleas and Ticks in Outdoor Areas around Patios, Warehouses, Restaurants, Food Processing Plants, Horse Barns, Dairy Barns, Shelter and Milk Sheds; and for use as a Fog or Mist Solution in Outdoor Areas for flies, mosquitoes and gnats. Use in barns to contact Face flies, Stable flies, House flies and Horn flies with the Spray Mist. The use of Vapona 1% in households, restaurants, warehouses, food processing plants and industrial plants is limited to pest control operators.

### 100% ACTIVE INGREDIENTS

2, 2-Dichlorovinyl Dimethyl Phosphate  
Related Compounds  
Mineral Oil

### BY WEIGHT

0.92%\*  
0.08%\*  
99.00%

\*Equivalent to 1.00% w Vapona Insecticide  
RVapona Insecticide T. M. Shell Chemical Co

### WARNING: Combustible Mixture

Keep Out of Reach of Children

SEE FRONT PANEL FOR ADDITIONAL WARNINGS

**CUSTOM FARM SERVICES, INC.**

ROWLAND-GORDON FARM CHEMICALS DIVISION

BOX 1449

PLAINVIEW, TEXAS 79072

U.S.D.A. REG. 60742

ACCEPTED

APR 10 1967

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND FERTILIZER ACT  
FOR ECONOMIC PEST CONTROL REGISTERED  
UNDER F.P.C. 60742 SUBJECT  
TO ATTACHED COMMENTS.

## DIRECTIONS

**CATTLE** (Beef and Dairy) To control Hornflies, Ho Flies, Face Flies, apply 1 to 2 fluid ounces of Vapona 1 as a mist spray daily with hand or automatic sprayer, as to wet the hair. DO NOT apply in excess of 2 fl animal per day.

**ANIMAL BUILDINGS** (Horse Barns, Dairy Barns, Shelter Houses, Milk Sheds). Outdoor areas. To control Flies, Ticks, Roaches, Fleas, Ants, Sowbugs, Brown Dog Ticks, Moths, and other small flying insects. Cockroaches, An Silverfish. Apply Vapona Mist as a coarse wet spray a pt. of 1.0% solution per 1,000 sq. ft. for mill sheds, st barns, loafing sheds, pig pens, and outdoor areas; treat ways, feed storage rooms, alleyways and window sills taminate milk or milking utensils, water, food or foodst of Fly Maggots in Maggot breeding areas apply at the pts. per 100 sq ft. In poultry houses apply to manure outside of penned enclosures and on floor of feed

**OUTDOOR USE** (For control of the above listed insect **DOOR-LIVING AREAS OF HOMES, BACK YARDS, P GROUNDS, OPEN-AIR THEATERS, OUTDOOR LATRINE STANDS, PARKING AREAS; OUTDOOR AREAS OF THEATERS, FOOD PROCESSING PLANTS, INDUSTRIAL HOUSES, SERVICE STATIONS, LOADING DOCKS; O BAGE COLLECTION AND DISPOSAL AREAS.** Use a spray. When fogging around the above, to kill flies, A gnats use Vapona 1% at the rate of 5 pints of V 0.05 lbs. per acre of actual DDVP. In food process Vapona 1% as a space spray only when plant is not Foods should be removed or covered during treatr processing surfaces should be covered or thoroughly using.

In treating Dog Kennel's apply to outside dog runways and ledges. Use Vapona mist as a coarse wet spray at 1 per 1,000 sq. ft. for control of flies, mosquitoes, gnats, flying insects. For control of fleas and brown dog ticks i to 1 qt. of Vapona Mist per 1,000 sq. ft. Treat infest boards, cracks, walls, doors. DO NOT allow children until dry.

For control of fly maggots in maggot breeding areas, a of 1 to 2 pts. per 100 sq. ft.

**WARNING: DO NOT TREAT PETS WITH VAPONA 1% NON-WARRANTY**

Notice to buyer: Seller makes no warranty, of any kin implied, concerning the use of this product. Buyer assume or handling whether in accordance with directions or not.

Contents 1 Gallon

ACCEPTED

APR 10 1967

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC PESTICIDE REGISTRATION  
REGISTERED UNDER NO. 6074-2 SUBJECT  
TO ATTACHED COMMENTS.



INSEC-TI-FOG

### FOR FOGGING OR SPRAYING

Against flies, mosquitoes, gnats), Mist Sprayers Ready to use coarse spray solution for ticks in Outdoor Areas around Patios, Warehouses, Food Processing Plants, Horse Barns, Dairy Sheds; and for use as a Fog or Mist Solution for flies, mosquitoes and gnats. Use in barns, Stable flies, House flies and Horn flies. Use of Vapona 1% in households, restaurants, food processing plants and industrial plants is limited to pest control operators.

INGREDIENTS	BY WEIGHT
2, 2-Dichlorovinyl Dimethyl Phosphate	0.92%*
Related Compounds	0.08%*
Mineral Oil	99.00%

Vapona Insecticide  
T. M. Shell Chemical Co

**G: Combustible Mixture  
Out of Reach of Children  
SEE BACK PANEL FOR ADDITIONAL WARNINGS  
CUSTOM FARM SERVICES, INC.**

ROWLAND-GORDON FARM CHEMICALS DIVISION  
BOX 1449  
PLAINVIEW, TEXAS 79072

USDA Reg 6074-2

### DIRECTIONS

**CATTLE** (Beef and Dairy) To control Hornflies, Houseflies, Stable Flies, Face Flies, apply 1 to 2 fluid ounces of Vapona Mist per animal as a mist spray daily with hand or automatic sprayer. Apply spray so as to wet the hair. DO NOT apply in excess of 2 fluid ounces per animal per day.

**ANIMAL BUILDINGS** (Horse Barns, Dairy Barns, Shelter Sheds, Poultry Houses, Milk Sheds). Outdoor areas. To control Flies, Gnats, Mosquitoes, Roaches, Fleas, Ants, Sowbugs, Brown Dog Ticks, Wasps, Flying Moths, and other small flying insects. Cockroaches, Ants, Spiders and Silverfish. Apply Vapona Mist as a coarse wet spray at the rate of 1 pt. of 1.0% solution per 1,000 sq. ft. for milking sheds, stables, livestock barns, loafing sheds, pig pens, and outdoor areas; treat around doorways, feed storage rooms, alleyways and window sills. DO NOT contaminate milk or milking utensils, water, food or foodstuffs. For control of Fly Maggots in Maggot breeding areas apply at the rate of 1 to 2 pts. per 100 sq ft. In poultry houses apply to manure, window sills outside of penned enclosures, and on floor of feed storage rooms.

**OUTDOOR USE** (For control of the above listed insects). Treat OUTDOOR LIVING AREAS OF HOMES, BACK YARDS, PATIOS, PICNIC GROUNDS, OPEN-AIR THEATERS, OUTDOOR LATRINES, ICE CREAM STANDS, PARKING AREAS, OUTDOOR AREAS OF RESTAURANTS, THEATERS, FOOD PROCESSING PLANTS, INDUSTRIAL PLANTS, WAREHOUSES, SERVICE STATIONS, LOADING DOCKS; OUTDOOR GARBAGE COLLECTION AND DISPOSAL AREAS. Use a coarse wetting spray. When fogging around the above, to kill flies, Mosquitoes, and gnats use Vapona 1% at the rate of 5 pints of Vapona 1% for 0.05 lbs. per acre of actual DDVP. In food processing plants use Vapona 1% as a space spray only when plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using.

In treating Dog Kennels apply to outside dog runways, window sills, and ledges. Use Vapona mist as a coarse wet spray at the rate of 1 pt. per 1,000 sq. ft. for control of flies, mosquitoes, gnats and other small flying insects. For control of fleas and brown dog ticks increase dosage to 1 qt. of Vapona Mist per 1,000 sq. ft. Treat infested areas, baseboards, cracks, walls, doors. DO NOT allow children in treated area until dry.

For control of fly maggots in maggot breeding areas, apply at the rate of 1 to 2 pts. per 100 sq ft.

**WARNING: DO NOT TREAT PETS WITH VAPONA 1%.**

#### NON-WARRANTY

Notice to buyer: Seller makes no warranty, of any kind, expressed or implied, concerning the use of this product. Buyer assumes all risk of use or handling whether in accordance with directions or not.

Net Contents 1 Gallon

VAPONA



INSEC-TI-FOG

### FOR FOGGING OR SPRAYING

For use in Foggers (against flies, mosquitoes, gnats), Mist Sprayers and Hand Sprayers. Ready to use coarse spray solution for control of Fleas and Ticks in Outdoor Areas around Patios, Warehouses, Restaurants, Food Processing Plants, Horse Barns, Dairy Barns, Shelter and Milk Sheds; and for use as a Fog or Mist Solution in Outdoor Areas for flies, mosquitoes and gnats. Use in barns to contact Face flies, Stable flies, House flies and Horn flies with the Spray Mist. The use of Vapona 1% in households, restaurants, warehouses, food processing plants and industrial plants is limited to pest control operators.

100% ACTIVE INGREDIENTS	BY WEIGHT
2, 2-Dichlorovinyl Dimethyl Phosphate	0.92%*
Related Compounds	0.08%*
Mineral Oil	99.00%

\*Equivalent to 100% w Vapona Insecticide  
T. M. Shell Chemical Co

**WARNING: Combustible Mixture  
KEEP OUT OF REACH OF CHILDREN  
SEE BACK PANEL FOR ADDITIONAL WARNINGS  
CUSTOM FARM SERVICES, INC.**

ROWLAND-GORDON FARM CHEMICALS DIVISION  
BOX 1449  
PLAINVIEW, TEXAS 79072

USDA Reg 6074-2