


CAUTION: KEEP OUT OF THE REACH OF CHILDREN



THORO



SANITIZING SOLUTION

ACTIVE INGREDIENT: N-Alkyl (C₁₄, C₁₂, C₁₆) Dimethyl Benzyl Ammonium Chlorides10.25 %

INERT INGREDIENT: Water89.75 %

AS A SANITIZER FOR RESTAURANTS, BARS, CAFES, SODA FOUNTAINS, ETC.

For sanitizing glasses, dishes and silverware, immerse for 30 seconds in solution of 1 ounce THORO to 4 gallons of water in final rinse tank of dishwashing procedures. This product may be employed in hot or cold water.

Eating and drinking utensils properly washed and submerged in product solutions, prepared as directed, will be sanitized in accordance with health department requirements. One ounce in four gallons of water will supply 200 p.p.m. of active ingredients.

GENERAL DISINFECTANT

For the disinfection of inanimate objects and surfaces such as floors, walls, wood or metal furniture, etc., use one ounce of product to two gallons of water, hot or cold. All surfaces and objects should be cleaned thoroughly with a good detergent prior to disinfection.

This product has a phenol co-efficient of 50 against *S. typhosa* and 75 against *M. pyogenes var. aureus* at 20°C.

"This product is effective at 200 p.p.m. quaternary strength in waters up to 550 p.p.m. (32 grains per gallon) hardness calculated as CaCO₃ when tested by the Chambers' modification of the Weber-Black Technique as outlined by the U.S. Public Health Service."

THORO CHEMICAL COMPANY USDA Reg. No.

14831 BESSEMER STREET

VAN NUYS, CALIFORNIA

6255-1

Telephones: State 5-5751 TRIANGLE 3-2772

Caution Avoid Contamination Of Food

6255-1

ACCEPTED
Jan 19 1968
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. SUBJECT
TO ATTACHED COMMENTS.