

**CAUTIONARY STATEMENTS
TO HUMANS AND
DOMESTIC ANIMALS**

Keep away from humans, children, and pets. If swallowed, this product may reduce the clotting ability of the blood, causing internal or external bleeding.

PHYSICIAN — If ingested, give 10 mg. of Vitamin K₁ intramuscularly or 10 mg. orally in bishydroxycoumarin form as necessary based on prothrombin times.

ENVIRONMENTAL HAZARD — Do not apply this product directly to water, fish, or to intertidal areas.

USE AND DISPOSAL

CONTAMINATION OF WATER, FOOD, OR DRUGS OR DISPOSAL. — Wastes resulting from the use of this product may be disposed of in accordance with approved waste disposal procedures.

CONTAINER DISPOSAL — Empty container: Securely wrap in plastic and dispose in trash.

Manufacturer assumes all risks of use. The user is responsible for the proper application of this material not in accordance with directions given on the label.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to roosting livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unplaced bait as specified on this label.

USE RESTRICTIONS - For control of Norway rats, roof rats, and house mice in and around homes, industrial buildings, and similar man-made structures. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS - Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawing openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much food as possible.

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 oz. of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 15 days.

MICE: Apply 1/4 to 1/2 oz. (1-2 level tablespoons) of bait at 8- to 12-foot intervals. Larger placements (up to 2 oz.) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS/ MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations or replenish as needed.

ACCEPTED
JUN 6 1995
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 6409-1