

ACCEPTED

DIRECTIONS:

RAT CONTROL: 4 ozs. of bait per station should be the minimum amount used. Place bait in locations not accessible to children, pets, domestic animals, or in tamper-proof bait boxes. An uninterrupted supply of bait should be maintained for 10 to 14 days.

MOUSE CONTROL: 1/4 to 1/2 oz. of bait should be placed at 8-12 foot intervals. Place bait in locations not accessible to children, pets, domestic animals, or in tamper-proof bait boxes. An uninterrupted supply of bait should be maintained for at least 15 days.

Place bait in protected areas, corners, along walls, under eaves, where gnawing or damage is seen, tracks, trails, smears or droppings. Replace spoiled or contaminated bait promptly. Fill stations until feeding stops. Keep stations filled to control re-infestation.

**THRIFTY ANTI-COAGULANT
RAT AND MOUSE BAIT**

CONTAINS FUMARIN

ACTIVE INGREDIENTS:	
3 Alpha-acetonylfurfuryl	4 hydroxycoumarin 0.025
INERT INGREDIENTS	
TOTAL	

NET WEIGHT, 50 LBS.



Four or five pounds of bait should be enough to control rats on the average infested farm. For heavy infestations, several pounds may be necessary in each building. On premises subject to reinfestations, keep bait continuously available.

CAUTION:

Keep Out of the Reach of Children
If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of Vitamin K combined with blood transfusions may be indicated, as in the case of hemorrhage caused by overdoses of bishydroxycoumarin (Dicumarol). Protect baits from children, pets and livestock by bait boxes or other suitable means.

W. H. MORTON CO. ATLANTA, GEORGIA

USDA Reg. No. 6561

(See Side Panel for Directions)