

TIDE 10-40 SEVIN* - SULFUR

AGRICULTURAL INSECTICIDE

ACTIVE INGREDIENTS:

Carbaryl (1-Nphtthyl N-methylcarbamate) 10%

Sulfur 40%

INERT INGREDIENTS:

50%

100%

*Trade-Mark of UCC for 1-naphthyl N-methylcarbamate

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to us during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

OBSERVE THE FOLLOWING USE PRECAUTIONS

Sulfur will cause severe fruit and leaf injury to sulfur-sensitive crops.

Do not apply to apricots, comice pears, cranberries, cucurbits (cucumber, cantaloupes, melons, squash), walnuts, or to other sensitive plants.

Sulfur may burn foliage and fruit during periods of high temperature and under certain other climatic conditions. Do not apply sulfur to sensitive crops when shade temperatures exceed or are likely to exceed 90° F.

Do not use sulfur on any crop unless local use has proved safe.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified.

Consult local State Agricultural Extension Service authorities for additional information, as the timing, number, and rate of applications needed will vary with local conditions.

DIRECTIONS FOR USE

Apply when air is calm, using ground equipment, aircraft, or hand duster. Make thorough coverage of plants with a uniform deposit of dust. Apply at the first sign of infestation and repeat at regular intervals when needed to maintain control.

RECOMMENDATIONS

This dust is formulated for the control of the following insects on COTTON.

For control of early-season Thrips and Cotton Fleahopper:

Apply 8 to 15 lbs. of Sevin-Sulfur 10-40 Dust per acre when insects first appear.

For control of Mid and Late Season Boll Weevil, Bollworm, Cotton Leafworm, Lygus Bugs, and Cotton Leaf Perforator.

Apply 15 to 25 lbs. of Sevin-Sulfur 10-40 Dust per acre at 5-7 day intervals when insects first appear.

For control of Pink Bollworms:

Apply 22-25 lbs. of Sevin-Sulfur 10-40 Dust per acre at 5-7 days intervals.

Sulfur Fineness: 99% Passing No. 325 USS Sieve

BURN OR BURY BAG — DO NOT REUSE — STAY OUT OF SMOKE OR FUMES

A Product of

TIDE PRODUCTS, INC.
EDINBURG, TEXAS

CAUTION

KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

- Harmful if inhaled or swallowed.
- Do not take internally.
- Do not breathe dust
- Avoid prolonged or repeated breathing of dust
- Skin contact may be harmful. Avoid unnecessary contact.
- Wash hands before eating. Take shower or bath after work.
- Wear regular long sleeved work clothing. Change to clean clothing daily.
- Note for Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

ACCEPTED
6/27/68
UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE AND RODENTICIDE ACT
FOR LISTING IN PESTICIDE REGISTER
ED UNDER NO. 100-1-100-100
NO ATTACHED COMMENTS

NOTICE: SELLER WARRANTS THAT THE PRODUCT CONFORMS TO ITS CHEMICAL DESCRIPTION AND IS REASONABLY FIT FOR THE PURPOSE STATED ON THE LABEL WHEN USED IN ACCORDANCE WITH DIRECTIONS UNDER NORMAL CONDITIONS OF USE, BUT NEITHER THIS WARRANTY NOR ANY OTHER WARRANTY OR MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, EXPRESS OR IMPLIED, EXTENDS TO THE USE OF THIS PRODUCT CONTRARY TO LABEL INSTRUCTIONS, OR UNDER ABNORMAL CONDITIONS, OR UNDER CONDITIONS NOT REASONABLY FORSEEABLE TO SELLER, AND BUYER ASSUMES THE RISK OF ANY SUCH USE.