



TIDE ZINEB 6.5 DUST

AN AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENTS:

Zineb (Zinc Ethylene Bisdithiocarbamate) 6.5%
(Zinc expressed as metallic 1.42%)

INERT INGREDIENTS

93.5%

100.0%

CAUTION

Keep Out of Reach of Children

HARMFUL IF SWALLOWED.

AVOID BREATHING DUST. AVOID CONTACT WITH EYES, SKIN AND CLOTHING.

WASH THOROUGHLY AFTER HANDLING.

DO NOT CONTAMINATE FOOD AND FEED.

OBSERVE THE FOLLOWING USE PRECAUTIONS

Observe use limitations given on the label for specific crops.

In order that pesticidal residues on food and forage crops will not exceed tolerance established by the Federal Food and Drug Administration, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or to adjoining food, fiber or pasture crops. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others. Consult state agricultural experiment stations or state agricultural extension service for additional information, as the timing, number and rate of application, need it will vary with local conditions.

DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground equipment or aircraft. Apply at the first sign of infection and repeat at 5 to 10-day intervals, depending on severity of infection, but observe use limits given for specific crops. Apply at the rate of 30 to 40 lb. per acre (except when stated differently) depending on plant size.

EPA Reg. No. 6735-192

RECOMMENDATIONS

VEGETABLES

BEANS: Anthracnose, Downy Mildew, Rust -- Do not apply within 7 days of harvest on Lima, snap or string beans.

CABBAGE, CAULIFLOWER: Alternaria Leaf Spot, Cercospora, Downy Mildew -- Do not apply within 7 days of harvest.

CELERY: Early and Late Blight -- Remove residues by washing and stripping. Do not apply within 14 days of harvest.

CUCURBITS: (Cantaloupe, Cucumbers, Squash, Melons): Angular Leaf Spot, Alternaria Blight, Anthracnose, Downy Mildew, Cucurbit Stem Blight. Do not apply within 5 days of harvest.

EGGPLANT, PEPPERS: Anthracnose and Cercospora on peppers; Alternaria, Anthracnose, and Phomopsis on eggplant.

LETTUCE: Downy Mildew -- Do not apply within 10 days of harvest.

POTATOES: Anthracnose, Early and Late Blight, Septoria Leaf Spot, Stemphylium. Do not apply within 5 days of harvest.

CORN: Helminthosporium Leaf Blight -- Do not feed treated foliage to dairy animals or animals being finished for slaughter. Do not apply within 40 days of harvest.

FIELD CROPS:

PEANUTS: Cercospora Leaf Spot -- Apply 20 lbs. per acre. Begin treatment when disease is first noted and continue applications at 1 to 2 weeks intervals depending upon severity of the disease. Do not use treated peanut hay for feed.

PLANT BEDS

TOBACCO: Blue Mold -- In plant beds, use 6 to 11½ lbs. per 100 square yards. Begin when plants are the size of a dime or when disease is first reported in the area. Repeat at 3 to 4 day intervals until transplanting time.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty or merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

50 LBS. NET

GUS-4-77-500-14421

BURN EMPTY BAG -- KEEP OUT OF SMOKE DO NOT REUSE

A PRODUCT OF

TIDE PRODUCTS, INC.

EDINBURG, TEXAS 78539