



TIDE MALATHION 5 DUST

AGRICULTURAL INSECTICIDE
(Contains 5% Premium Grade Malathion)

Active Ingredient:	
Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate)	5%
Inert Ingredients	95%
	100%

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED

AVOID BREATHING DUST.

AVOID CONTACT WITH SKIN.

WASH THOROUGHLY AFTER USING.

AVOID CONTAMINATION OF FEED AND FOOD.

NOTE TO PHYSICIANS: Malathion, upon use, may cause cholinesterase inhibition. Atropine is antidotal.

OBSERVE THE FOLLOWING USE PRECAUTIONS

Observe use limitations given on the label for specific crops.

In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. **The grower is responsible** for residues on his crops as well as for damage caused by drift from his property to that of others.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information as the timing of applications will vary with local conditions.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground equipment or aircraft. Apply at the first sign of infestation, and repeat at regular intervals when needed to maintain control, but observe use limits given for specified crops.

Maximum permissible rate per acre, expressed as Malathion 5 Dust, is given in parentheses () after each crop.

BURN OR BURY EMPTY BAG -- DO NOT REUSE -- STAY OUT OF SMOKE OR FUMES

E.P.A. REG. NO. 6735-201

A PRODUCT OF
TIDE PRODUCTS, INC.
EDINBURG, TEXAS 78539

RECOMMENDATIONS

ALSO READ USE PRECAUTIONS & DIRECTIONS FOR USE

ALFALFA Alfalfa weevil, larvae, army worm, aphids, potato leafhopper, spider mites, cutworms and velvet beetles. Apply 30 lbs. per acre. Apply to alfalfa in bloom only in the evening or early morning when bees are not working in the field or are not landing on outside of bays. (30 lbs.)

Note May be applied on day of harvest or grazing.

BEANS Aphids, cucumber beetle, Japanese beetle, Mexican bean beetle, midribbed beetle, pea leaf weevil, potato leafhopper, spider mites. Apply 30 lbs. per acre. Make 2 or more applications as needed. Do not apply within 24 hours of harvest. (35 lbs.)

CARROTS, RADISHES Aphids, leafhoppers. Apply 30 lbs. per acre. Do not apply within 7 days of harvest. (40 lbs.)

CLOVER Aphids, spider mite. Apply 25-30 lbs. per acre. Do not apply to clover in bloom. May be applied on day of harvest or grazing. (30 lbs.)

COLE CROPS Broccoli, Brussels sprouts, cabbage, cauliflower, kale, mustard and turnips. Aphids, cabbage looper, flea beetle, imported cabbage worm. Apply 20-30 lbs. per acre. Make full coverage applications repeated as necessary. Do not apply within 7 days of harvest of Brussels sprouts, cabbage, cauliflower, kale and mustard 3 days on broccoli and turnips. (50 lbs.)

CUCUMBERS, SQUASH AND MELONS Aphids, pickleworms and spider mites. Apply 30 lbs. per acre. For cucumber beetles and squash vine borers use 35 lbs. per acre. Apply weekly to stems and vines at base of plants to control squash vine borers. CAUTION: Do not apply unless plants are dry. Make no application within 1 day of picking. (45 lbs.)

LETTUCE Aphids, cabbage looper, leafhoppers, mites. Apply 25-35 lbs. per acre. Do not apply to leaf lettuce within 14 days of harvest, 7 days on head lettuce. (60 lbs.)

ONIONS Onion thrips, onion maggot. Apply 20-25 lbs. per acre. For maggot control apply when maggots appear and repeat every 4 days. Do not apply within 3 days of harvest on green onions. (40 lbs.)

PASTURE & RANGE GRASS, GRASS & GRASS HAY Aphids, leafhoppers. Apply 30-40 lbs. per acre. Armyworms on grass and grass hay. Apply 30-40 lbs. May be applied on day of harvest or grazing. (40 lbs.)

PEANUTS Potato leafhopper. Apply 15-20 lbs. per acre. Do not apply within 30 days of harvest. (20 lbs.)

PEAS Pea aphid and pea weevil. Apply 20-25 lbs. per acre. Make 2 or more applications within 1 day of harvest. (50 lbs.)

POTATOES Flea beetle, Colorado potato beetle, Colorado potato beetle, Colorado potato beetle. Apply 20-30 lbs. per acre. (30 lbs.)

SOYBEANS Mexican bean beetle. Apply 30-40 lbs. per acre. Do not apply within 7 days of harvest. (40 lbs.)

SPINACH Aphid. Apply 20-30 lbs. per acre. Do not apply within 7 days of harvest. (40 lbs.)

TOMATOES Aphid, flea beetle, spider mites. Apply 30 to 40 lbs. per acre. Make no application within 24 hours of harvest. (40 lbs.)

NOTICE: SELLER WARRANTS THAT THE PRODUCT CONFORMS TO ITS CHEMICAL DESCRIPTION AND IS REASONABLY FIT FOR THE PURPOSE STATED ON THE LABEL WHEN USED IN ACCORDANCE WITH DIRECTIONS UNDER NORMAL CONDITIONS OF USE, BUT NEITHER THIS WARRANTY NOR ANY OTHER WARRANTY OR MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, EXPRESS OR IMPLIED, EXTENDS TO THE USE OF THIS PRODUCT CONTRARY TO LABEL INSTRUCTIONS OR UNDER ABNORMAL CONDITIONS OR UNDER CONDITIONS NOT REASONABLY FORESEEABLE TO SELLER AND BUYER ASSUMES THE RISK OF ANY SUCH USE.

GPS 10-11M-1010

ACCEPTED WITH COMMENTS