


TIDE CAPTAN 7.5 DUST

AGRICULTURAL FUNGICIDE



ACTIVE INGREDIENT:
CAPTAN: N-(trichloromethylthio)-4-cyclohexene-1,2-dicarboximide 7.5%
INERT INGREDIENTS: 92.5%
100.0%

RECOMMENDATIONS

GRAPES: Mold Control on Fruit on Drying Trays in the Field (Penicillium spp., Aspergillus spp., Botrytis cinerea, Rhizopus spp.) - Apply 13 lbs. per acre. Apply to fruit on trays for protection of fruit and prevention of above molds. Apply to give even coverage. During damp weather make a second application after trays are turned. Apply this dust by hand or ground dusters only. Do not apply within 5 days of a pre-harvest application of Captan.

GRAPES: Botrytis Rot (Bunch Rot) - 50 to 65 lbs. per acre. Make 2 applications before bloom and 1 application just after bloom. Repeat periodically up to 3 cover applications before bunches close.

PEACH, NECTARINES: Brown Rot, Peach Scab - 50 to 65 lbs. per acre. Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make pre-harvest application where necessary, but do not apply within one day of harvest on peaches.

STRAWBERRIES: Botrytis Rot - 40 to 50 lbs. per acre beginning as soon as new growth starts. Repeat at about weekly intervals, up to harvest. Apply after each picking if conditions favor fruit rot.

BEANS, PEAS, Soil Treatment - Damp-off (Beans only), Fungus Root Rot - Work into top 3 or 4 inches of soil before planting. Apply 60 to 75 lbs. per acre.

CELERY: (Plant bed only) - Damp-off, Pink Rot - Apply 50 to 60 lbs. per acre when all seeds have germinated. Repeat if necessary. Repeat at 5 to 10-day intervals until danger of infection is past.

CUCURBITS (Cantaloupe, Cucumber, Melon, Squash): Angular Leaf Spot, Anthracnose, Downey Mildew - Start application when runners begin to form or at the first sign of disease. Apply at rate of 40 to 60 lbs. per acre on young plants increasing to 40 to 50 lbs. per acre on mature plants.

POTATOES: Early and Late Blight - Make first application when plants are 2 to 4 inches high. Repeat at 7 to 10 day intervals as long as disease threatens. Apply 45 to 55 lbs. per acre on young plants. Use 65 lbs. per acre on mature plants.

POTATO SEED PIECE TREATMENT: Damp-off, Seed Rot Fungi, Brown Eye Disease, Verticillium Wilt - Use at rate of 15 ozs. per bushel cut seed pieces (16 ozs. or 1 lb. to 100 lbs. of cut seed pieces). Apply in convenient container or by duster attachment over belt. Coat each seed piece evenly with a light film of dust. Treat pieces within 6 hours after cutting. If planting is delayed, store treated seed pieces in open crates two days after bagging.

TOMATOES: Anthracnose, Early and Late Blight, Gray Leaf Spot, Septoria Leaf Spot - Apply 30 to 45 lbs. per acre at weekly intervals on young plants, 55 to 65 lbs. per acre on mature plants.

SOIL AND GREENHOUSE BENCH TREATMENT - 2 1/4 lbs. per 1,000 sq. ft. (80 lbs. per acre). Work into upper 3 or 4 inches of soil before planting. For damp-off and root rot disease of seedlings and transplants of roses, other shrubs, flowers, lawn seedbeds and vegetables such as beans, peas, cabbage, eggplant, pepper and tomato.

ORNAMENTALS: Rose, Black Spot; Carnation, Rust, Leaf Spot; Chrysanthemum, Flower Blight - Apply thoroughly prior to or at first signs of disease. Repeat applications at 7 to 10 day intervals. Shorten interval during periods of frequent rain and heavy dews.

LAWNS, TURF: Brown Patch, Damp-off, Leaf Spot, Root Rot Fungus - Apply 2 to 3 lbs. per 1,000 sq. ft. Repeat at 7 to 10 day intervals as needed. Melting-out - Apply 2 to 3 lbs. per 1,000 sq. ft., beginning when grass resumes growth in the Spring (March-April). Repeat at 7 to 14 day intervals as long as cool, wet weather prevails. Do not graze treated areas or feed clippings to livestock.

NOTICE: SELLER WARRANTS THAT THE PRODUCT CONFORMS TO ITS CHEMICAL DESCRIPTION AND IS REASONABLY FIT FOR THE PURPOSE STATED ON THE LABEL WHEN USED IN ACCORDANCE WITH DIRECTIONS UNDER NORMAL CONDITIONS OF USE. BUT NEITHER THIS WARRANTY NOR ANY OTHER WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, EXPRESS OR IMPLIED, EXTENDS TO THE USE OF THIS PRODUCT CONTRARY TO LABEL INSTRUCTIONS, OR UNDER ABNORMAL CONDITIONS OR UNDER CONDITIONS NOT REASONABLY FORESEEABLE TO SELLER. AND BUYER ASSUMES THE RISK OF ANY SUCH USE.

CAUTION

Keep Out of Reach of Children

Feb. 25, 1974
6785-213

HARMFUL IF TAKEN INTERNALLY

AVOID INHALING DUST.

AVOID CONTACT WITH SKIN.

AVOID CONTAMINATION OF FEED AND FOOD.

STORE IN A COOL DRY PLACE; PROTECT FROM EXCESSIVE HEAT.

USE PRECAUTIONS

The Federal Food And Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information, as the timing, number, and rate of applications needed will vary with local conditions.

DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground equipment or aircraft. Apply at the first sign of infection and repeat at 5-10 day intervals as needed to maintain control.

BURN EMPTY BAG - KEEP OUT OF SMOKE

E.P.A. Reg. No. 6735-203

A PRODUCT OF
TIDE PRODUCTS, INC.
EDINBURG, TEXAS 78539

GPS 1-74 1M- (7804) B-74