

OCTAGON PROCESS INC.

596 RIVER ROAD

EDGEWATER, NEW JERSEY

RODENTICIDE, BAIT, ANTICOAGULANT, PIVAL 0.025%

OCTAGON PIVAL 0.025%

FEDERAL SPECIFICATION O-R-500B

NET WEIGHT - 5 LBS.

ACTIVE INGREDIENTS 2-pivalyl-1, 3-indandione 0.025%

INERT INGREDIENTS 99.975%

Liquid petrolatum 4.800%

Sucrose 4.800%

Dye 0.050%

Breakfast Rolled Oats 90.325%

ACCEPTED WITH COMMENTS

CAUTION: KEEP OUT OF REACH OF CHILDREN

DIRECTIONS:

BAIT CONTROL: Place 4 to 8 ounces of Rodenticide, Bait, Anticoagulant, Bait stations. Place so as to protect children, pets, and livestock in locations where rats or mice feed. Replenish baits as necessary, and maintain stations for not less than 15 days. Baiting may be continued indefinitely, and increased time of exposure of baits may result in better control of rodents. If bait molds, decomposes, or becomes sour, replace with fresh material.

SPECIAL DIRECTIONS FOR MOUSE CONTROL:

a) Tablespoon amounts (14 to 12 ounce) of bait should be placed at 8-12 foot intervals.

b) An uninterrupted supply of bait should be maintained for at least 15 days.

CAUTION: Keep away from humans, domestic animals, and pets. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated in the case of hemorrhage caused by overdose of bis-hydroxycoumatin.

Baits should be placed in areas inaccessible to children, pets, wildlife, and domestic animals, or in tamper-proof containers.

Do not use in containers. Destroy when empty. EPA REG. NO. 6830-26

ACCEPTED