

[6830-79]

OCTAGON PROCESS INC.
596 RIVER ROAD
EDGEWATER, NEW JERSEY 07020

OCTAGON DAIRY & LIVESTOCK SPRAY

A Ready-To-Use Pyrethrin Insecticide on Beef Cattle, Dairy Cattle, Horses, Swine, Sheep and Goats.

Acceptable for Use on Dairy and Meat Animals and in Dairy Barns, Milk Rooms and Dairies.

Kills and Repels Houseflies, Stable Flies, Horse Flies, Horn Flies, Mosquitoes, Gnats, and kills blood-sucking lice.

ACTIVE INGREDIENTS:

Pyrethrins	0.10%
*Piperonyl Butoxide, Technical	1.00%
Kerosene	98.90%

*Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% of related compounds.

CAUTION

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. Avoid contamination of feed or foodstuffs. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of spray equipment or disposal of wastes.

Remove pets, birds and cover fish aquariums before spraying.

Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place.

DIRECTIONS

ON ANIMALS: To protect animals from **horn flies, house flies, mosquitoes, and gnats**, apply a light mist sufficient to wet the tips of the hair. To control **stable flies, horse flies, deer flies**, apply 2 ounces per adult animal sufficient to wet the hair but not to soak the hide. Repeat treatment once or twice daily, or at intervals necessary to give continued protection.

To control **blood-sucking lice**, apply the spray lightly to treated areas using a stiff brush to get the spray to the base of the hair. Use 2 ounces or less per animal. Repeat every two or three weeks as indicated.

BARN, MILK ROOMS, DAIRY PLANTS: For **horn flies, mosquitoes, and gnats**, apply as a fog or fine mist directed toward the ceiling and upper corners until the area is filled with spray and about 2 ounces per 1000 cubic feet of space. For best results, close windows and doors before spraying and keep closed for ten to fifteen minutes. Ventilate the area and ventilate before reoccupying. Repeat application as necessary.

EPA REG. NO. 6830-79