


CHEMICAL & EQUIPMENT SALES & SERVICE CO. TM
Cessco TM
 P. O. Box 12452 / Charlotte, N. C. 28205



***TECHNICAL DATA-FLY CONTROL INSIDE WITH CURTAINS CLOSED**

Dosage required 59 sec. per 1000 cu. ft.
 Cost of treatments Approx. 1/2¢ per 1000 cu. ft.
 Delivery rate @ 70°F 1 gram per sec. per nozzle
 Cylinder Steel
 Propellent Gas High pressure

***TECHNICAL DATA-FLY CONTROL INSIDE WITH CURTAINS OPEN**

Dosage required 10 sec. per 1000 cu. ft.
 Cost of treatments Approx. 5¢ per 1000 cu. ft.
 Delivery rate @ 70°F 1 gram per sec. per nozzle
 Cylinder Steel
 Propellent Gas High Pressure

*Meets specifications when used in accordance with instructions in Specific Dosage Insecticide Aerosol Piping system with Automatic Timer.

CAUTION

Harmful or fatal if swallowed, inhaled or absorbed through the skin. Keep out of the reach of children, fowl and domestic animals. Do not contaminate feed, water or food stuffs. If clothing becomes contaminated with VAPONA[®], change clothing immediately and wash before reuse. Exposure to VAPONA[®] may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid. Atropine is antidotal.

WARNING

Flammable! Contents under pressure. Do not use near fire, sparks, or flame. Do not puncture or incinerate container. Exposure to temperatures about 130 F may cause bursting.

SPECIFIC DOSAGE TREATMENT

aerosol insecticide

FOR POULTRY CAGED LAYING HOUSES

To be used only with specific dosage insecticide aerosol piping system with automatic timer.

WARNING

Keep out of reach of children.

DO NOT REFILL.

When empty, close valves and return to manufacturer or dealer.

See side panel for additional cautions.

INGREDIENTS

| | |
|-----------------------------------------------|---------|
| Active Ingredient(s) | |
| **2, 2-Dichlorovinyl Dimethyl Phosphate | 93.0% |
| **Related Compounds | .070% |
| Vegetable Oil | 13.500% |
| Inert Ingredient | 85.500% |
| **Equivalent to 1.00% by weight SHELL VAPONA. | |

E.P.A. No. F959-34
 NET WEIGHT 250 LBS

DIRECTIONS


This treatment should not be made more often than once daily. Timer should be changed when curtains are raised or covered to coincide with directions. Nozzles should be strategically located and directed in the gable of the caged laying house so as to give complete coverage in the gable area where the air circulation is at a minimum and flies congregate, especially in the late afternoon. To avoid contamination of food, water and fowl, do not place nozzle within 5 feet of food, water or fowl.

FIRST AID


EXTERNAL: If skin or eyes become contaminated, change clothes and wash thoroughly with soap and water. For eyes, flush with water for 15 minutes and get medical attention. **INTERNAL:** If swallowed, induce vomiting by giving a tablespoon of salt in a glass of warm water. Repeat until vomit fluid is clear. CALL A PHYSICIAN IMMEDIATELY. SYMPTOMS of poisoning are pin-point pupils, drooling, labored breathing, vomiting, marked tremors, diarrhea, and convulsions. If these symptoms should occur after handling or being exposed to VAPONA[®], CALL A PHYSICIAN IMMEDIATELY. NOTE TO PHYSICIAN: If the above symptoms are present, atropine sulfate is the drug of choice. 2-PAM is also antidotal and may be administered in conjunction with atropine.

COVERAGE

Each lb. of aerosol insecticide will effectively treat approximately the same area as 7 gallons of an oil base spray without the use of over 600 gallons of oil.



POULTRY CAGED LAYING HOUSE



MANUFACTURED BY AEROSOL DIV. CESSCO, INC. CHARLOTTE, N. C.

AI 273 R(P573)

1959-34