


**ACCEPTED** Turf Fungicide 50W

14 JUL 1975

**contains DURENE**



**ACTIVE INGREDIENT:**

4000 m.s.m. N. of Laramie, 1300  
ft. above timberline.

## **INERT INGREDIENTS**

..... 50%  
..... 50%  
**TOTAL** 100%

100% owned subsidiary of Maranatha Corporation

EPA Reg. No. 7001-212-AA

EPA Est. No. 7001-CA-1

Fungicide 50 W is a fast-acting fungicide that is safe on lawns and turf. Its broad spectrum activity controls most diseases of lawns and turf. It has very good suspension stability and is suitable for all types of spray equipment. Turf Fungicide 50 W will control spot, powdery mildew, and other diseases of the sprayer for a preventive treatment. It is effective on a wide variety of grasses. It should be applied during the spring and summer months. Do not use the Turf Fungicide 50 W on lawns or turf during the fall.

1. *Phragmites australis* (Cav.) Trin. ex Steud.  
2. *Phragmites australis* (Cav.) Trin. ex Steud.  
3. *Phragmites australis* (Cav.) Trin. ex Steud.  
4. *Phragmites australis* (Cav.) Trin. ex Steud.  
5. *Phragmites australis* (Cav.) Trin. ex Steud.  
6. *Phragmites australis* (Cav.) Trin. ex Steud.  
7. *Phragmites australis* (Cav.) Trin. ex Steud.  
8. *Phragmites australis* (Cav.) Trin. ex Steud.  
9. *Phragmites australis* (Cav.) Trin. ex Steud.  
10. *Phragmites australis* (Cav.) Trin. ex Steud.

#### **DIRECTIONS FOR USE**

• The first part of the project involved the collection of data related to the characteristics of the population under study. The results of this phase are presented in the following section.

For example, the first report of a case of the disease at present is from Australia and reports have come in from 14 May. The earliest report of the disease in Europe and Central America came in from Mexico on 17 June. The first report of the disease in South America was made on 20 June.

“The first step in the development of a new field is to identify the key variables that will affect the outcome.”

## **RECOMMENDED APPLICATIONS**

**Lawns & Turf:** Brown Patch, Copper Spot, dollar Spot, Leaf Spot, Melting-Out, Rust, Snow Mold, Tephritis, Thrip. Apply specified dosage in 5 gallons of water per 1,000 square feet. Repeat at 1- to 2-week intervals throughout the spring and summer months. For control of Snow Mold, apply at lower specified rate after the first killing frost in the fall. A second application should be made when the lawn is free of snow in mid-winter, and this should be followed by a third application after the final melting of snow in late winter or early spring.

#### **Container & Waste Disposal:**

Residue equipment and residue of waste by burning of crop-tarants away from water supply. Disposal of container by burning with wastes or burning scrap-wood smoke. Never re-use containers.

**CAUTION!**

**KEEP OUT OF REACH OF CHILDREN**

Both the USA and the UK have strict rules about the use of GM food. The UK has a law that says GM food must be labeled as such. The USA does not have a similar law.

### **REFERENCES AND NOTES**

It is also important to remember that the results of our study must be interpreted with care, as the sample size was relatively small and the study design was observational.

<sup>1</sup> See, e.g., *Ward v. City of Sacramento*, 100 F.3d 1321, 1326 (9th Cir. 1996) (en banc).

After the selection of a project, the author will receive a copy of the properties specified and should make all necessary arrangements to have the same delivered to the author. All expenses for shipping, otherwise naturally occurring, insurance, and handling shall be paid by the author.

(GL 16)

NET WT. 4 LBS.



#### DEFINITION OF MEAN