

PM23

7401-40

145



Makers of fine lawn & garden and farm chemicals / Natural Guard Brand / Farmers Cooperative Brand / ferti-lome Brand / AMERICAN Brand / Hi-Yield Brand

Feb. 22, 1993

Joanne I. Miller, Product Manager 23  
U.S. EPA  
OPP  
Registration Division (H7505C)  
401 M Street SW  
Washington, D.C. 20460

I, Being an authorized representative of Voluntary Purchasing Groups, Inc. certify that all containers of ferti-lome Broad Spectrum Weed Killer, 7401-40, produced by June 15, 1994 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products. I further certify that all containers of said products sold or distributed by this company by Jan. 1, 1995 will bear revised labeling in accordance with the revised labeling required for Task Force Technical and Manufacturing-use products.

Sincerely,

A handwritten signature in dark ink, appearing to read "Raymond H. Smith".

Raymond H. Smith  
Product Manager

REV  
2 of 5

(Front Panel)

ferti-lome (R)

Broad Spectrum  
Weed Killer

- \* CONTROLS BROADLEAF WEEDS
- \* CONTROLS VINING TYPE WEEDS
- \* 5 TABLESPOONFULS MAKES 6 GALLONS OF SPRAY THAT WILL COVER 1000 SQ. FT.

Active Ingredients:

Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)* -----	3.43%
Dimethylamine Salts of related acids -----	.61%
Dimethylamine Salt of 2,4-dichlorophenoxyacetic acid** -----	20.00%
Inert Ingredients: -----	75.96%
Total -----	100.00%

\*Equivalent to 2.85% 3,6-dichloro-o-anisic acid (dicamba)

\*\*Equivalent to 16.61% 2,4-dichlorophenoxyacetic acid.

NET CONTENTS 32 FL. OZ. (ONE QUART) ✓

EPA Est. 7401-TX-1 ✓


EPA Reg. No. 7401-40 ✓

KEEP OUT OF REACH OF CHILDREN (12 point type)

CAUTION (18 point type)

See Back Panel for Additional Cautions

VOLUNTARY PURCHASING GROUPS, INC. BONHAM, TEXAS ✓



**BEST AVAILABLE COPY**

38

345

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** May be harmful or fatal if swallowed. Avoid contact with skin, eyes and clothing. When using this product, wear long-sleeved shirt, long pants, socks, shoes and chemical resistant gloves. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dust has settled.

**STATEMENT OF PRACTICAL TREATMENT**

- IF SWALLOWED-** Call a physician or Poison Control Center immediately. if possible, vomiting should be induced under medical supervision. Drink one or two glasses of water and induce Vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
- IF INHALED** - Remove victim to fresh air. Apply artificial respiration if indicated.
- IF ON SKIN** - Remove contaminated clothing and wash affected areas with soap and water.
- IF IN EYES** - Flush eyes with plenty of water. Call a physician immediately.

**ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates. Drifts or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment washwaters. Most case of groundwater contamination involving phenoxy herbicides such as 2,4-D have associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

**VOLUNTARY PURCHASING GROUPS, INC.**

**BONHAM, TEXAS 75418**

**NET CONTENTS \_\_\_\_\_ FL. OZS.**

**BEST AVAILABLE COPY**

DIRECTIONS FOR USE

GENERAL CLASSIFICATION: It is a violation of Federal Law to use this product in a manner inconsistent with it's labeling.

CONTROLS

- |                |                     |                  |
|----------------|---------------------|------------------|
| Dandelion      | Henbit              | Spotted Spurge   |
| Clovers        | Oxalis              | Stitch Wort      |
| Chicory        | Mustard             | Wild Aster       |
| Docks          | Pennywart           | Wild Carrot      |
| Dog Fennel     | Peppergrass         | Wild Garlic      |
| English Daisy  | Pigweed             | Wild Onion       |
| Frenchweed     | Plantain (Buckhorn, | Wild Radish      |
| Erect Knotweed | Broadleaf, Rugel's) | Wood Sorrel      |
| Ground Ivy     | Poison Oak          | Yarrow           |
| Hawkweed       | Prostrate Knotweed  | Puncturevine     |
| Healall        | Purslane            | Milkweed         |
| Lambsquarter   | Ragweed             | Bull Thistle     |
| Mallow         | Sheep Sorrel        | Shepherd's Purse |
| Morning Glory  | Mouseear Chickweed  |                  |

1. WHAT IS THIS PACKAGE FOR?

This package of ferti-lome BROAD SPECTRUM WEED KILLER is for control of weeds in lawns, golf courses, and similar turf areas. See side of container for a partial list of weeds controlled.

2. HOW DO YOU USE IT?

Mix 5 tablespoonfuls in 6 gallons of water. Apply thoroughly to 1000 sq. ft. (10' x 100'). Heavily infested lawns may require repeat application to kill missed weeds. Best results are obtained when weeds are young and actively growing. Application in early Spring and Fall when moisture and temperature make weeds grow rapidly is a good time to spray. Watering and fertilizing schedules should be maintained before and after spraying. Do not apply during periods of drought. If hose end sprayer is used, follow manufacturer's directions for proper mixing. Course sprays are less likely to drift. Do not apply within the dripline of desirable trees and shrubs.

BEST AVAILABLE COPY

5 of 5

For use on St. Augustine Grass mix 2 1/2 tablespoonfuls in 12 gals. of water. Apply thoroughly to 2000 sq. ft. of lawn area. Repeat application in 10-14 days if necessary.


Direct spray at undesirable plants. Do not use the same spray equipment for applying other material to plants, as injury will result.

If lawn begins to brown temporarily, water heavily. Do not use on Centipede grass, Dichondra, nor on lawns or turf where desirable Bent-grass or Clovers are present. Do not use on newly seeded lawns or turf.

3. HERE ARE THE RESULTS YOU MAY EXPECT!

ferti-lome BROAD SPECTRUM WEED KILLER used according to directions gives you economical and effective control of weeds in your lawn thus letting your lawn grasses grow vigorously to provide you with a beautiful weed free lawn.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.



170