

PREPARATION OF USE DILUTIONS

Dilute with water as directed.
Do not mix with soap.*

TO DILUTE 50% ROCCAL

First prepare a 10% solution of Roccal by carefully adding 1 gallon of the 50% concentrate to 4 gallons of water. Stir gently and, if necessary, filter solution before bottling. A porcelain, glass-lined or stainless steel mixing tank is recommended. Use this solution to prepare use dilutions. If siphon is used, use rubber bulb to start siphoning. DO NOT START SIPHON BY MOUTH.

TO PREPARE USE DILUTIONS OF ROCCAL

For Roccal 1:625 (1600 ppm), add 2 oz. (4 tablespoons) of Roccal 10% solution to 1 gallon of water and stir.

For Roccal 1:1250 (800 ppm), add 1 oz. (2 tablespoons) of Roccal 10% solution to 1 gallon of water and stir.

For Roccal 1:2500 (400 ppm), add ½ oz. (1 tablespoon) of Roccal 10% solution to 1 gallon of water and stir.

*When soap or an anionic detergent is used for cleansing, rinse with water before applying Roccal.

R-295

SOLD TO GRADUATE VETERINARIANS ONLY

50% SOLUTION

ROCCAL®

POWERFUL DISINFECTANT—SANITIZER—DEODORANT

Active ingredients — n-alkyl* dimethyl benzyl ammonium chlorides..... 50%

* (67% C₁₂, 20% C₁₄, 7% C₁₆, 2% C₁₈, 2% C₂₀, and C₂₂)

Inert ingredient — water..... 50%

NOTE: This product must be diluted as directed before use. After use of entire contents, rinse empty container thoroughly with water and discard it.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

DANGER: Keep Out of Reach of Children.

Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling the concentrate. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

5 GALLONS

Winthrop

DIRECTIONS FOR USE

FOR GENERAL DISINFECTION OF FLOORS, WALLS, KENNELS, ANIMAL CAGES, FURNITURE, BOOTS, AND APRONS: Clean thoroughly and rinse all surface areas before disinfecting. Then apply Roccal dilution 1:625 (1600 ppm) by mop, sponge, or spray and allow to dry. For routine sanitization of environmental surfaces where terminal disinfection is not required, use Roccal dilution 1:2500 (400 ppm).

FOR SANITIZING UTENSILS, FOOD AND WATER DISHES, AND FOOD PREPARATION SURFACES: Scrub and prewash with soap or detergent, and rinse thoroughly with water. Then use Roccal dilution 1:2500 (400 ppm). Dishes and other small items can be sanitized by immersion for at least two minutes in Roccal dilution 1:1250 (800 ppm), whereupon they are rinsed with water and allowed to dry.

In recommended use dilutions, Roccal is effective without sequestrants in water up to 400 ppm of hardness when tested by the official procedure.

EPA Reg. No. 7738-4-AA

Veterinary Department
Winthrop Laboratories
Division of Sterling Drug Inc.
New York, N. Y. 10016

RW 8-J

MADE IN U. S. A.

ACCEPTED

JUL 16 1974