PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Hay be harmful if swallowed. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist.

IF SWALLOWED: Get Medical Attention

IF IN EYES:

Flush eyes with Water for 15 Hinutes and

Get Hedical Attention.

IF ON SKIN:

Wash Skin with Soap and Water

ENVIRONMENTAL HAZARDS

Do not allow solution or apray drift to contact desirable plants. Do not allow on or near desirable turf, valuable trees or other plants, or in area into which their roots may extend, or on slopes where "run-off" will move the chemical into the root areas of useful plants. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of waste.

PHYSICAL BAZARDS

This product is corrosive to many metals. Keep away itom such materials.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product to a manner inconsistent with its labeling,

HOM-SELECTIVE CONTACT AND RESIGNAL VEGETATION KILLER IS A LEADYto-use weed and grass killer for general application to prevent vegetation growth on driveways, parking lots, walks, around fences, curbs, patios, buildings, and similar non-crit areas where plant growth is not desired.

This produce can also be used around drainage ditch banks, fuel tanks, roadways, storage acres, machinery, pumps, guard rails airports, military installations, highway medians, pipelines, railroads, lumberyards, industrial sites, light poles, fire hydrants, right-of-way or similar areas.

NON-SELECTIVE CONTACT AND RESIDUAL VEGETATION KILLER is also an effective control spray for control of: Woody perennials such as poison by, poison oak, and poison surac. Perennials such as leafy spurge, bitter dock, goldenrod, perennial raqueru, milkweed, and blueweed. Annuals such as crabgrass, broomweed, mickweed, cocklebur, jimsonweed, lambsquarters, larkspur, prickly lettuce, requeed and shepherds-purse.

NON-SELECTIVE CONTACT & RESIDUAL VEGETATION KILLER READY-TO-USE LIQUID

A CONTACT AND RESIDUAL DUAL ACTION NON-SELECTIVE WEED, GRASS, AND VEGETATION KILLER THAT MAKES SOIL BARREN. ALSO CONTROLS POISON IVY, POISON BAK AND OTHER WOODY PLANTS

ACTIVE INGREDIENTS: Sodium Chlorate (NaClOs)* 2.3001 Diquat dibromide (6,7-dihydrod-pyrido (1,2-a;2',1'-c) pyrazinedtium dibromide)** 0.154% INERT INGREDIENTS: 97.5463 ------TOTAL 100,000%

*Equivalent to 0.196 lbs. of active ingredients per gallon **Equivalent to 0.013 lbs. of active ingredient per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE ADDITIONAL PRECAUTIONS ON SIDE PANELS

EPA REG. NO. 8123-EPA EST. NO. 8123-IL-1

NET CONTENTS: __

Sold By

FRANK MILLER & SONS, INC. E D1831 SOUTH EMERALD AVENUE CHICAGO, ILLINOIS 60626

JUL 1 3 1987

Under the Federal Insecticide, Fungi ide, and Rodenticide Act. as amended, for the pesticide EPA Res. No.8/3/3

Best moults are obtained when applications are made during normal consture conditions and on clear days then the temberature is around 75° F. Avoid application in that, dry weather, since the killing action of HON-SELECTIVE CONTACT AND RESIDUAL VEGETATION. KILLER is through both the roots and the tops of plants, and therefore moisture is necessary after application to take the chemical into the soil to kill deep-rooted permodals. Fall application should be made in areas of 20 inches or less rainfall. The degree of control and duration of effect will vary with weed species, rainfall, temperature, and other conditions. By preventing weed growth early, you are decreasing the ever-present fire hazard that can become dangerous due to dead and dry vegetation.

HOW TO USE AS A SPOT TREATMENT: Adjust spray mozzle to give coarse spray and prevent drifting. Shake well and aim at center of vegetation and spray to wet. Best results are obtained when weeds and grass are small. One application may be sufficient. Effects begin to thow after a few days and weed gradually die. A repeat application may be required in 3 - 4 weeks for hard to control weeds or if rain occurs within 24 hours following application.

| 1 quart controls | 15 sq. | It |
|--------------------|-------------|----|
| 2 quarts controls | 30 sq. | it |
| 1 gallon controls | 60 sq. | ft |
| 5 gallons controls | 300 sq. | ft |

AS AM EDGING LIQUID: Shake well and uniformly spray 1 gallon per 60 equare feet of soil surfact or a strip 4 inches wide, 180 feet long. To make a 4 Inch band hold applicator end about 6 inches from surface to be treated. To obtain wider band, raise container. Thoroughly wet all foliage and stems of undesired vegetation to insure contact action.

INPORTANT: Do not use this product for controlling weeds in vegetable gardens, flower beds or around shrubs or ornamental plants.

STORAGE AND DISPOSAL

STORAGE: Keep from freezing. To be stored in original container and placed in an area inaccessible to children.

PESTICIDE DISPOSAL: Produce remaining in original container should be disposed of by securely wrapping original container in several layers of neupapers and discarding in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Discard empty container in trash.

NOTICE: Buyer assumes all risks of use, storage and handling this product not in street accordance with directions given herein.