

**FRONT PANEL:****T & C****1 PINT****CRABGRASS KILLER**

ALSO KILLS: FOXTAIL, CHICKWEED, DALLIS GRASS, BARNYARDGRASS,  
SANDBUR, WITCHGRASS, JOHNSON GRASS


USE ON LAWNS OTHER THAN St. Augustine Centipede, Carpet and  
Bahia grass and NON-CROP AREAS.

**ACTIVE INGREDIENTS:**

Disodium Methaneearsonate . . . . .	12.3%
INERT INGREDIENTS: . . . . .	87.7%
TOTAL 100.0%	

Total Arsenic as elemental, all in water soluble form 5.00%  
Equivalent to 20.0% Disodium Methaneearsonate Hexahydrate.

**CAUTION: KEEP OUT OF REACH OF CHILDREN.  
FOR ADDITIONAL CAUTIONS SEE BACK PANEL.**

**PRODUCT NO. 24623****U.S.D.A. REG. NO. 8222-25****BACK PANEL:****DIRECTIONS FOR USE**

**HOW TO USE:** Apply at the rate indicated below. Spray when conditions for growth of turf grasses are good....During droughts water several days before treating and fertilize to restore green color. Keep lawn mowed 1½ to 2 inches during treatment. Spray newly seeded lawns at "Seedling" rate below after they are well established. Do not use on St. Augustine, Bahia Carpet or Centipedegrass.

**CAN BE USED ON:**

TYPE OF LAWN	PESTS	DIRECTIONS
New lawns, more sensitive grasses, such as, Fescues and Bentgrass in early Spring or late Fall during cool days and moist soil conditions.	Crabgrass in "seedling" stage, witch grass, Barnyardgrass,	Spray at rate of 14 tbsps. or 7 ounces to each 5 gallons of water per each 1000 sq. ft. (25 ft. x 40 ft.) of lawn in early summer when crabgrass is young. Apply 2 or more treatments at 7-day intervals. If new crabgrass growth appears later in the season, repeat applications may be necessary.
	Crabgrass in "Branching" stage.	Spray at rate of 14 tbsps. to 5 gallons of water per each 700 sq. ft. (20 ft. x 35 ft.). Apply 2 or more treatments at 7-day intervals.
Established lawns, such as, Kentucky, Merion Blue, Dichondra, Zoysia, Bermuda.	Crabgrass in "Branched" stage, witch grass, foxtail, Barnyard grass, dallisgrass, Sandbur, Chickweed..	Spray at rate of 24 tbsps. or 12 ounces to 5 gallons of water per each 1000 sq. ft. (40 ft. x 25 ft.). Repeat spraying in 7 days. Application may discolor, but will not permanently injure desirable grasses

BACK PANEL CONTINUED:

On non-crop or non-pasture areas, such as, drainage ditch banks, fence rows, rights-of-way, to control johnson grass, dallis grass, sandburs, barnyard grass, spray on grass to point of run-off diluted with water at the rate of 1 pint in 4 gallons of water, 1 quart in 8 gallons water, 1 gallon in 32 gallons of water.

**CAUTION:**

Harmful if swallowed. If taken internally, induce vomiting. CALL A PHYSICIAN AT ONCE. Avoid contact with skin or breathing of spray mist. Keep away from children and domestic animals. Avoid contamination of feed and foodstuffs. Wash hands thoroughly after using. Do not contaminate water used for irrigation or domestic purposes.

Keep livestock off treated area.

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

Packed for TOWN & COUNTRY ENTERPRISES, Union City, California  
By GENERAL PACKAGING COMPANY, INC.