

directions

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS

DOGS AND CATS: To control fleas apply sufficient dust to treat animal thoroughly using a garden type duster or shaker type dispenser. Rep at every 2 or 3 weeks as needed. Do not treat dogs and cats under four weeks old. Place clean bedding in kennels and pens after treatment.

ANIMAL QUARTERS: To control fleas in kennels, pens, yards, lawns, under houses, remove manure and other debris. Apply 1 or 2 lbs. of dust per 1000 sq. ft. using a garden type duster.

THOROUGH COVERAGE OF ENTIRE PLANT, INCLUDING LOWER LEAF SURFACES IS ESSENTIAL FOR GOOD INSECT CONTROL.

APPLES, PLUMS, PRUNES: To control clover mite, codling moth, European red mite, eye-spotted bud moth, green apple aphid, lecanium scale, red banded leaf roller, rosy apple aphid, 2-spotted mite, 4-spotted mite, apply when insects appear. Repeat as needed. Do not apply within 3 days of harvest.

ASPARAGUS: To control asparagus beetle apply when insects first appear. Repeat as necessary. For thrips apply immediately after cutting. Do not apply within 1 day of harvest.

BEANS, BLACKBERRIES, RASPBERRIES, TOMATOES: To control adult Japanese beetles, aphids, leafhoppers, Mexican bean beetles, mites, psyllid, raspberry fruit worm, serpentine leaf miner, apply when insects appear. Repeat as needed. Do not apply within 1 day of harvest.

BEETS, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, SPINACH: To control aphids, flea beetle, garden webworm, leaf miner, mites, and thrips, apply when insects appear. Repeat as needed. Do not apply within 7 days of harvest.

BROCCOLI, EGGPLANT, ONIONS, PEPPERS, STRAWBERRIES, TURNIPS: To control aphids and thrips apply when insects appear. Repeat as needed. For maggot make four weekly applications beginning in July. Do not apply within 3 days of harvest.

CUCUMBERS, MELONS, SQUASH: To control aphids, borers, cucumber beetles, squash bugs, stink bugs, apply when insects appear. Repeat as needed. Apply only when foliage is dry. Do not apply within 1 day of harvest.

LETTUCE: To control leafhoppers and tarnished plant bugs apply when insects appear. Repeat as needed. Do not apply within 14 days of harvest on leaf lettuce or within 7 days of harvest on head lettuce.

PEACHES: To control cottony peach scale, European red mite, lecanium scale, oriental fruit moth, plum curculio, 2-spotted spider mite, apply when insects appear. Repeat as needed. Do not apply within 7 days of harvest.

PEARS: To control pear psylla and plum curculio, apply when insects appear. Repeat as needed. Do not use on Bosc variety or apply within 1 day of harvest.

PEAS: For pea aphids apply when insects appear. Repeat at 7 day intervals as needed. Do not apply within 3 days of harvest.

POTATOES: To control aphids apply when insects appear. Repeat as needed.

FLOWERS AND ORNAMENTALS: Use on many flowers and ornamentals to control adult Japanese beetles, aphids, bagworms, chrysanthemum fly, leafhoppers, mealy bugs, psyllids, mites, thrips. Use when insects first appear. Repeat as needed. May cause injury to certain ferns including Boston, Maidenhair, and Pteris, as well as some species of Carssula.

RATE OF APPLICATION FOR ABOVE USES: Apply at a rate to give a thin uniform covering to the plants. One pound of dust will treat approximately 400 feet of row depending on plant size.

directions

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS

DOGS AND CATS: To control fleas apply sufficient dust to treat animal thoroughly using a garden type duster or shaker type dispenser. Repeat every 2 or 3 weeks as needed. Do not treat dogs and cats under four weeks old. Place clean bedding in kennels and pens after treatment.

ANIMAL QUARTERS: To control fleas in kennels, pens, yards, lawns, under houses, remove manure and other debris. Apply 1 or 2 lbs. of dust per 1000 sq. ft. using a garden type duster.

THOROUGH COVERAGE OF ENTIRE PLANT, INCLUDING LOWER LEAF SURFACES IS ESSENTIAL FOR GOOD INSECT CONTROL.

APPLES, PLUMS, PRUNES: To control clover mite, codling moth, European red mite, eye-spotted bud moth, green apple aphid, lecanium scale, red banded leaf roller, rosy apple aphid, 2-spotted mite, 4-spotted mite, apply when insects appear. Repeat as needed. Do not apply within 3 days of harvest.

ASPARAGUS: To control asparagus beetle apply when insects first appear. Repeat as necessary. For thrips apply immediately after cutting. Do not apply within 1 day of harvest.

BEANS, BLACKBERRIES, RASPBERRIES, TOMATOES: To control adult Japanese beetles, aphids, leafhoppers, Mexican bean beetles, mites, psyllid, raspberry fruit worm, serpentine leaf miner, apply when insects appear. Repeat as needed. Do not apply within 1 day of harvest.

BEEETS, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, SPINACH: To control aphids, flea beetle, garden webworm, leaf miner, mites, and thrips, apply when insects appear. Repeat as needed. Do not apply within 7 days of harvest.

BROCCOLI, EGGPLANT, ONIONS, PEPPERS, STRAWBERRIES, TURNIPS: To control aphids and thrips apply when insects appear. Repeat as needed. For maggot make four weekly applications beginning in July. Do not apply within 3 days of harvest.

CUCUMBERS, MELONS, SQUASH: To control aphids, borers, cucumber beetles, squash bugs, stink bugs, apply when insects appear. Repeat as needed. Apply only when foliage is dry. Do not apply within 1 day of harvest.

LETTUCE: To control leafhoppers and tarnished plant bugs apply when insects appear. Repeat as needed. Do not apply within 14 days of harvest on leaf lettuce or within 7 days of harvest on head lettuce.

PEACHES: To control cottony peach scale, European red mite, lecanium scale, oriental fruit moth, plum curculio, 2-spotted spider mite, apply when insects appear. Repeat as needed. Do not apply within 7 days of harvest.

PEARS: To control pear psylla and plum curculio, apply when insects appear. Repeat as needed. Do not use on Bosc variety or apply within 1 day of harvest.

PEAS: For pea aphids apply when insects appear. Repeat at 7 day intervals as needed. Do not apply within 3 days of harvest.

POTATOES: To control aphids apply when insects appear. Repeat as needed.

FLOWERS AND ORNAMENTALS: Use on many flowers and ornamentals to control adult Japanese beetles, aphids, bagworms, chrysanthemum fly, leafhoppers, mealy bugs, psyllids, mites, thrips. Use when insects first appear. Repeat as needed. May cause injury to certain ferns including Boston, Maidenhair, and Pteris, as well as some species of Carex.

RATE OF APPLICATION FOR ABOVE USES: Apply at a rate to give a thin uniform covering to the plants. One pound of dust will treat approximately 400 feet of row, depending on plant size.



AGWAY

**4% malathion
flower and
vegetable
dust**

**FOR INSECTS OF MANY FLOWERS,
FRUITS, ORNAMENTALS,
VEGETABLES AND PETS**

ACTIVE INGREDIENTS:

*Malathion	4.0%
INERT INGREDIENTS, not more than	<u>96.0%</u>
	100.0%

*0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate.

caution:

KEEP OUT OF REACH OF CHILDREN

See additional cautions and warnings on side panel.

NET WEIGHT 4 LBS.

CAUTION: Avoid breathing dust. Wash thoroughly. Avoid contact with skin and clothing.

This product is toxic to fish and aquatic life. Do not discharge into water by direct application. Proper disposal of wastes and containers is required.

This product is highly flammable. Do not use near open flame or heat. Do not use in enclosed spaces. Do not use in areas where food is obtained from your service.