

A C W A Y P R O C U R T I D U A L C O M P A N Y

ITEM: TABLE COPY + Big Red Flea and Tick Dust

DATE: 4/3/68

((FRONT PANEL))

NET WEIGHT
Four Ounces

((ACWAYT.M.))

BIG RED FLEA AND TICK DUST


ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-methylcarbamate) ... 5.0%

INERT INGREDIENTS: 95.0%
100.0%

CAUTION: ((18 pt. type)) KEEP OUT OF REACH
OF CHILDREN. ((12 pt. type)) See additional
cautions and warnings on ((fill in)) panel.

ACCEPTED



ASIAN PESTICIDE COMPANY - CHINA TRADE CO., LTD.

ONE DOSE, COPY = 1/4 Bag Flea and Tick Dust

4/3/63

(W. REED PARK))

~~DOGS AND CATS~~
for Control of
Brown dog ticks and Fleas:

Apply Big Dog Flea and Tick Dust to the animal and rub into skin. Apply in sleeping quarters weekly. Do not treat kittens under four weeks old.

WARNING: Seller warrants that the product conforms to its chemical description etc. is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

NOTE: This pesticide is to be sold ONLY in this original, unbroken package.

MANUFACTURED BY AGCHY INC. U.S.D.A. REG. NO.
SYRACUSE, N.Y. 3596-

ACCEPTED

5-24-68

8570-291

Big Red Flea and Tick Dust

Product of U.S.A. - Item No. 100-1000

((WALKER PANEU))

C A T I O N : Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Skin contact may be harmful. Wash hands before eating or smoking. Avoid contamination of food, feeding troughs and watering receptacles.

N O T E T O P H Y S I C I A N S : Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WILKINSON OR
SALM. USE STRICTLY IN ACCORDANCE WITH LABEL
CAUTIONS, WARNINGS AND DIRECTIONS.

BIG RED FLEA, HORSES, SHEEP, BOGS

Big Red Flea and Tick Dust may be applied up to seven days before slaughter on beef cattle, horses, sheep and hogs. Treatments may be repeated once every four days, if necessary. Big Red Flea and Tick Dust should not be used on dairy animals. A rate of two ounces of 5 per cent dust, Big Red Flea and Tick Dust will control horn flies on livestock.

POULTRY

Big Red Flea and Tick Dust is highly effective against the northern fowl mite even where it has shown resistance to phosphate insecticides. Apply up to seven days before slaughter. Avoid direct contamination of eggs. Do not treat nest litter. Apply one pound of 5 per cent dust per 100 birds or in dust bath boxes apply 2½ pounds of five per cent dust for each 50 birds.

Control of Northern fowl mite, chicken mite,
and fleas: ((Insect following "POULTRY"))
Dust box treatment is a substitute to other
method for chicken mite and flea control. On range,
no natural walls may be substituted for such fence
box.

ACCEPTEE
5-24-68
8590-291