AGWAY GREENLAWN PLUS 20-5-7

WITH 5% DACTHAL

FOR ESTABLISHED LAWN USE ONLY.

23# treats 5,000 sq. ft. (10.43 kg treats 465 sq. meters) ((46# treats 10,000 sq. ft. (20.87 kg treats 930 sq. meters))

Contains 5% Dacthal for established lawns. Crabgrass Preventer preemergent Herbicide plus Lawn Fertilizer, Iron added for Greener Lawns

ACTIVE INGREDIENT		
Dimethy! Tetrachloroterephthalate*		5 00%
INERT INGREDIENTS**		95 00 %
TOTAL		100 000 0
* Use as a preemergence herbicide patented unde	r U.S. Pat. No. 2,923 634	
" FERTILIZER GUARANTEED ANALYSIS		
Total Nitrogen (N)		20 00%
6.7% Water Insoluble Nitrogen		
Available Phosphoric Acid (P ₂ O ₄)		5 00°°
		7 00%
tron (Fe)		0.10%

KEEP OUT OF REACH OF CHILDREN CAUTION

Read entire label before using this product

FOR ESTABLISHED LAWNS ONLY

1. Prevents Crabgrass and other annual lawn weeds

Registered Trademark of Diamond Shamrock Corp.

2. Plus Fertilizes your Lawn with 20-5-7

AGWAY GREENLAWN PLUS 20-5-7 WITH 5% DACTHAL is a very effective fertilizer and urabgrass control preamergent herbicide compound. The fertilizer fracedients provide for prous hardy turf growth. 3.92 pounds of Nitrogen is applied to each 1,000 square feet. -0.45 kg per 100 sq. meters). Slow release greaform is combined with quick acting water so uble. Nitrogen for steady in form feeding. Phosphorus and Potash are available for good. of and Stem development

The 5% Day that in this product willig selexication to reamergence control of crabgrass and the following weeds in turf and ornamental plantings

Carpetweed Common Chickweed Fall Panicum Florida Pussley, Green Foxtail Hairy Crabgrass, Johnsongrass, Lambsquarters, Lovegrass, Purslane, Smooth Crabgrass, Texas Millet, Witchgrass and Yellow Foxtail

DIRECTIONS FOR USE. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

Do Not Hand Spread

Do Not use on Dichondra Lawns or Newly sended lawns

STORAGE AND DISPOSAL. Do not contaminate water, food, feed, seed or other pesticides by storage, disposal or cleaning of equipment

PESTICIDE DISPOSAL. Pesticide ispray mixture or rinse water that cannot be used according to label instructions must be disposed of ail ording to applicable Federal istate or local procedures

CONTAINER DISPONAL Completely empty bag by shaking and tapping sides and bottom to loosen chinging particles. Empty residue into application equipment. Then dispose of hag in a sanitary landfill or by incineration if allowed by state and local authorities

WHEN TO APPLY Agway Greeniawh Phys 20.5.1 with 5% Daothal kills crabgrass and other annual weeds as they germinate. But not after they have rooted. To be effective, application. must be before weed seed germination in early scring. Crabgrass seed germination usually coincides with the time Forsythia blossoms start to fall. A single နည်းပြသမ်းက in နarty သုံးကရ before weed seed germination usually provides control up to increamonths in areas which commonly experience stands of late-germinating respicass and other annual veedy grasses, a second application at half the regular rate can be safely made two months after the first application

HOW MUCH TO APPLY Apply contents of bag (23# (10 43 kg)) evenly to 5,000 sq. ft. (465 sq. meters) or 4.6# per 1,000 sq. ft. (1.95 kg per 100 sq. meters). Use any lawn spreader in good condition. Do not spread over larger area than recommended as such application would give poor results

HOW TO APPLY Before making application, lawn should be raked to remove extraneous material. In order to obtain expected results, Agway Greenlawn Plus 20-5-7 with 5% Dacthal. should be evenly applied with no skips or misses.

it is suggested that the area be covered twice, the second time at right angles to the first to insure thorough coverage (see attached table)

SPREADER SETTING GUIDES'

	Rate Per 100 Sq 1000 Meters Sq Ft		Cyclone B1 Spinner Type		Ortho Spinner Type		Scotts Spinner Type		Ortho Drop Type		Scotts Drop Type	
1			Once Over	Twice	Once Over	Twice Over	Over Over	Oher Imice	Once Over	Twice Over	Once Over	Twice
ń	325	4 6										
×	kg	lbs	5	4	6	3	[E	В	4	3₩	6	4 7

NOTE: These settings are furnished as a guide only, age, condition of spreader, speed of operation and roughness of terrain can cause variation in rates applied

Do not rake the lawn or disturb soil surface after application

Water in lightly if no rain falls within a week after application

RE-SEEDING. When some spring seeding is necessary, there should be a delay of approximately 60 days after the application. This delayed seeding improves the survival of desirable grass seedlings. If a re-seeding is done before 60 days after application, the amount of seed should be increased 50% to insure thrifty stands.

An early spring application may be made to newly seeded lawns after they have exhibited greening of newly-sprouted lawn grasses

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Do not graze animals on treated areas or feed plant refuse to livestock

In case of contact, flush eyes with water for 15 min , for skin, wash with plenty of soap and water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS. Keep out of takes, streams or ponds. Do not contaminate water used for irrigation or domestic purposes

DISCLAIMER Buyer assumes all risks of use, storage, or handling not in strict accordance with directions given herewith

NOTE. This pesticide is to be sold ONLY in this original unbroken package

EPA Reg. No. 8590-482

Code Number 86-4775

EPA Est No 8590-NY-1



AGWAY INC., BOX 4933, SYRACUSE, NY 13221

NET WT. 23 LBS. (10.43 kg)

A 2.82

