

AGWAY

GREENLAWN PLUS 21-5-7 FERTILIZER WITH BROADLEAF WEED KILLER FORMULA III

Contains 2,4-D and MCPP

Weeds and Feeds 465 sq. meters (5,000 sq. ft.) Kills dandelions, clover, plantain, chickweed, and many other broadleaf weeds as listed. Feeds and builds beautiful, thick green lawns.

ACTIVE INGREDIENTS:

* 2-Ethylhexyl Ester of 2,4-Dichlorophenoxyacetic	
Acid	1.19%
**Isooctyl ester of 2-(2-methyl-4-chlorophenoxy)	
propionic acid	1.20%
INERT INGREDIENTS:	<u>97.61%</u>
Total	100.00%

*Equivalent to 0.79% 2,4-dichlorophenoxyacetic acid **Equivalent to 0.79% 2,(2-methyl-4-chlorophenoxy)propionic acid

Isomer specific by AOAC Method No. 6.Dol-5

EPA Reg. No. 8590-522 EPA Est. No. 8590-NY-1

ACCEPTED ACCEPTED ACCEPTED Under the Federal Insociatido, Fungicido, and Rodontialido Act. as americal, for the publicido registered under EPA Reg. No. 0590-522

KEEP OUT OF REACH OF CHILDREN CAUTION

See Additional Precautionary Statements On Rear Panel

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS AND DIRECTIONS.

Covers 5,000 sq. ft. (465 sq. meters)

NET WEIGHT: 30 Pounds

Manufactured for:

AGWAY INC. P.O. Box 4741 Syracuse, NY 13221

. HUG-29-'95 13:12 ID:UNIVERSHL-CHEMICAL

auf 4

(Back)

(

Weeds and Feeds 465 square meters (5,000 scuare feet) Net Wt. 13.61 kg (30 lbs.)

AGWAY GREENLAWN PLUS 21-5-7 FERTILIZER WITH BROADLEAF WEED FILLER - FORMULA III

GUARANTEED ANALYSIS:

Total Nitrogen (N)	10.00%
1.75 Coated Slow Release Urea Nitrogen	
8.25 Water Soluble Nitrogen	
Available Phosphate $(P_2 0_3)$	6.00%
Soluble Potash (K ₂ 0)	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Causes eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid b eathing dust. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

FIRST AID

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce voluiting.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates Drift or runoff may adversely affect aquatic invertebrates and non target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

Protective Clothing

When using this product, wear long-sleeved shirt, long pants, socks and shoes.

Worker Reentry

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dust has settled.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

344

STORAGE & DISPOSAL

والالالا بالالالية بالمراجعة

Do not contaminate water, food, or feed by storage or disposal.

A REPORT A MERICAN REPORT OF REPORT OF REPORT

Keep bag closed when not in use. Store only in original container in a cool, dry area that can be locked and is out of the reach of children.

Do not reuse empty bag. Discard bag in trasl.

This product rejuvenates undernourished and weedy lawns by:

- 1. Providing a controlled feeding fertilizer which gives a green lawn, and
- 2. Controlling unsightly broadleaf weeds such as:

ĺ

aster	field bindweed	pigweed
buckhorn	filaree	pineapple weed
buttercress	galinsoga	plantain
black medic	ground ivy	prickly lettuce
burdock	heal all	purslane
buttercup	henbit	ragweed
buttonweed	hop clover	shadowflower
chickweed	knotweed (seedling)	shepherdspurse
chicory	lambsquarters	sow thistle
carpet weed	matchweed	smartweed
cinquefoil	creeping charlie	stitchwort
cudweed	moneywort)	thyme leaf
curly dock	morning glory	(sandwort)
daisy fleabane	mustard	vetch
a dandelion	oxalis	white clover
english daisy	pennywort	wild radish
evening primrose	pepper weed	yellow rocket

WHEN TO APPLY: This product should be applied in the spring or fall when weeds are young and growing vigorously. Do not use this product during hot summer months since injury may occur. To avoid injury to new growth, new lawns should not be treated until after the fourth mowing. Established lawns may be injured if treated when in need of water. Water thoroughly 3 to 4 days prior to treatment and again the day before treatment. Grass may be injured if treated in periods of high heat and humidity. For this reason, apply only when the daytime temperature is not expected to exceed 80°F for 48 hours. To avoid drift, apply when air is calm. Do not apply if rain is expected within 24 hours.

HOW TO APPLY: The contents of this bag will cover an area of 5,000 sq. ft. Apply at rate of 2.92 kg/100 sq. meters (6.0 lbs./1,000 sq. ft.). Apply to lawn when grass is wet (after rain, sprinkling, or dew) so pellets will stick to the weeds. Do not water or mow lawn for at least 2 days after treatment, then resume normal watering program. The maximum number of broadcast applications per treatment site is 2 Jer year. Supplement with additional feeding of Turfood Special 10-6-4 Pertilizer as recommended for year-round nutrition of the lawn.

444

IMPORTANT NOTICE: Since this product kills desirable as well as undesirable broadleaf plants, it may injure flowers, ornamentals, and trees. Vapors from this product may injure susceptible plants in the immediate vicinity. I've not exceed recommended rate. Avoid contact with flower beds or shrubs. If accidentally deposited on flowers or shrubs, hose off immediately. Do not use on Bentgrass, Dichondra, Carpetgrass, St. Augustine, or ground covers. Delay planting grass seed until 4 weeks after treatment. Read and follow directions carefully.

DISCLAIMER: Buyer assumes all risks of use, storage, or handling not in strict accordance with directions given herewith.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

SPREADER SEITING GUIDE** FOR AGWAY GREENLAWN PLUS 21-5-7 FERTILIZER WITH BROADLEAF WEED KILLER - FORMULA III

	Cyclone B1	Ortho	Scotts	Ortho	Scotts
Rate Per	Spinner Type	Spinner Type	Spinner Type	Drop Type	Drop Type
100 Sq. 1000	Once Twice	Once Twice	Once Twice	Once Twice	Once Twice
Meters Sq.Ft.	Over Over	Over Over	Over Over	Over Over	Over Over

2.92kg 6.0 lbs

(

NOTE: These settings are furnished as a guide <u>only</u>. Age, condition of spreader, speed of operation, and roughness of terrain cause variation in rates applied.