

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: AGWAY TREE FRUIT SPRAY

DATE: 7/2/82

ACCEPTED
with conditions
in EPA Letter 8590-12-1

NET CONTENTS: 8 oz.

EPA REGISTRATION
NO. 8590-12-1

NOV 18 1982

ACCEPTED
NOV 18 1982
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 8590-12-1

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 8590-12-1
AGWAY TREE FRUIT SPRAY

Controls aphids, mites, codling moth, on apples & peaches and other tree fruits.

ACTIVE INGREDIENTS:

Captan N-((trichloromethyl)thio)-4-cyclohexene-1,2-dicarboximide	9.46%
Related Compounds	0.54%
Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate).	6.00%
*Methoxychlor, technical	12.00%
Carbaryl (1-naphthyl N-methylcarbamate).	3.00%
Xylene range aromatic solvent	59.83%
INERT INGREDIENTS:	9.17%
TOTAL	100.00%

*Equivalent to 10.56% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 1.44% of other isomers and related compounds.

KEEP OUT OF REACH OF CHILDREN (12 pt.)

CAUTION (18 pt.)

See side panel for additional precautionary statements.

EPA Reg. No. 8590-
Epa Est. No. 8590-12-1

Agway Inc.
Chemical Division
Box 4933
Syracuse, NY 13211

OK BY

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR:

AGWAY TREE FRUIT SPRAY

DATE: 7/2/82

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS (& DOMESTIC ANIMALS)

CAUTION: Harmful if swallowed or inhaled. Avoid breathing spray mist or contact with skin, eyes or clothing. May be absorbed through skin. Do not use on household pets or humans. Wash hands and contaminated clothing after use. Do not smoke while applying. Store out of reach of children.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

If on skin, remove contaminated clothing and wash affected areas with soap and water.

If in eyes, flush eyes with plenty of water.

Call a physician if irritation persists.

Remove residue on edible fruit by wiping or washing before canning or eating. Avoid contamination of foodstuffs, feed or feeding troughs

FOR INFORMATION
CALL 1-800-4-A-AGWAY

OKSY

SHEET NO. 2 of 6

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR:

AGWAY TREE FRUIT SPRAY

DATE: 7/2/82

and water receptacles or pets or animals, outdoor fish ponds, streams and water supplies.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to lakes, streams or ponds.

Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as recommended on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

EASY TO USE: Mixes with water instantly; no plugged nozzles; no messy powders to handle, measure or mix; no pre-mixing or straining necessary. Designed especially for home orchards to protect apples, cherries, grapes, peaches, pears, plums and prunes from the ravages of insects and diseases listed on spray chart below.

DIRECTIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

PLA 50 CHARACTERS
ELITE 100 CHARACTERS

BY

3

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR:

AGWAY TREE FRUIT SPRAY

DATE: 7/2/82

SHAKE PRODUCT THOROUGHLY BEFORE USING. Contains micronie particles which settle upon standing and require reblending by agitation. Choose a cool, calm period, preferably early morning or evening, when trees will not be subjected to hot direct rays of the sun. Do not apply if rain is expected before leaf surfaces dry. It is easier to prevent disease and scale damage than to cure same, therefore a preventative 10-14 day spray schedule is recommended. When insects are present, use maximum dilution. When active disease spores are present (warm, humid days) shorten spray intervals to 5-7 days at maximum dilution.

Apply spray to leaves, fruit branches, trunk and base of tree until lightly dripping. Shake sprayer occasionally or agitate to keep spray particles in suspension during application.

Follow spray schedule given below in tablespoons per gallon of water. For larger quantities, use 1 gallon per 100 gallons of water. Observe caution statement and cautions for fruit to be treated.

STORAGE & DISPOSAL (12 point)

Do not reuse empty container. Wrap container and put in trash collection.

PICA - 50 CHARACTER
ELITE - 30 CHARACTER

COPY

SHEET NO. 4 of 6

KEY CHART

APPLY THE FOLLOWING AMOUNTS (GIVEN IN TABLESPOONS) IN ONE GALLON OF WATER.

Fruit	Insects/Diseases	Dormant	Green Tip	Pre Bloom	Full Pink	Bloom	Petal Fall	1st Cover	2nd Cover to Harvest	Post Harvest	
Apple:	apple maggots, codling moths, Jap beetles, leafhoppers, plum curculios, aphids, bud moth, Forbes scale, eastern tent caterpillar, red banded leaf roller, mites, bitter rot, black rot, frog-eye leaf spot, black pox, botryosphaeria (white rot), botrytis rot, bullseye rot, Brooks fruit spot, flyspeck, cedar rust, quince rust, scab, sooty blotch	Dormant - absence of growth-fall, winter or spring. Dormant sprays are recommended. Use Agway Dormant Oil or Dormant Oil with Ethion up to 7 days before Green Tip application.	TWO	THREE	THREE		THREE	THREE	THREE	TWO	
			Start and repeat at 10 day intervals until bloom				Repeat at 14 day intervals up to 14 days before harvest				
			Green Tip -- when branch tips show green tissue	Pre-bloom -- when blossom buds begin to show color	Full Pink -- when blossom buds show full color before opening		Petal Fall -- when last petals have fallen from blossom	1st Cover -- 7 - 10 days after petal fall	2nd Cover -- to cut off date before harvest	Post Harvest -- to leaf drop	
Cherry:	plum curculios, cherry fruit flies, cherry fruit worms, black cherry aphid, fruit tree leaf roller, bud moth, Jap beetle, Botrytis rot, brown rot, leaf spot			TWO	THREE	THREE		THREE	THREE	THREE	TWO
				Start and repeat at 10 day intervals until bloom				Repeat at 14 day intervals up to 14 days before harvest			
Grape:	flea beetles, Jap beetles, grape leaf skeletonizers, rose chafers, grape berry moths, spider mites, mealy bugs, leafhoppers, black rot, downy mildew (berries only), botrytis rot		TWO	THREE	THREE		THREE	THREE	THREE	TWO	
			Start and repeat at 10 day intervals until bloom				Repeat at 14 day intervals up to 14 days before harvest				
Peach:	plum curculio, Jap beetle, spider mite, oriental fruit moth, green, peach, black cherry, black peach, rusty plum aphids, brown rot, rhizopus, scab, brown rot (blossom blight), coryneum blight		TWO	TWO	THREE		THREE	THREE	THREE	THREE	
			Start and repeat at 10 day intervals until bloom				Repeat at 14 day intervals up to 21 days before harvest				
Pear:	codling moth, pear slug, fruit tree leaf roller, pear psylla, plum curculio, red banded leaf roller, mites, mealybugs, fruit spot, scab		TWO	THREE	THREE		THREE	THREE	THREE	TWO	
			Start and repeat at 10 day intervals until bloom				Repeat at 14 day intervals up to 7 days before harvest				
			*Russeting may be induced on Bosc pears. Do not use on D'Anjou pears.								
Plum, Prune:	plum curculios, mealy plum aphid, bud moth, brown rot, russet or lacy scab		TWO	TWO	THREE		THREE	THREE	THREE	THREE	
			Start and repeat at 10 day intervals until bloom				Repeat at 14 day intervals up to 7 days before harvest				

Bloom - When blossoms are open. Don't spray pollinators.
*Fungicide alone may be applied for disease

BEST DOCUMENT AVAILABLE

5 of 6

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR:

AGWAY TREE FRUIT SPRAY

DATE: 7/2/82

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

PICA - 500-APR-1982
82 72 - 60 CHARACTERS

CK 6Y

SHEET NO. 6 OF 5