US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (18-767) WASHINGTON, DC 20460

TERM OF ISSUANCE NAME OF PESTICIDE PROTUCT

NOTICE OF PESTICIDE: REGISTRATION REREGISTRATION

(Under the Federal Insecticide, . ungicide, and Rodenticide Act, as amended)

BGS-116 WILD RODEUT OATS

A REGISTRATION NO.

8612-110

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

B&G Company 10539 Maybank P.O. Box 20372 Dallas, TX 75220 iel 188,33

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration paust be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFPA section 3(c)(7)(A) provided that you:

- 1. Submit/cite all data required for registration/reregistration of your product under PIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
- 2. Make the labeling changes listed below before you release the menduct thirm n't
 - Add the obrase "PPS registration No. 3612-110."

BEST AVAILABLE COPY

- andlule in the Endangered Species considerations the "Salt Marsh Harvest House."
- 3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.
- If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

SIGNATURE OF APPROVING OFFICIAL

EPA Form 8570-6 (Rev. 5-76)

PREVIOUS EDITION MAY BE USED UNTIL SUPPLY IS EXHAUSTED.

2

A stamped copy of the label is enclosed for your records.

please note the data requirements listed in our letter of October 10, 1986. The due date for the 1-year data for subject product is October 24, 1987. In the near future we will ask you to revise your label to include the species of ground squirrels claimed on your label. The species claimed on your label will be determined by the data submitted to the Agency.

Your Storage Stability data have been reviewed and are acceptable.

William H. Miller

Product Manager (16) Insecticide-Redenticide Branch Registration Division (TS-767)

Enclosures



ACCEPTED with COLLIENTS in EPA Louis 100 15

MAC 1987

Fig. 2 - 110

BEST AVAILABLE REPY

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

CONVULSIVE POISON! Poisonous if swallowed. Do not breathe dust. Do not contaminate feed and foodstuffs. Keep away from children, pets and domestic animals. Wash thoroughly with soap and water after handling and before eating or smoking. Clean clothing should be used daily.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Bests exposed on soil surface may be hazardous to birds and other wildlife. Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

ENDANGERED SPECIES CONSIDERATIONS

NOTICE: The killing of a member of an endangered species during strychnine baiting operations may result in a fine under the Endangered Species Act. Before baiting, the user is advised to contact the Regional U.S. Fish a Wildlife Service (Endangered Species Specialist) or the local Fish a Game Office for specific information on endangered species, Strychnine balts should not be used in the geographic ranges of the following species except under programs and procedures approved by the USEPA: California Condor, San Joaquin Kit Fox, Aleutian Canada Goose, Morro Bay Kangaroo Rat, Gray Wolf and Gristly Bear.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. JACK RABBIT CONTROL

USE RESTRICTIONS: For use against jackrabbits in rangeland, pasture and cropland, and non-agricultural areas (around airports only). Do not use for jackrabbit control in areas occupied by the Utah prairie dog in Garfield, Iron, Kane, Pitte, Sevier and Wayne counties, Utah. Do not use for jackrabbit control in areas occupied by the Masked Bobwhite Quali in Pina and Sonta Cruz counties, Artions. Do not use for jackrabbit control in habitats occupied by Attwater's Greater Prairie Chicken in the following Texas counties: Aransas, Austin, Brazoria, Colorado, Fort Bend, Galveston, Gollad, Refuglo and Victoria.

PREBAITING: Before exposing poisoned bait, it is necessary to probait (offer untreated bait) for 3-5 days to condition the rabbits to feed in locations other than where they are doing damage. Probabiling is most effective when done in the late afternoon.

At a distance of about 100 yards from the field being damaged, place a small handful amount of untreated bait at intervals of 10-15 steps, preferably in trails.

BAITING: In the late afternoon, remove or cover all prebait and replace with a heaping tableapoon of poisoned bait at the same apots where prebait has been accepted.

Do not expose balts in a manner which presents a potential hazard to pels, poultry or livestock. Pick up and there or bury all visible carcasses of animals found in or near treated areas.

GROUND SQUIRREL CONTROL

USE RESTRICTIONS: For use against ground squirrels in rangeland, pasture and cropland, and non-agricultural areas. Do not use for ground squirrel control within 200 yards of prairie dog colonies. Do not use within one mile of a prairie dog colony where the presence of a Black-footed Perret has been confirmed within a five year period. Do not use for ground aquirrel control in areas occupied by Utah prairie dogs in Garfield, Iron, Kane. Piute. Sevier and Wayne counties.

BAITING: Evenly scatter a tablespoon quantity of bait (about 60 baits per pound) on bare ground over \$-3 square fest at side or behind each active burrow. Do not overbalt. Do not expose baits in a manner which presents a potential hazard to pets, poultry or livestock. Do not place bait in plies. Pick up and burn or bury all visible carcasses of animals found in or near treated areas.

RESTRICTED USE PES

For retail sale to and use only by Certified Applicators or person and only for those uses covered by the Certified Applicators Cert





STRYCHNINE TREATED GRAIN

ACTIVE INGREDIENTS

KEEP OUT OF REACH OF CI DANGER - POISON



See Side Panel For Additional Ca

STATEMENT OF PRACTICAL TRE

IF SWALLOWED: Call a physician or Poison Control Center imme have passed since the poison was taken, give 1 or 2 glusses o touching back of throat with finger. Repeat until vomit fluid is quiet, darkened room and keep him warm and quiet. Do not indumouth to an unconscious person.

IF INHALED: Remove victim to Iresh air. Apply artificial respirate ON SKIN: Remove contaminated clothing and wash affected ar IF IN EYES: Flush eyes with plenty of water. Get medical attentions

NOTE TO PHYSICIAN

(A). Administer 1001 OXYGEN by positive pressure to provide a

as possible, despite actures.

(B). Administer ANTICONYULSANT DRUGS intravenously to conmay be difficult or impossible to stop the seizure activity wiprepared to maintain pulmonary ventilation mechanically. Tra seizures are prolonged.

EPA REG.NO. 8812-RRN EPA EST.NO. 8612-TX-1

MANUFACTURED BY BAG COMPANY DALLAS, TX 75220

Label → BGS-118-08

and

and

be

not

ions

l to

fish

1844

veá

Bay

>led

not

rus

ter

na.

for ge.

of

ing

k.

end

dog

in

ies.

und

aits

i កែ

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicators Certification.



STRYCHNINE TREATED GRAIN BAIT

ACTIVE INGREDIENTS:	
Strychnine Alkaloid	0.351
INERT INCREDIENTS	
TOTAL	174 44

KEEP OUT OF REACH OF CHILDREN DANGER - POISON



See Side Panel For Additional Cautions

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED. Call a physician or Poison Control Center immediately. If less than 10 minutes have passed since the poison was taken, give I or 2 glasses of water and induce vomiting by touching back of throat with finger, Repeat until vomit fluid is clear. Have patient lie down in quiet, darkened room and keep him warm and quiet. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED. Remove victim to fresh sir. Apply artificial respiration if indicated.

IF ON SKIN Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES. Flush eyes with plenty of water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN.

(A). Admir.ster 1001 OXYGEN by positive pressure to provide is much pulmonary gas exchange as possible, despite seltures.

(B). Administer ANTICONVULSANT DRUGS intravenously to control convulsions. CAUTION: II may be difficult or impossible to stop the selzure activity without stopping respiration. Be prepared to maintain pulmonary ventilation mechanically. Tracheotomy may be necessary if seizures are prolonged.

EPA REG.NO. 8612-RRN

EPA EST.NO. 8612-TX-1

NET CONTENTS

MANUFACTURED BY BAG COMPANY DALLAS, TX 75220

Label # BGS-116-08

POUNDS

KANGAROO RAT CONTROL

USE RESTRICTIONS: For use against rate in rangeland, pasture and cropland. Do not use in non-agricultural areas. Do not use for kangaroo rat control in areas occupied by the Utah prairie dog in Garfield, Iron, Kane, Plute, Sevier and Wayne countles, Utah. Do not use for kangaroo ret control in areas occupied by the Masked Bobwhite Quail in Pime and Santa Cruz counties, Arizona. Do not apply where the Morro Bay Kangaroo Rat, Dipodomys merriami morroensis, is present in or near a 1.7 square mile of Morro Bay, San Luis Obispo County,

BAITING - Spot Bailing: Evenly acatter 3-4 tablespoons of bait on bare ground near active burrows. Kangaroo rats living within 30 yards of each bailing spot are likely to find the bait. Scatter grain as much as possible to prevent one or two animals from taking all the bait. Do not

Broadcast Balting: Spread evenly by hand, mechanical apreader, or aircraft at the rate of 1/2 to 3/4 pound per swath acre through infested ares. Application of balt should be done in strips 100 feet wide spaced 100 feet apart. Do not expose baits in a manner which presents a likely hazard to pets, poultry, or livestock. Do not place bait in piles. Pick up and burn or bury all visible carcasses of animals found in or near treated areas.

COTTON RAT CONTROL

USE RESTRICTIONS: For use against cotton rats in rangeland, pasture and cropland. Do not use in non-agricultural areas. Do not use for colton rat control in areas occupied by the Mississippi Sandhill Crane in Jackson County, Mississippi. Do not use for cotton rat control in areas occupied by the Cape Sable Sparrow in Collier, Dade and Monroe counties, Florida, Do not use for cotton est control in habitats occupied by Attwater's Greater Prairie Chicken in the following Texas counties: Aranses, Austin, Brazoria, Coloredo, Fort Rend, Galveston, Gollad, Refugio and Victoria.

BAITING - Spot Builting: Lightly scatter tesspoon quantities of bait (about 80 baits per round) in runways near active burrows

Broadcast Balting: Spread bait evenly by hand or mechanical aprevie- through the infested area at the rate of 5-10 pounds per acre, depending on the density of infestation. Breveut tait will fall through most vegetation to the ground surface. Do not apply ball-where trees or glass are wet, or when rain is likely to occur within 24 hours. Do not expose baits in a manner which presents a likely hazard to pets, poultry or livestock. Do not place in piles, Pick un and burn or bury all visible carcasses of animals found in or near treated area",

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store only in original container, in a dry place inacressible to children. pats and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal isn. If these mastes cannot be disposed of by use according to label instructions, contact your State Jes icide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Paper Bag: Completely empty bag by shaking and tapping sides and cottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by state and local authorities. Plastic Containers: Triple rinse (or equivalent) then offer for recycling or reconditioning. or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke,

NOTICE

Seller's guarantee shall be limited to the terms of this label and subject thereto, the buyer sasumes any risks to persons or property arising out of the use or handling and accepts the product on these conditions.

BEST AVAILABLE COPY

1 > 1)

C