

RESTRICTED USE PESTICIDE

For sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

METHYL PARATHION 4 E.C.
Insecticide

Active Ingredients:
O,O-dimethyl O-p-nitrophenyl phosphorothioate* 44.4%
Aromatic Petroleum Solvent 39.5%
Inert Ingredients 16.1%
*methyl parathion

(1 gallon contains 4 lbs. methyl parathion)

DIRECTIONS FOR USE

DISCLAIMER

Staple Cotton Services Association A.A.I. warrants only that the material contained herein conforms to the description on the label and is reasonably fit for the use thereon described when used in accordance with the directions for use.

Any damage arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential, commercial damages such as loss of profits or value, etc.

Staple Cotton Services Association A.A.I. makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

BUYER assumes the risk of any use contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable by Staple Cotton Services Association A.A.I.

This product is to be applied by experienced personnel using ground or aircraft equipment. The hazards and precautions on this container are equally applicable to it after dilution with water for spray application. Pour specified amount of this product into nearby filled spray tank. Add balance of water to fill tank. Mix thoroughly either by means of a tank agitator or pump by-pass.

For best results, thoroughly cover all surfaces to be treated with spray. Consult your State Experiment Station or State Agricultural Extension Service for specific recommendations regarding application, dosage and timing of sprays. When applying by ground equipment, use specified dosage in not less than 10 gallons of water per acre; by aircraft equipment in 2-5 gallons of water per acre.

PEACH AND APRICOT: European red mite and green peach aphid - Apply 1/2 pt. per 100 gallons of water. Apply twice at 5-day intervals for control of European red mite. Do not apply within 14 days of harvest.

VEGETABLE AND FIELD CROPS: States are given in terms of plants of METHYL PARATHION 4 E.C. per acre in addition water for good coverage. The number shown in parentheses () indicates the number of days which should be allowed between last application of harvest or grading. Unless otherwise specified, this product will control aphids, armyworms (up to 3rd instar), Dog, Fleas, Fleas, Fleas and Fleas on the following crops:


DANGER
Keep out of reach of children

POISON
See this panel for antidote and precautions



CAUTION

This product is highly toxic to humans. Keep out of reach of children. Do not apply near water. Do not apply to drift from area. Do not contain equipment, or



Staple Cotton

DANGER - POISON - PRECAUTIONS

NOT FOR HOME USE



CAN KILL YOU IF SWALLOWED

This product can kill you if swallowed even in small amounts; spray mist or dust may be fatal if swallowed.



CAN KILL YOU BY SKIN CONTACT

This product can kill you if touched by hands or splashed on skin, in eyes or on clothing (liquid goes through clothes).



CAN KILL YOU IF BREATHED

This product can kill you if vapors, spray mist or dust are breathed.



Physicians' Note: Warning symptoms include weakness, headache, tightness in the chest, blurred vision, uncoördinate pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

Treatment: Give atropine, preferably by injection, grains 1/300 three or more tablets at once and every hour as required until pupils dilate. Thirty or more tablets may be required during the first 24 hours. Never give morphine. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until cholinesterase regeneration has been allowed as determined by blood tests. I-PAM is also antidotal and may be administered in conjunction with atropine.

NOTE: Protect from temperatures below 32° F.

Do Not Get On Skin, On Clothing
Wear clean, heavy rubber gloves and clean waterproof or freshly laundered protective clothing (coveralls, cap, etc.) Clothing must not get wet through. Destroy and replace gloves frequently.
Wash thoroughly with soap and warm water before eating or smoking. Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and hot water before re-use.
In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water.

Do Not Breathe Vapor Or Spray Mist
Wear a mask or respirator of a type specified by the U. S. Bureau of Mines for methyl parathion protection. Wear goggles. Airplane pilots should wear full-face canister-type mask.
If handled indoors, provide mechanical exhaust ventilation.
Remove unprotected persons and domestic animals from operating and adjacent areas where insecticide may drift, prohibit re-entry until drift and vapor dissipate.

Do Not Store Near Food Or Food Products
Cover spillage with an absorbent such as a soda ash, lime, clay or sawdust. Sweep up and bury. Wash area thoroughly with a full-strength liquid household chlorine bleach.
Apply this product only as specified on this label.
DECONTAMINATE EMPTY CONTAINERS AS FOLLOWS: Add 1 pint of liquid chlorine bleach, clean container and shake for 5 minutes. Allow to stand for 15 minutes. Flush with water and mutilate to prevent re-use. When decontaminating spills or empty containers, observe personal safety precautions listed above.

Atropine is an antidote - Consult physician for emergency supply.

FIRST AID TREATMENT

Call A Physician At Once In All Cases Of Suspected Methyl Parathion Poisoning. If symptoms or signs of poisoning (See Physicians' Note) include blurred vision, abdominal cramps and tightness in the chest, don't wait for a doctor but give two atropine tablets (each 1/300 gr.) at once.

Remove patient immediately from the area where methyl parathion is present. Remove contaminated clothing and wash the skin clean with plenty of soap and water to remove all traces of methyl parathion. If available, induce vomiting by giving warm, salty or soapy water. Have patient lie down and keep quiet.

Never Give Anything By Mouth To An Unconscious Person