

Yellow Copy

File # 8660-51

10/2/92

OCT 2 1992

Mr. Daniel Paradiso, Jr.
The Andersons
d/b/a Parker Fertilizer Company
201 4th Street
Sylacauga, AL 35150

Dear Mr. Paradiso:

Subject: Revised Labeling
Vertagreen 50% Methoxychlor Wettable Powder
EPA Registration No. 8660-51
Your Application Dated July 24, 1992

The labeling for the product referred to above submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable subject to the comments listed below. A stamped copy of the label is enclosed for your records.

1. The term "For Professional/Commercial Use Only" must be deleted when storage and disposal directions other than for the homeowner are used for this product.

Sincerely yours,

DHE

Dennis H. Edwards, Jr.
Product Manager 19
Insecticide-Rodenticide Branch
Registration Division (H-7505C)

236

8660-51
1 of 5

(FRONT PANEL)

VERTAGREEN®

50% METHOXYCHLOR

WETTABLE POWDER

ACCEPTED
with **COMMENTS**
in EPA Letter Dated

OCT 2 1992

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
8660-51

ACTIVE INGREDIENTS:

Methoxychlor: 1,1,1-trichloro-2,2-bis
(p-methoxyphenyl) ethane

50.00%

INERT INGREDIENTS:

50.00%

Total

100.00%

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT


IF ON SKIN: Wash skin with plenty of soap and water.

IF IN EYES: Flush with water for 15 minutes and get medical attention.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET CONTENTS 8 OZ.

This Container Will Make Up To 17 Gallons of Spray



(BACK PANEL)

NET CONTENTS 8 OZ.

This Container Will Make Up To 17 Gallons of Spray

VERTAGREEN®

50% METHOXYCHLOR

WETTABLE POWDER

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes, or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Aerial application over such sites is permissible only when they are not visible from above the tree canopy. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

Do not graze or feed treated pea vines or pea hay to livestock.
Do not graze or feed treated cowpea vines or cowpea hay to livestock.
Do not graze or feed treated soybean hay or straw to livestock.
Do not feed treated corn to livestock.

WORKER PROTECTION

Do not enter or allow entry into treated areas until sprays have dried/dusts have settled/vapors have dispersed, as applicable, to perform hand labor tasks. A person may enter the area to perform other tasks only if the person is wearing the personal protection equipment listed on the label for a pesticide handler.

Do not apply this product in a way that will contact unprotected workers, either directly or through drift. Only protected handlers wearing long pants, long-sleeved shirts, socks, shoes and chemical resistant gloves may be in the area during application.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

INSECT CONTROL ON PLANTS: Mix 2 to 3 tbsp. per gallon of water (2 to 3 lbs. per 100 gals.). Apply thoroughly to cover all surfaces of foliage and fruit. Repeat at 7 to 14 day intervals as needed, to control the following insect, but do not exceed pounds of the product per acre indicated in parenthesis after each crop. do not apply later than 7 days (*3 days, **14 days, *** 21 days) before harvest.


VEGETABLES: Asparagus* (4-1/2), Beans (4-1/2), Beets** (4-1/2), Blackeyed Peas (6), Broccoli** (4-1/2), Brussels Sprouts** (4-1/2), Cabbage (4-1/2), Carrots** (4-1/2), Cauliflower (4-1/2), Collards** (4-1/2), Corn (4-1/2), Cucumbers** (4-1/2), Eggplant (4-1/2), Kale** (4-1/2), Kohlrabi (4-1/2), Lettuce** (4-1/2), Melons (4-1/2), Peas (4-1/2), Peppers (4-1/2), Potatoes (4-1/2), Pumpkin (4-1/2), Radish (4-1/2), Rutabaga (6), Spinach** (4-1/2), Squash (4-1/2), Summer Squash (4-1/2), Tomatoes (4-1/2), and Turnips** (4-1/2) for control of Asparagus Beetle, Bean Leaf Beetle, Blister Beetle, Cabbageworm, Colorado Potato Beetle, Cucumber Beetle, Fall Armyworm, Flea Beetles, Japanese Beetle, Leafhoppers, Mexican Bean Beetle, Pea Weevil, Squash Vine Borer and Webworm.

FLOWERS AND ORNAMENTALS: Including Chrysanthemums, Dahlias, Roses, Shrubs, Shade trees and most other flowers for control of Blister Beetle, Cankerworm, cucumber Beetle, Flea Beetle, Fleahoppers, Rose Chafer, Rose slug (Sawfly), Tent Caterpillar, and many other insects.

TREE FRUITS: Apples (30), Apricots*** (12), Cherries (16), Nectarines*** (4), Peaches*** (12), Pears (16), Plums (16), Prunes (16), and Quinces (12) for control of Apple Maggot, Cankerworm, Cherry Fruit Fly, Cherry Fruitworm, Codling Moth, Japanese Beetle, Leafhoppers, Pear Slug, Plum Curculio and Tent Caterpillar.

SMALL FRUITS: Blackberries** (6), Blueberries** (4-1/2), Cranberries** (4-1/2), Currants** (6), Gooseberries** (6), Grapes** (12), Loganberries** (6), Raspberries** (6), and Strawberries** (4-1/2), for control of Cherry Fruitworm, Cranberry Fruitworm, Grape Berry Moth, Grape Leaf Skeletonizer, Japanese Beetle, Leafhoppers, Plum Curculio, Rose Chafer, San Jose Scale (Crawlers), Spittlebug and Strawberry Weevil.

FIELD AND FORAGE CROPS: Alfalfa (3), Cowpeas (3), Clover (3), Peanuts (3), Soy Beans (3), and foliage grasses (3) for control Alfalfa Caterpillar, Alfalfa Weevil Larvae, Blister Beetle, Armyworm, Cowpea Curculio, Clover Leaf Weevil, Fall Armyworm, Flea Beetles, Japanese Beetle, Leafhoppers, Pea Weevil, Soybean Caterpillar, Spittlebug and Velvet Bean Caterpillar. Do not apply later than 7 days before harvest or grazing.



INSECT CONTROL ON ANIMALS: Spray or dip cattle, hogs and sheep to control Horn Flies, Lice, Ticks (Keds) and Fleas, using 2 to 4 lbs. per 25 gals. of water. Make sure animals are wetted to the skin; usually, one treatment is effective for about 3 weeks. Dust dairy animals (for Horn Fly control) by using a single heaping tablespoonful per animal, sprinkled over the animal and rubbed into the hair or wool. On cattle, apply to poll, neck, back and upper sides, which will last two to three weeks. On Sheep, if dusting is desirable because of weather use 1 tbs. per animal at shearing time.

INSECT CONTROL IN BUILDINGS:

FARM BUILDINGS: To control Stable Flies, House Flies and Fleas, mix 2 lbs. in 2-1/2 gals. of water, and apply 1 gal. of mixture to 500 sq. ft. of surface; treating walls, floors, cracks, crevices, baseboards, bedding, coops, stalls, and pens. Such heavy deposits usually are effective for 3-4 weeks, unless removed by weathering or other means. Proper sanitary measures should also be followed, including spraying or removal of breeding sources such as manure in and around buildings. CAUTION: When treating dairy barns, dairy animals should be removed, and feed and water protected during the treatment.

GRAIN STORAGE BINS: Spray empty bins thoroughly (after thorough cleaning) before storage of barley, corn, oats, rice, rye, sorghum (milo) or wheat, with spray composed of 2 lbs. mixed in 5 gals. of water at the rate of 1 gal. of spray per 500 sq. ft. of surface. This treatment will aid in control of Cadelle, Flat Grain Beetle, Saw-Toothed Grain Beetle, Lesser Grain Borer, Red Flour Beetle, Long-Headed Flour Beetle, Rice Weevil, Granary Weevil, Confused Flour Beetle, Foreign Grain Beetle and Hairy Fungus Beetle.

STORAGE AND DISPOSAL

FOR HOMEOWNER USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

FOR PROFESSIONAL/COMMERCIAL USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.**

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use condition. **FOLLOW DIRECTIONS CAREFULLY.** Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured By:

**PARKER FERTILIZER CO.
201 4TH STREET
SYLACAUGA, AL 35150**

Date Printed: 07/23/92

Date Approved:

Supersedes:

Label Change:

