



Ring Around Brand

READ LABEL BEFORE USING!

NOT FOR HOME USE



DA N G E R

Keep Out of Reach of Children

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

CONTAINER DISPOSAL

See instructions for Container disposal on this label

Ring Around Brand

EPN EMULSIFIABLE INSECTICIDE

Manufactured By RING AROUND PRODUCTS, INC. MONTGOMERY, ALABAMA

NOTE TO PHYSICIAN

Antidote--administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as convulsions are overcome. Repeat at 5 to 10 minute intervals until signs of atropinization appear. 2 PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS

EPN is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement. VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS. TREAT SYMPTOMATICALLY.

DIRECTION FOR USE

Ring Around Brand EPN Emulsifiable Insecticide contains 2 lbs. of EPN for use on cotton insects listed.

Apply at the following rates in sufficient water to obtain thorough uniform plant coverage; for aircraft application, use 3 to 10 gallons of water per acre.

Cotton fleahopper, cotton leafworm, thrips, yellow-striped armyworm, spider mites, (certain species): Apply 1-1/4 to 1-3/4 pints per acre. Begin application when species appear. Repeat at 5-day intervals as needed.

Boll Weevil: Apply 2 to 3 pints per acre. Begin when cotton is fruiting and 10% to 25% of squares are punctured. Repeat at 5 day intervals as needed.

Bollworm, tobacco hornworm. Apply 1 pint per acre. Begin when eggs and 4 to 6 worms per 100 terminals are found. Repeat at 5-day intervals as needed.

CAUTION: Do not apply within 3 days of hand picking. Workers entering fields within 24 hours after application should wear protective clothing. Do not feed forage or grain to livestock.

USE ONLY IN ACCORDANCE WITH INSTRUCTIONS ON THIS LABEL

DO NOT REUSE EMPTY CONTAINERS FOR ANY PURPOSE. Decontaminate and destroy all empty containers. To decontaminate, proceed as follows. Completely empty contents. For containers of 1 gallon or less, use 1 quart of water and 4 tablespoons of household detergent; for containers larger than 1 gallon, use 5 gallons of water, 2 pounds caustic soda (household lye), and 1 cupful detergent as rinse solution. Pour into container and carefully rotate to wet all inner surfaces with the solution. Bury the unused chemical and rinse solution at least 18 inches deep in an isolated location away from water supplies. Break or punch holes in top and bottom of container and bury deeply in an isolated location.

Δ 100 100

100

100
100

100

100
100

100 100 100 100 100