1 7 11

FEB | 1995

FE₅

Ms. Michelle Bell
The Andersons Lawn Fertilizer Division
DBA Pree Flow Fertilizer
P.O. Box 115
Maumee, Oh 43537

Dear Ms. pell:

Subject: The Anderson: Fertilizer with 0.87% Team/0.58% Dursoan EPA Registration No. 9195-102
Application Dated December 30, 1394, Request To
Amend Labeling by Adding APS Exclusionary Statement and Using Split Labeling

The proposed labeling submitted with your upplication dated March 25, 1994 has been reviewed and round acceptable under the Federal Insecticide, Fungicide and Rodenticide Act (rIRA) as amended, provided that:

- 1. Add the same exclusionary statement to the "homeowner ose only" labeling as the one you added to the "Commercial" Use Only" labeling submitted with this application.
- Submit one (1) copy of the final printed labels before releasing the product for shipment under the subject laceling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, Section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copies of the labeling are enclosed for your records.

Sincerely yours,

BEST AVAILABLE COPY

Joanne I. Biller Product Manager (23) rungicide-Herbicide Branch Registration Division (7565C)

cholosure

E.Wilson: Diskette ABC17:02-01-35

(FRONT PANEL)

HOMEOWNER USE ONLY

THE ANDERSONS FERTILIZER WITH 0.87% TEAM /0.58% DURSHAN

(Logo)

ACTIVE INGREDIENTS:

Total	100.00%
INERT INGREDIENTS	. 98.55+
Chlorpyrifos [0,0-diethyl 0-(2,4,6-trichloro-2-pyridyl)phosphorothioate]	. 0.50+
Trifluralin: alpha,alpha,alpha-trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine	0.289
N-butyl-N-ethyl-~,~,~-trifluoro-2,6-dinitro-p-toliudine	0.395

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional precautionary statements

Net Weight 50 lbs. (22.7 kg.)

Treats Up to 12,500 Sq. Ft.

BEST AVAILABLE COPY

EPA Reg. No. 9198-102

ACCEPTED
with Comments
in EPA Lames Is to de

which is the second

Under the Federal Decrease of Foregoids and Holographics of the tipe of the specific strength and on LPV the $\sim 1.016 \, \mathrm{GeV} - 10.2$

EPA Est. 9156--0H-1

9199 102H Page 2 of 5

(BACK PANEL)

HOMEOWNER USE CNLY

Net Weight 50 lbs. (22.7 kg.)

Treats Up To 12,500 Sq. Ft.

THE ANDERSONS FERTILIZER WITH 0.87% TEAM /0.58% DURSBAN

Guaranteed Analysis

Total Nitrogen (N)	25•
1.2: Ammoniacal Nitrogen	
17.8. Urea Nitrogen	
6.0+ Coated Slow Release Urea Nitrogen	
Available Phosphoric Acid (P2O5)	٠,٠
Soluble Potash (K2O)	
Plant Mutrients Derived From: Ammoniated Encaphates, Urea, Sultur	Coates

Urea, and Muriate of Potash.

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

CAUTION

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Avoid contact with eyes. If in eyes, flush with plenty of water. Harmful if swallowed, inhaled or absorbed through the skin. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with scap and water after handling. Remove contaminated clothing and wash refere reuse.

(BACK PANEL)

HOMEOWNER USE ONLY

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Fold eyelids open and flush with a steady, gentle stream of

water for 15 minutes. Call a physician.

IF SWALLOWED: Call a physician or Poison Control Center. Drink one or two

glasses of water and induce vomiting by touching finger to the back of throat. Never give anything by mouth to an unconscious

or convulsing person.

IF ON SKIN: Wash with soap and plenty of water. Get medical attention if

if irritation persists.

IF INHALED: If breathing difficulty occurs, move individual to fresh air

and get medical attention.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase innibitor. Treat

symptomatically. Atropine is an antidote by injection

only.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish, birds, and other wildlife. Do not apply to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate spills. Do not contaminate water when cleaning of equipment washwaters.

DIRECTIONS FOR USE

It is in violation of Federal law to use this product in a manner inconsistent with its labeling.

The Andersons Fertilizer With 0.87% Team/0.58% Dursban is a combination of growth promoting fertilizer ingredients, a preemergent herbicide, and a turf insecticide. It is designed to control smooth and hairy crabgrass, goosegrass (silver crabgrass or crowsfoot), barnyard grass (watergrass), green and yellow foxtail, and annual bluegrass (Poa Annua), as well as ants, armyworms, billbugs, brown dog ticks, chinch bugs, crickets, cutworms, carwigs, grasshoppers, and sod webworms.

When applied as directed, the product can be used on the following turfs: Kentucky Bluegrass, perennial ryegrasses, centipedegrass, fescue, zoyslagrass, bermudagrass, St. Augustinegrass, and bahiagrass. The product will not harm nearby established ornamental trees and shrubs.

(BACK PANEL)

HOMEOWNER USE ONLY

APPLICATION DIRECTIONS

For preemergence weed control and early insect control, apply 4 pounds per thousand square feet in the spring one to two weeks prior to the enset of conditions favorable for annual weed grass seed germination. A second application should be made eight to ten weeks after the first to provide season control of these annual grass species and to provide a second insect kill.

USE PRECAUTIONS

Apply The Andersons Fertilizer with 0.87% Team/0.58% Dursban to the turf prior to the germination of weed grasses since The Andersons Fertilizer With 0.87% Team/0.58% Dursban will not control established weeds. The Andersons Fertilizer With 0.87% Team/0.58% Dursban should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application of The Andersons Fertilizer With 0.87% Team/0.58% Dursban should not be used on putting greens. The Andersons Fertilizer With 0.87% Team/0.58% Dursban should not be used on putting dreens. The Andersons Fertilizer With 0.87% Team/0.58% Dursban should not be used on putting dreens. The Andersons Fertilizer With 0.87% Team/0.58% Dursban should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass of Centipedegrass until these turfs are well established.

RESEEDING

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applying The Andersons Fertilizer With 0.87% Team/0.58% Dursban. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization be followed. For satisfactory reseeding results after using The Andersons Fertilizer With 0.87% Team/0.58% Dursban the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as Rogers Aero Seeder) should be employed. When using The Andersons Fertilizer With 0.87% Team/0.58% Dursban reseeding should be delayed 12 to 16 weeks after application.

SUGGESTED SPREADER SETTINGS

(This section will be completed for final labeling)

9198-102H Page 5 of 5

(BACK PANEL)

HOMEOWNER USE ONLY

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash. CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WAPPANTY EXPRESS OF IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLIOW DIRECTIONS CAREFULLY. Timing and method of application, weather and turf conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer.

Manufactured By: THE ANDERSONS

Lawn Fertilizer Division

P.O. Box 119 Maumee, OH 43537

Date Printed: 11/02/94

Date Approved: Supersedes: Label Change:

(FRONT PANEL)

COMMERCIAL USE ONLY

THE ANDERSONS FERTILIZER WITH 0.87% TEAM /0.58% DURSBAN

(Logo)

ACTIVE INGREDIENTS:

N-butyl-N-ethyl-~,~,~-trifluoro-2,6-dinitro-p-toliudine	•	•	•	•		0.59+
Trifluralin: alpha,alpha,alpha-triflucro-2,6-dinitro-N,N-dipropyl-p-toluidine						0.28%
Chlorpyrifos [0,0-diethyl 0-(2,4,6-trichloro-2-pyridyl)phosphorothioate]	•				•	0.58•
INERT INGREDIENTS		٠				98.55*
Total						100.00•

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional precautionary statements

Net Weight 50 lbs. (22.7 kg.)

Treats Up to 12,500 Sq. Ft.

BEST AVAILABLE COPY

EPA Reg. No. 9198-102

ACCEPTED
with COMMENTS
in EPA Letter Dated:

FEB | 1905

Under the Federal In countle, Funziole, and Rodents, no Act as amended, for the perbude registered under EPA Reg. No.

EPA Est. 9198-OH-1

9198-OH-1

, ,

(BACK PANEL)

COMMERCIAL USE ONLY

Net Weight 50 lbs. (22.7 kg.)
Treats Up To 12,500 Sq. Ft.

THE ANDERSONS FERTILIZER WITH 0.87% TEAM /0.58% DURSBAN

Guaranteed Analysis

Total Nitrogen (N)	25.
1.2% Ammoniacal Nitrogen	
17.8% Urea Nitrogen	
6.0 Coated Slow Release Urea Nitrogen	
Available Phosphoric Acid (P2O5)	3 •
Soluble Potash (K2O)	÷.

Plant Nutrients Derived From: Ammoniated Phosphates, Urea, Sulfur Coated Urea, and Muriate of Potash.

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

CAUTION

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Avoid contact with eyes. If in eyes, flush with plenty of water. Harmful if swallowed, inhaled or absorbed through the skin. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

S' TEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Call a physician.

water for 15 minutes. Call a physician.

IF SWALLOWED: Call a physician or Poison Control Center. Drink one of two glasses of water and induce vomiting by touching finger to the back of throat. Never give anything by mouth to an unconscious

unconscious or convulsing person.

IF ON SKIN: Wash with soap and plenty of water. Get medical attention if

if irritation persists.

BEST AVAILABLE COPY

9198-102C Page 3 of 5

(BACK PANEL)

COMMERCIAL USE ONLY

IF INHALED: If breathing difficulty occurs, move individual to fresh air

and get medical attention.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat

symptomatically. Atropine is an antidote by injection

only.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish, birds, and other wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate spills. Do not contaminate water when cleaning or equipment washwaters.

DIRECTIONS FOR USE

It is in violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

The Andersons Fertilizer With 0.87% Team/0.58% Dursban is a combination of growth promoting fertilizer ingredients, a preemergent herbicide, and a turf insecticide. It is designed to control smooth and hairy crabgrass, goosegrass (silver crabgrass or crowsfoot), barnyard grass (watergrass), green and yellow foxtail, and annual bluegrass (Poa Annua), as well as ants, armyworms, billbugs, brown dog ticks, chinch bugs, crickets, cutworms, earwigs, grasshoppers, and sod webworms.

When applied as directed, the product can be used on the following turfs: Kentucky Bluegrass, perennial ryegrasses, centipedegrass, fescue, zoysiagrass, bermudagrass, St. Augustinegrass, and bahiagrass. The product will not harm nearby established ornamental trees and shrubs.

APPLICATION DIRECTIONS

For preemergence weed control and early insect control, apply 4 pounds per thousand square feet in the spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination. A second application should be made eight to ten weeks after the first to provide season control of these annual grass species and to provide a second insect kill.

BEST AVAILABLE COPY

9195 1 200 Page 4 of 5

1. 7 11

(BACK PANEL)

COMMERCIAL USE ONLY

USE PRECAUTIONS

Apply The Andersons Fertilizer With 0.87* Team/0.58* Dursban to the turf prior to the germination of weed grasses since The Andersons Fertilizer With 0.87* Team/0.58% Dursban will not control established weeds. The Andersons Fertilizer With 0.87% Team/0.58* Dursban should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application of The Andersons Fertilizer With 0.87% Team/0.58% Dursban should not be used on putting greens. The Andersons Fertilizer With 0.87% Team/0.58% Dursban should not be used on dichondra. The Andersons Fertilizer With 0.87% Team/0.58% Dursban should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.

RESEEDING

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applying The Andersons Fertilizer With 0.87. Team/0.58. Dursban. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization be followed. For satisfactory reseeding results after using The Andersons Fertilizer With 0.87. Team/0.58. Dursban the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as Rogers Aero Seeder) should be employed. When using The Andersons Fertilizer With 0.87. Team/0.58. Dursban reseeding should be delayed 12 to 16 weeks after application.

SUGGESTED SPREADER SETTINGS

(This section will be completed for final labeling)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfit or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

9198-102C Page 5 of 5

(BACK PANEL)

COMMERCIAL USE ONLY

WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and turf conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer.

Manufactured By: THE ANDERSONS

Lawn Fertilizer Division

P.O. Box 119 Maumee, OH 43537

Date Printed: 11/02/94

Date Approved: Supersedes: Label Change:

•

• • •