

Net Weight 50 Pounds

BERRIEN

ACCEPTED

JUN 16 1967

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 9275-20

5% SEVIN-75% SULPHUR DUST

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-Methylcarbamate) 5.00%

Sulphur, as elemental 75.00%

INERT INGREDIENTS: 20.00%

TOTAL 100.00%

(R) Trademark of Union Carbide Corp. for 1-naphthyl N-methylcarbamate.

Keep Out of Reach of Children

CAUTION: Harmful if Swallowed or Inhaled. Do not take internally. Avoid breathing dust. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs. This product is toxic to honeybees. Avoid use during flowering and pollination period.

DIRECTIONS

VEGETABLES

APPLICATION: Begin applications when insect or damage is first observed and repeat on a regular 5 to 7 day schedule or as necessary for good insect control.

BEANS: To control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Leafhoppers, Flea Beetles, Japanese Beetles, Corn Earworm and Rust — apply at the rate of 20 pounds per acre. To control Lygus Bug, Stink Bug and Lima Bean Pod Borer — apply 40 pounds per acre.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI: To control Flea Beetle and Harlequin Bug — apply at the rate of 20 pounds per acre. To control Corn Earworm, Armyworm, Cabbage Caterpillars and small Cabbage Loopers — apply 20 to 40 pounds per acre.

OKRA: To control Corn Earworm and Stink Bugs — apply at the rate of 20 to 40 lbs. per acre.

TOMATO, EGGPLANT, PEPPER: To control Tomato Fruitworm, Fall Armyworm, Flea Beetles and Leafhoppers — apply at the rate of 20 pounds per acre. For control of Tomato Hornworm, European Corn Borer and Stink Bugs — apply at the rate of 20 to 40 pounds per acre.

POTATO: To control Colorado Potato Beetle, Leafhoppers, and Flea Beetles, apply at 20 pounds and for Lygus Bug, apply at 40 pounds per acre.

INTERVAL BEFORE HARVEST: This product can be used to and including the day of harvest on all vegetables listed except Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kohlrabi. When applied to these vegetables allow 3 days between the last application and harvest.

CAUTION: Do not apply this product to Cucumbers, Squash or Melons because sulphur injury will result.

FORAGE CROPS

APPLICATION: Apply when insects or signs of damage appear and repeat 7 to 14 days later if necessary.

PEANUTS. To control Mexican Bean Beetle, Blister Beetle, Hoppers, Three-Cornered Alfalfa Hopper, Cucumber Beetle, Flea Beetle, Japanese Beetle, Alfalfa Caterpillar, Velvet Bean Caterpillar, Leaf Spot — apply at the rate of 20 pounds per acre. To control Armyworms, Webworms, Corn Earworm, Stink Bugs, and Grasshoppers — apply at the rate of 20 to 30 pounds per acre. Application may be made up to and including the day of harvest. Some leaf injury may occur if applications are made when tender foliage is wet.

ALFALFA, CLOVERS, SOYBEANS, COWPEAS: To control the insects listed under peanuts apply at the rate of 20 to 30 pounds per acre. Some leaf injury may occur if applications are made when foliage is wet.

CORN: (Forage and Fodder): To control Corn Earworm, European Corn Borer, Fall Armyworm, Saw Beetles and Japanese Beetles, apply at the rate of 20 to 40 pounds per acre. Applications may be made up to and including 7 days of harvest.

SORGHUM: (Grain and Forage): To control Armyworm, Webworm, Corn Earworm, Stink Bugs, and Grasshoppers, apply at the rate of 20 to 30 pounds per acre. Allow 21 days between last application and harvest of grain, no time limit on sorghum forage.