

Sunniland Corporation
c/o Mr. H. R. McLane
7210 Red Road, Suite 206
Miami, FL 33143-5321

FEB 16 1993

Dear Mr. McLane:

Subject: Amendment - Change to "Caution" Formula
Sunniland 25% Diazinon Liquid Concentrate
EPA Reg. No. 9404-65
Your label submission dated July 30, 1992
and CSF submission dated January 26, 1993

The amendment referred to above, submitted in connection with a registration under the Federal Insecticide, Fungicide, and Rodenticide Act is acceptable provided the following revisions are made prior to release for shipment under the amended labeling:

- 1) On page 2 of the booklet label, revise the end of item "5" in the "Notes" under the heading "General Information" to read: "... in greenhouses.", rather than "... in green houses."
- 2) On page 5 of the booklet label, in the first sentence under the heading "Vegetables", use either "this product" or the correct name for your product, rather than "D.z.n Lawn and Garden Insect Control".

A stamped copy of the label is enclosed for your records. The replacement Confidential Formula (CSF) dated January 26, 1993 is acceptable and has been added to this product's registration file.

NOTE - The label and associated Confidential Statement of Formula accepted at this time supersede all previous labels and formula statements on file for this product.

Sincerely,

George T. LaRocca
Product Manager (13)
Insecticide Rodenticide Branch
Registration Division (H7505C)

2 of 13

Sunniland®

25% DIAZINON LIQUID CONCENTRATE

**THIS PRODUCT MUST NOT BE USED
ON GOLF COURSES AND SOD FARMS.**

ACCEPTED
with COMMENTS
in EPA Letter Dated

**FOR OUTDOOR USE ON THE HOME LAWN
AND IN THE HOME GARDEN ONLY.** FEB 16 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

**A SPRAY CONCENTRATE
CONTAINS DIAZINON INSECTICIDE**

9404-65

**Protects - Lawns, Vegetables, Fruits & Nuts, Roses, Flowers,
Trees, Shrubs**

**Controls - Ants, Aphids, Chinch Bugs, Cutworms, **Fire Ants,
Flea Beetles, Grubs, Mites, Sod Webworms and
many other insects. **aids in control.**

ACTIVE INGREDIENT:

**DIAZINON: [O,O-Diethyl O-(2-isopropyl-6-methyl-
4-pyrimidinyl) phosphorothioate] - 25.00%**

INERT INGREDIENTS:* - - - - - 75.00%
TOTAL 100.00%

***Contains Aromatic Petroleum Distillates
KEEP OUT OF REACH OF CHILDREN
CAUTION**

See end of booklet for additional precautionary statements

EPA REG. NO. 9404-65

EPA EST. NO. 9404-FL-1

NET CONTENTS 1 PINT;(1 QUART; 1 GALLON)

SUNNILAND CORPORATION

SANFORD, FL 32771

3 of 13

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

NOTICE: This product conforms to its chemical description and is reasonably fit for the purpose stated on the label, when used in accordance with directions under normal conditions of use. Manufacturer is not responsible for the use of this product contrary to the label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the manufacturer and/or seller and the buyer assumes the risk of any such use. 06-30-82 rev.

DIRECTIONS FOR USE

Violation of federal law to use this product in a manner inconsistent with its labeling. **FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

General Information

This Product is an emulsifiable concentrate product which must be diluted with water before application. Refer to the section How to Mix and Apply. This product kills many outdoor insects which damage lawns, roses, flowers, trees, shrubs, vegetables, fruits and nuts, and insects which annoy pets and people. Contains diazinon insecticide which provides positive control of many "hard-to-kill" sucking and chewing insects. Works fast and applications should be repeated only as directed to maintain control.

Notes: (1) USE CONTENTS WITHIN 6 MONTHS AFTER OPENING. SEE STORAGE AND DISPOSAL DIRECTIONS. (2) Do not use in the home. (3) Do not use on ferns, pansy, hibiscus, papaya, pine, gardenia or orchids since injury may occur. (4) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties. (5) Do not apply to food crops grown in green houses.

Precautions: (1) Wear a long-sleeved shirt and long-legged pants when spraying. (2) Spray with the wind to your back. (3) Do not spray on windy days. (4) If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water, and launder clothing before wearing again.

How to Mix and Apply

Step #1: Determine size of spraying job. For lawns and vegetables, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear. Applications should be repeated only as directed to maintain control.

Step #2: Use a clean sprayer. Carefully measure and mix as suggested in the rate chart.

Step #3: Spray as directed in rate chart. Thorough coverage is important. Flush sprayer with clean water after each use.

See chart for complete list of use sites, crops, and insects controlled. Use the rate given in the chart for the type of insect you want to control on the site or crop you are spraying.

Measurement Conversions

- 1 tablespoon = 3 teaspoons
- 1 fl. oz. = 6 teaspoons or 2 tablespoons
- 6 fl. oz. = ¾ cup
- 1 ½ tablespoons in 3 gals. of water = 1 ½ teaspoons in 1 gal. of water
- 2 tablespoons in 3 gals. of water = 2 teaspoons in 1 gal. of water

Lawn Pest Control

Precautions: Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as directed on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid over-

lapping of sprays. Where irrigation (watering) is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), water lawn with at least ¼ inch of water immediately after spraying this product; however, stop watering before puddling occurs.

Use on Bent, Bermuda, Bluegrass, Dichondra, Fescue, Menon and St. Augustinegrass. To control pests listed below, apply recommended rates.

Pests	Rate		Remarks
	This Product	Water	
Lawn Chinch Bugs	1 ½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Thoroughly wet down grass a few hours before applying. In the North, if Chinch Bugs are the only problem, a lower rate of application can be used: 3 teaspoons in 3 gals. of water covers 125 sq. ft. or 4 fl. oz. in 24 gals. of water per 1,000 sq. ft.
Ants, Armyworms, Bermudagrass Mites, Clover Mites, Springtails (Colembola), Crickets, Cutworms, Digger Wasps, Earwigs, Frit Flies, Lawn Billbugs, Leafhoppers, Sod Weevils (Lawn Moths), Sowbugs (Pillbugs)	1 ½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Thoroughly wet down grass a few hours before applying. For ants and digger wasps, spot spray ant hills and wasp ground nest openings. For digger wasps, apply at dusk, when wasps are not active. Billbug control: Apply when activity is first observed or when chewed or brown grass is present indicating damage from the insect. For frit flies, mow grass and water thoroughly before treatment, but delay application until grass is dry.
Brown Dog Ticks, Chiggers, Fleas	1 ½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	NOTE: Do not apply this product to animals. For brown dog ticks, spray grass and (if possible) shrubs v. particularly near house.
Hyperodes Weevil	1 ½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Treat problem areas in mid-afternoon and apply in mid-day. Water grass thoroughly after application.
Milipedes	1 ½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Thoroughly wet grass a few hours before spraying.
Rhodesgrass Scales	1 ½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Apply when crawlers first emerge.
White Grubs or Japanese Beetle, European Crane, Southern Crane	1 ½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Apply any time between late July and early October. Water grass thoroughly after application.

BEST AVAILABLE COPY

4713

Nuisance Pests Outside the Home

<p>Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Sowbugs (Pillbugs), Spiders, Springtails (Collembola)</p>	<p>Mix 6 fl. oz. in 3 gals. of water and thoroughly spray window and door screens, sills, foundations, porch and patio walls, entranceways, walks, garbage cans, tree trunks, and into cracks and other places where insects can hide.</p> <p>To prevent entry into houses, spray a two-foot band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 feet. Repeat application as necessary.</p> <p>Do not treat animals with this product. Do not treat plants (other than tree trunks) at this nuisance pest rate.</p>
--	---

<p>Fire Ants</p>	<p>To aid in the control of fire ants in home lawns and other outdoor home recreation areas, apply 2 tablespoons in a minimum of 1 gal. of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities of solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gal. per mound. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness.</p> <p>NOTE: For best results, apply in cool weather 65-80°F, or in early morning or late evening hours. Treat new mounds as they appear.</p>
------------------	---

Insect Control on Ornamental Trees, Shrubs and Flowers

Trees and Shrubs	Insects	Rate per 3 Gals. of Water	Timing/Special Directions
<p>Arboreal, Azaleas, Birch, Boxwood, Camellias, Douglas Fir, Elm, Hawthorn, Holly, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Rose, Willow</p>	<p>Aphids, Bagworms, European Pine Shoot Moths, Holly Bud Moths, Dipsas Leafminers (Arboreal), Azalea, Birch, Boxwood, Holly, Oak, Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European Red, Privet, Two-spotted Spider), Oosure Root Weevils, Omnivorous Leafhoppers, Scale Crawlers (Cottony-cushion, Euonymus, Lecanium, Pine Needle, Pr. San Jose, Soft)</p>	<p>2 table- spoons</p>	<p>Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to penetrate dense foliage with spray</p>
	<p>Webworms (Juniper, Mimosa), Whiteflies</p>	<p>2 table- spoons</p>	<p>Apply as a thorough foliar spray. Mimosa Webworms apply as a thorough foliar spray when adults first appear. Make two additional applications at 4-5 week intervals</p>
	<p>Apple-and-Thorn Scale, Hemlock Chermes, Oak Looper, Opossum-eared Leafhopper, Pear Slug, Tent Caterpillars, Webworms (Colonizer, Fall)</p>	<p>2 table- spoons</p>	<p>Apply as a thorough foliar spray</p>

Roses and Flowers	Insects	Rate per 3 Gals. of Water	Timing/Special Directions
<p>Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilac, Rhododendron, Roses</p>	<p>Aphids, Euonymus Scale Crawlers, Flea Beetles, Dipsas Leafminers, Leafhoppers, Mealybugs, Mites (Carnation Bud, Carnation Shoot, Cyclamen, European Red, Two-spotted Spider), Thrips, Whiteflies</p>	<p>2 table- spoons</p>	<p>Apply as a thorough foliar spray. Try to spray underside of leaves and penetrate dense foliage.</p>

Fruits and Nuts

Start spraying after blooming. Applications should be made according to infestation or when aphids, mites, and crawler stages of scale insects are first noted.

Observe stated time intervals between last application and harvest of the crop, as well as intervals between applications and total number of applications that can be made per season.

Crop	PHI*
Almonds	**
Apricots	21
Blackberries	7
Boysenberries	7
Cherries	21
Dewberries	7
Grapes	28
Loganberries	7
Nectarines	21
Peaches	21
Plums	21
Prunes	21
Raspberries	7
Strawberries	15
Walnuts	45

*PHI = Pre-Harvest Interval, or the number of days between the last application and harvest

** Dormant spray only

Note: To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds around the trees are in bloom

Crop	Pest	Rate	Timing/Special Directions
<p>Almonds (CA only)</p>	<p>San Jose scale, Panatoria scale</p>	<p>2 table- spoons in 3 gals. of water</p>	<p>Apply only as a dormant spray. Do not apply more than 6 gallons per 500 sq. ft. of orchard.</p> <p>This product may be used on almonds only in CA. Do not use this product on almonds grown in other states.</p>

BEST AVAILABLE COPY 1

5 of 13

Apricots	Aphids. Apricot Mealybugs. Mees (Brown, Two-spotted Spider). Scale Crawlers (Olive, San Jose)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Cane- berries (CA, OR, WA only): Black- berries, Boysen- berries, Dew- berries, Logan- berries, Rasp- berries	Aphids. Leafhoppers. Raspberry Fruitworms. Two-spotted Spider Mees. Trips	2 tablespoons in 3 gals. of water	Apply as a full cover spray. Do not apply more than 5 gallons per 500 sq. ft. of field. Do not make more than 5 applications per season a minimum of 14 days apart. Do not apply within 7 days of harvest. This product may be used on caneberries only in CA, OR and WA. Do not use this product on caneber- ries grown in other states.
Cherries	Black Cherry Aphids. Cherry Fruitflies. Cherry Fruitworms (Maggots). Fruitree Leafrollers. Mees (Cherry Rust). Scale Crawlers (San Jose)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 7 days apart. Do not apply within 21 days of harvest.
Grapes	Aphids. Leafhoppers. Grape Berry Moths Grape Leafhoppers	4 1/2 teaspoons in 3 gals. of water	Apply as a full cover spray. Do not apply more than 3 gallons per 500 sq. ft. of vineyard. Do not make more than 5 applica- tions per season a minimum of 7 days apart. Do not spray within 28 days of harvest.
Nectarines	Aphids Mees (Brown, Two-spotted Spider) Scale Crawlers (Olive, San Jose)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Peaches	Aphids. Mees (Brown, Two-spotted Spider) Scale Crawlers (Olive, San Jose)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 14 days apart. Do not apply within 21 days of harvest.

Plums and Prunes	Aphids (Leafhopper Plum, Mealy Plum, Thistle). Mees (Brown, European Red)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 7 days apart. Do not apply within 21 days of harvest.
Straw- berries	Aphids. Two-spotted Spider Mees. Strawberry Leafrollers	2 tablespoons in 3 gals. of water	Apply as a full cover spray. Do not apply more than 3 gallons per 500 sq. ft. of field. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 5 days of harvest.
Walnuts (CA only)	Aphids. Scale Crawlers (Walnut, Italian Pear, Frosted). Spider Mees. Twig Borers	2 tablespoons in 3 gals. of water	Apply as a full cover spray. Do not apply more than 6 gallons per 500 sq. ft. of orchard. Do not make more than 3 applica- tions per season 14 days apart. Do not apply after fruits open or split. Do not apply within 45 days of harvest. This product may be used on walnuts only in CA. Do not use this product on walnuts grown in other states.

BEST AVAILABLE COPY

6813

Vegetables

For foliar or soil applications. Diazinon Lawn and Garden Insect Control should be applied in the volume of water recommended in the table below to assure thorough coverage of the foliage or soil.

Observe stated time intervals between last application and harvest as well as intervals between applications and maximum numbers of applications per season. Do not apply to seed crops grown in greenhouses.

Pre-Harvest Interval (PHI*) for Foliar Applications

Crop	PHI* (Days)
Beans ¹ (succulent varieties only: Lima, Pole and Snap)	7
Broccoli	7
Cabbage	21
Cantaloupes	3
Carrots	14
Cassias	2
Cauliflower	7
Chinese Broccoli, Chinese Cabbage, Collards	10
Crenshaws	3
Cucumbers	7
Envy (Escarole)	14
Green Melons	3
Kale	10
Lettuce	14
Muskmelons	3
Onions (Bulb and Green), Parsnips	14
Peas ¹	7
Peppers	3
Potatoes (Irish)	35
Radishes, Spinach	14
Squash, Summer	7
Squash, Winter	3
Sweet Corn	7
Tomatoes	7
Turkips (Roots and Tops)	14
Watermelons	3

* Pre-harvest interval, in the number of days between the last application and harvest.

¹ Do not use this product on dried pea or bean varieties such as dried beans, dried lima, split peas or dried black-eyed peas.

Vegetables

For application to the foliage, thoroughly spray all plant parts. Application to the soil should be mixed into the soil immediately after application.

Crop	Pest	Rate	Timing/Special Directions
Beans, Lima, Pole and Snap (succulent varieties only)	Aphids, Cucumber beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Do not apply within 7 days of harvest.
	Curworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface curworms, 3-5 inches for subterranean curworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
Beets, Red (Table)	Wireworms (snap beans only)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-6 inches.
	Curworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface curworms, 3-6 inches for subterranean curworms.
Broccoli	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or other type insecticides.
Cabbage	Curworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface curworms, 3-6 inches for subterranean curworms.
	Root Maggots	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-6 inches.
	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 21 days of harvest.

BEST AVAILABLE COPY

7 of 13

Carnes	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 14 days of harvest. NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for sur- face cutworms, 3-6 inches for subterra- nean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before plant- ing. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches.
Cauliflower	Aphids, Diamondback Moths Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 7 days of harvest. NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
Chinese Broccoli	Aphids, Diamondback Moths Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 10 days of harvest. NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
Chinese Cabbage (Box Choy and Napa)	Aphids, Diamondback Moths Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 10 days of harvest. NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
Cucumbers	Aphids, Diamondback Moths Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 10 days of harvest. NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.

Com (see sweet corn)			
Cucumbers	Aphids, Thrips, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 7 days of harvest.
Encive (Escarole)	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 14 days of harvest. NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
Kale	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 10 days of harvest. NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
Lettuce	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 14 days of harvest. NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for sur- face cutworms, 3-6 inches for subterra- nean cutworms.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches.
Metons (Carni- touses Cabbages Cen- shews Honey- cres Must- ardens Parsnips and Hydras of these) and Water- melons	Aphids, Thrips, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 3 days of harvest.

BEST AVAILABLE COPY

8 7 13

Onions (Bulb and Green)	Onion Thrips	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.
Parsnips	Aphids. Flea Beetles. Leafhoppers. Dipterous Leafminers. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat applications every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Peas (succulent only)	Aphids. Cucumber Beetles. Grasshoppers. Leafhoppers. Mexican Bean Beetles Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Allow at least 7 days between the last application and harvest
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting immediately work into soil 4-8 inches
NOTES: (1) Do not apply this product to dried pea varieties such as split peas or dried black-eyed peas. (2) To protect bees, do not apply foliage treatments if bees or bees in the treatment area are in bloom.			
Peppers	Aphids. Colorado Potato Beetles. Flea Beetles. Leafhoppers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 5 days of harvest
Potatoes	Aphids. Flea Beetles. Leafhoppers. Colorado Potato Beetles. Southern Armyworms. Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 35 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.

Radishes	Aphids. Flea Beetles. Leafhoppers. Imported Cabbageworms. Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 3 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
Squash	Aphids. Flea Beetles. Leafhoppers. Dipterous Leafminers. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Squash, Summer	Aphids. Thrips. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.
Squash, Winter	Aphids. Thrips. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest.
Sweet Corn	Aphids, Thrips. Flea Beetles. Grasshoppers Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days following last application.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting immediately work into soil 4-8 inches
NOTE: To protect bees, do not apply to corn foliage during the pollen shed period.			

BEST AVAILABLE COPY

9813

Tomatoes	Aphids, Colorado Potato Beetles, Fall Armyworms, Flea Beetles, Leafhoppers, Southern Armyworms, Vinegar Flies (Drosophila spp.)	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 1 day of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for sur- face cutworms, 3-6 inches for subterra- nean cutworms.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting Immediately work into soil 4-8 inches.
Beans and Turnip Tops	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 14 days of harvest. NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for sur- face cutworms, 3-6 inches for subterra- nean cutworms

Storage and Disposal

Storage

Do not store near heat or open flame. Store in original container in an area inaccessible to children and pets. **USE CONTENTS WITHIN 6 MONTHS AFTER OPENING.**

Disposal

Do not reuse empty container. Thoroughly rinse container, securely wrap it in several layers of newspaper and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye injury. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils such as spoons and measuring cups must not be used for food purposes after use in measuring pesticides.

Statement of Practical Treatment

If swallowed: Call a physician immediately. **DO NOT INDUCE VOMITING UNLESS NO OTHER TREATMENT IS AVAILABLE.** Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If inhaled: Remove victim to fresh air and apply artificial respiration if indicated. Get medical attention.

If on skin: Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

If in eye: Rinse eyes with plenty of water and call a physician immediately.

Note to Physician: This product is an organophosphorus insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains aromatic petroleum distillates which may present an aspiration hazard. Gastric lavage may be indicated if product was taken orally.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migration habits of certain Atlantic Coast waterfowl, do not apply this product to lands in Nassau County, New York, between November 1 and May 20. Do not exceed the minimum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lands where irrigation (watering) is recommended in the directions or if waterfowl, i.e., duck or geese, can be expected in the treated area after treatment (except spot treatment), water down with at least 1/4 inch of water immediately after spraying this product, however, stop watering before packing occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.

BEST AVAILABLE COPY

10 of 13

Sunniland®

25% DIAZINON LIQUID CONCENTRATE

**THIS PRODUCT MUST NOT BE USED
ON GOLF COURSES AND SOD FARMS.**

ACCEPTED
with COMMENTS
in EPA Letter Dated

**FOR OUTDOOR USE ON THE HOME LAWN
AND IN THE HOME GARDEN ONLY.**

FEB 16 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

9404-65

**A SPRAY CONCENTRATE
CONTAINS DIAZINON INSECTICIDE**

**Protects - Lawns, Vegetables, Fruits & Nuts, Roses, Flowers,
Trees, Shrubs**

**Controls - Ants, Aphids, Chinch Bugs, Cutworms, **Fire Ants,
Flea Beetles, Grubs, Mites, Sod Webworms and
many other insects. **aids in control.**

See full label in booklet attached to this container
for complete directions for mixing and applying
this product and a complete list of pests controlled.

ACTIVE INGREDIENT:

**DIAZINON: [O,O-Diethyl O-(2-isopropyl-6-methyl-
4-pyrimidinyl) phosphorothioate] -**

25.00%

INERT INGREDIENTS* - - - - -

75.00%

TOTAL 100.00%

***Contains Aromatic Petroleum Distillates**

KEEP OUT OF REACH OF CHILDREN

CAUTION

See end of booklet for additional precautionary statements

EPA REG. NO. 9404-65

EPA EST. NO. 9404-FL-1

NET CONTENTS 1 PINT;(1 QUART; 1 GALLON)

SUNNILAND CORPORATION

SANFORD, FL 32771

BEST AVAILABLE COPY

11 of 13

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

General Information

This Product is an emulsifiable concentrate product which must be diluted with water before application. This product kills many outdoor pests which damage lawns, roses, flowers, trees, shrubs, vegetables and fruits and insects which annoy pets and people.

SEE THE BOOKLET UNDER CAP FOR COMPLETE DIRECTIONS FOR USE AND PESTS CONTROLLED.

Notes: (1) USE CONTENTS WITHIN 6 MONTHS AFTER OPENING. SEE STORAGE AND DISPOSAL DIRECTIONS. (2) Do not use in the home. (3) Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia or orchids since plant injury may occur. (4) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties. (5) Do not apply to food crops grown in greenhouses.

After spraying, wait the following number of days before harvesting fruits, nuts, or vegetables:

1 day	Tomatoes
3 days	Melons, Winter Squash
5 days	Peppers, Strawberries
7 days	Beans, Broccoli, Blackberries, Boysenberries, Cauliflower, Cucumbers, Dewberries, Loganberries, Peas, Raspberries, Summer Squash, Sweet Corn
10 days	Chinese Broccoli, Chinese Cabbage, Collards, Kale
14 days	Carrots, Endive (Escarole), Lettuce, Onions (Bulb and Green), Parsnips, Radishes, Spinach, Turnips
21 days	Apricots, Cabbage, Cherries, Nectarines, Peaches, Plum, Prunes
28 days	Grapes
35 days	Potatoes (Irish)
45 days	Walnuts
*	Almonds (one dormant, spray only)

Precautions: (1) Wear a long-sleeved shirt and long-legged pants when spraying. (2) Spray with the wind to your back. (3) Do not spray on windy days. (4) If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water and launder clothing before wearing again.

* This table is only a partial listing of the pests controlled by this product. See booklet under cap for complete directions for use and pests controlled.

Problem Area	Pests*
Nuisance Pests in Outside Areas	Ants, Brown Dog Ticks, Chiggers, Cockroaches, Crickets, Fleas, Fire Ants, Spiders and others*
Lawns	Chinch Bugs, Ants, Sod Webworms, Brown Dog Ticks, Fleas, White Grubs, Lawn Billbugs and others*
Vegetables: Foliar Application	
Beans	Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles
Peas (except dried varieties)	Spider Mites
Broccoli, Chinese Broccoli, Cabbage, Chinese Cabbage, Cauliflower, Collards, Kale	Aphids, Flea Beetles, Dipterous Leafminers, Diamondback Moths, Imported Cabbageworms
Endive (Escarole), Lettuce, Spinach	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites

BEST AVAILABLE COPY

12 8 13

Peppers. Tomatoes	Aphids. Colorado Potato Beetles. Fall Armyworms (Tomatoes). Flea Beetles. Leafhoppers. Southern Armyworms (Tomatoes). Vinegar Flies - <i>Drosophila</i> spp. (Tomatoes)
Onions (Bulb and Green)	Onion Thrips
Radishes. Turnips. Carrots. Parsnips. Potatoes	Aphids. Flea Beetles. Leafhoppers. Dipterous Leafminers. Colorado Potato Beetles (Potatoes). Southern Armyworms (Potatoes)
Sweet Corn	Aphids. Flea Beetles. Grasshoppers. Spider Mites
Melons. Squash (Summer, Winter)	Aphids. Thrips. Spider Mites
Vegetables: Preplant Soil Application	
Beans. Peas (except dried varieties)	Cutworms. Mole Crickets. Wireworms (Snap Beans)
Beets. Red (Table)	Cutworms. Mole Crickets
Cabbage	Cutworms. Root Maggots. Wireworm
Carrots. Sweet Corn	Cutworms. Mole Crickets. Wireworms
Lettuce. Tomatoes	Cutworms. Wireworms
Radishes. Turnips	Cutworms
Fruits & Nuts	
Almonds (California only)	Scale Crawlers
Walnuts (California only)	Aphids. Scale Crawlers. Spider Mites and others*
Apricots. Peaches. Nectarines	Aphids. Apricot Mealybugs (Apricots). Spider Mites. Scale Crawlers
Cherries	Black Cherry Aphids. Cherry Fruitflies. Cherry Rust Mites. Scale Crawlers and others*
Plums. Prunes	Aphids. Spider Mites
Canberries (CA. OR. WA only) Blackberries. Boysenberries. Dewberries. Loganberries. Raspberries	Aphids. Leafhoppers. Two-Spotted Spider Mites. Thrips. Raspberry Fruitworms
Strawberries	Aphids. Two-Spotted Spider Mites. Strawberry Leafrollers
Grapes	Aphids. Grape Berry Moths. Grape Leafrollers. Leafhoppers
Roses & Flowers	
Azaleas. Camellias. Chrysanthemums. Carnations. Gladiol. Lilac. Rhododendron. Roses	Aphids. Flea Beetles. Dipterous Leafminers. Leafhoppers. Mealybugs. Two-spotted Spider Mites. Thrips. Whiteflies and others*

BEST AVAILABLE COPY

13 813

Ornamental trees & Shrubs

Arborvitae, Aphids, Mites, Scale Crawlers, Whiteflies, Tent Caterpillars and others*
 Azaleas, Birch,
 Boxwood, Camellias,
 Douglas Fir, Elm,
 Hawthorn Holly,
 Juniper, Lilac,
 Locust, Maple,
 Mimosa, Oak, Pine,
 Ornamental Plum,
 Poplar,
 Rhododendron,
 Spruce, Willow

Storage and Disposal**Storage**

Do not store near heat or open flame. Store in original container in an area inaccessible to children or pets. **USE CONTENTS WITHIN 6 MONTHS AFTER OPENING.**

Disposal

Do not reuse empty container. Thoroughly rinse container, securely wrap in several layers of newspaper, and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

Precautionary Statements**Hazards to Humans and Domestic Animals****CAUTION**

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye injury. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils such as spoons and measuring cups must not be used for food purposes after use in measuring pesticides.

Statement of Practical Treatment

If swallowed: Call a physician immediately. **DO NOT INDUCE VOMITING UNLESS NO OTHER TREATMENT IS AVAILABLE.** Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If inhaled: Remove victim to fresh air and apply artificial respiration if indicated. Get medical attention.

If on skin: Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

If in eyes: Rinse eyes with plenty of water and call a physician immediately.

Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains aromatic petroleum

distillates which may present an aspiration hazard. Gastric lavage may be indicated if product was taken internally.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlap, ping of sprays. For lawns, where irrigation (watering) is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply at least 1/4 inch of water immediately after spraying this product, however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.

BEST AVAILABLE COPY