

UNDER THE
FUNGICIDE ACT
FOR ECONOMIC
ED UNDER NO. 9431-2

comment

NET CONTENTS

Hanco

DAIRY-TERG

DAIRY DETERGENT-SANITIZER

DIRECTIONS FOR USE

To prepare a standard cleaner-sanitizer solution for general dairy farm use dissolve 1 ounce (one measure) in 2 gallons of warm water.

Before Morning Milking:

1. Flush each milker unit with clean safe hot water and rinse other utensils.*

After Morning Milking:

1. Immediately draw at least one pailful of clean warm water through each unit.
2. Dismantle machine, including air hose, and immerse with other utensils in cleaner-sanitizer solution (1/2 ounce per gallon of water). Soak one-half hour if possible.
3. Brush all parts inside and out. Assemble milker.
4. Allow to dry without rinsing. Protect all equipment from contamination.

*NOTE: The hot water used for rinsing the utensils before milking can then be used for cleaning the udders by adding 1/2 ounce of cleaner-sanitizer per gallon of water. "Safe" means water of drinking quality.



ACTIVE INGREDIENTS:

Sodium carbonate monohydrate and n-alkyl (C14, C12, C16) dimethyl benzyl ammonium chlorides 40%

INERT INGREDIENTS:

Sodium tripolyphosphate and octyl phenoxy polyethoxy ethanol, water 60%

CAUTION
KEEP OUT OF REACH OF CHILDREN.
SEE CAUTION ON SIDE PANEL.

U.S.D.A. REG. NO. 1074-24

HANCO MANUFACTURING CO.
MEMPHIS, TENNESSEE

Before Night Milking:

1. Flush each milker unit with clean safe hot water and rinse other utensils.*

After Night Milking:

1. Rinse milker and other utensils with warm water.
2. Flush milker with cleaner-sanitizer solution. (1/2 ounce per gallon of water).
3. Wash other utensils.
4. Allow to dry without rinsing. Protect all equipment from contamination.

CAUTION

Avoid contamination of food. Harmful if swallowed. Keep out of eyes. In case of contact, flush with plenty of water. Dilute as directed. Do not mix with soap or anionic detergents. Avoid getting the concentrate on skin. If irritation persists, get medical attention.

Use DAIRI-TERG Detergent-Sanitizer for cleaning and sanitizing milking machines and utensils, and for washing cow's udders. This product reduces cleaning time to a minimum by combining cleaning and sanitizing in one operation. The daily use of DAIRI-TERG will aid in the production of cleaner milk with low bacteria counts.

ACCEPTED WITH COMMENTS

NET CONTENTS

ONS

HANCO 950 is recommended for use in hospitals for cleaning, disinfecting, and deodorizing. Use on floors, walls, ceilings, equipment and fixtures in areas such as operating rooms, clinics, isolation wards, nurseries, diet kitchens, rest rooms, etc.

Use HANCO 950 for cleaning and disinfecting and deodorizing in schools and public institutions all hard surface areas such as corridors, class rooms, offices, rest rooms, locker rooms, showers. Use in public places where cleanliness and bacterial control are vital for the protection of health.

Use HANCO 950 for cleaning and sanitizing and deodorizing food handling equipment in food, milk, and beverage processing plants, restaurants, farms. Such equipment would include blanchers, evaporators, homogenizers, holding tanks, pipes, valves, fittings, utensils, etc. All equipment should be flushed with clear water just prior to use.

HANCO 950 is non-corrosive in use dilutions.

PHENOL COEF A.O.A.C Method

s. aureus.....27
s. typhosa.....19

HANCO 950 CONCENTRATED DETERGENT DISINFECTANT - SANITIZER DEODORANT

ACTIVE INGREDIENTS:

n-Alkyl (C₁₂ 50% C₁₄ 30%
C₁₆ 17% C₁₈ 3%) dimethyl
ethylbenzyl ammonium
chloride.....1.6%
n-Alkyl (C₁₄ 60% C₁₆ 30%
C₁₂ 5% C₁₈ 5%) dimethyl
benzyl ammonium chloride..1.6%
INERT INGREDIENTS.....96.8%

CAUTION
KEEP OUT OF REACH OF CHILDREN
See Caution Statement Side Panel

USDA REG NO 9931-x

HANCO MANUFACTURING CO
Memphis Tennessee

ACCEPTED
OCT 16 1968
UNITED STATES DEPARTMENT OF AGRICULTURE
FOR THE

DIRECTIONS

For cleaning and disinfecting floors, walls, surfaces as found in hospitals, institutions, public rooms, homes, industrial areas, conveyances, etc.

Dilute 2 ounces of HANCO 950 to each gallon of water and apply with a mop or cloth.

In instances where surfaces are heavily contaminated with soil and/or organic matter, (i.e. operating rooms), apply solution twice to ensure complete protection. For maximum germicidal and detergent activity, surfaces to be treated should be wet thoroughly.

For cleaning and Sanitizing apply HANCO 950 at least one ounce to one gallon of water. Allow at least two minutes of contact with surface before rinsing.

For Cleansing and Deodorizing apply one ounce diluted to one or two gallons of water.

Caution

HANCO 950 is a concentrated detergent-disinfectant and must be diluted before using. Solutions should not come in contact with food. Solutions should not be used with soap or anionic detergents.

Avoid getting concentrate in eyes. In case of contact, wash thoroughly with water. If irritation persists get medical attention.