JAN 1 0 1992

Ornbelt Chemical Company P.O. Box 410 GCOOK, NE 69001



hear Mr. Kohmetscher:

Subject: Revised labeling

Sevin 5% Pait

PPA Registration No. 10107-133

Your submission of October 15, 1991

The resident referred to above, submitted is connection with sugjectable provided that you:

1. Yok the labeling changes listed below before you relate the product for shipment bearing the smeaded lobeling:

Change the PMI for the following crops:

Crops	Rate	BHI	
Alfalfa, peas Cotton	30 lbs. 30 lbs.	7-dayo 7-days	
Apples, cherries, peaches, pears, plums Mectarines Citrus Apricots	40-48 lbs. 40-48 lbs. 40-48 lbs. 40-43 lbs.	1-61ys 3-62ys 5-63ys 28-63ys	

Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the Λ -79 Enclosure for further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with STERA sec. 6(e). Your release for chipment of the project bearing the amended labeling constitutes acceptance of these conditions.

			CONCURRENC	ES			
SYMBOL							
DIRMANE !	4	1	 1	J	į.	ł	1
DATE		,				<u>l</u>	<u>}</u>
EPA Farm	1320-1 (12-70)	<u></u>	 			OFFICE	AL FILE COPY

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

DHE

Dennis H. Edwards, Jr. Product Manager (12) Insecticide Rodenticide Branch Registration Division (H-7505C)

Enclosure

5% Sevin® Bait Under the Federal Insecticide. Act

generided, for the pesticide red under EPA Res. No.

AN ATTRACTANT INSECTICIDE __IOIO7-143

For the Control of Ants, Cutworms, Armyworms, Crickets, Darkling Beetle, Sowbugs and Grasshoppers

ACTIVE INGREDIENT:	BY WEIGHT
Carbaryl (1-naphthyl N-methylcarbamate)	5.00% by wt.
INERT INGREDIENTS	95.00% by wt.
	TOTAL100.0% by wt.

For MEDICAL and TRANSPORTATION Emergencies ONLY, Call Chemtrec 1-800-424-9300 **@SEVIN** is the registered trademark of Rhone Poulenc Ag Company for Carbaryl Insecticide.

KEEP OUT OF THE REACH OF CHILDREN AND ANIMALS CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Never give anything by mouth to an unconscious or convulsing person. If conscious and not convulsing, drink 1 to 2 glasses of water and induce vomiting by touching the back of the throat with finger.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash thoroughly with soap and water.

IF INHALED: Move from contaminated atmosphere and call a physician.

NOTE FOR PHYSIC'AN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

RE-ENTRY STATEMENT

Do not enter treated areas without protective clothing until dusts have settled. Be cause certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral precautions must be given to workers engaged in agricultural hand labor who are expected to be in a treated area or in an area bout to be treated with this product. These precautions are: CAUTION: Area treated with carbaryl on (date of application). Do not enter treated area the day of application without protective clothing and bloves. Protective clothing at a minimum should include clean long pants, long sleeved shirt, shoes and socks. For accidental exposure, including skin contact with pellets, see Statements of Practical Treatment.

Oral precautions must be given if there is reason to believe that written precautions cannot be understood by workers. When oral precautions are given, precautions shall be given in a language customarily understood by workers.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS. CAUTION

MAY BE HARMFUL IF SWALLOWED. Avoid breathing vapors (dust or spray-mist). Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water and wetlands, except under the forest canopy and use on rice. Discharge from rice fields may kill aquatic and estuarine invertebrates. Do not contaminate water by cleaning equipment of disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

EPA Reg. No. 10107-143

EPA Est. No. 10107-NB-1

Net Weight ____ **Pounds**

Manufactured for:

Combelt Chemical Company. • P.O. Box 410 • McCook, Nebraska 69001 **REV 10-01**

STORAGE AND DISPOSAL

PROHIBITION: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Dispose of bags according to approved Federal, State or local procedures under the Resource Conservation and Recovery Act.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION

Tolerances established under the Federal food, Drug and Cosmetic Act permit the sale of crops bearing probable SEVIN residues when SEVIN is used in accordance with label directions.

If SEVIN insecticide is used in accordance with label directions, these crops, including bean vines, carrot tops, pea vines, sugar beet tops, corn forage and cotton gorage may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in milk and meat.

CROP	RATE PER ACRE	RATE PER 1 1,000 SQ. FT.	DAYS BETWEEN BIAL APPLICATION AND HARVEST	CROP	RATE PER ACRE	RATE PER 1 1,000 SQ. FT.	DAYS BETWEEN FINAL APPLICATION AND HARVEST
Cucumbers, melons, pesnuts, pumkins, squesh	20 lbs.	7½ ozs.	0	Wheat (grain — Fall or Spring)	30 lbs.	11 28.	21
Birdsfoot trefoil (forage and hay), clover, wheat (pasture, forage)	30 lbs.	11 ozs.	0	Sorghums (mile, grain sorghum, sweet sorghum and hybrids	20-40 lbs.	71/1 -14 ² /1 ozs	. 21(grain)
Lewn and turf — GROUND APPLICATION ONLY	30 lbs.	11 ozs.	0	Oats, barley and nye (fall treatment)	20-40 lbs.	<u> </u>	<u> </u>
Pasture (rangeland and waste- lens), forage grasses, fence row and non-crop areas				Do not feed treated fodder to delay animals or animals being finis slaughter or allow them to graze on crops treated with this pro Sunflowers 20-40 lbs. 71/s -147/s ozs.			
Beans, carrots, com (field, sweet, pop), eggplant, okra, peppers, polatoes, aorghum forage, soybeans, sweet potatoes, tobacco beds, fornatoes	20-40 lbs.	7 ¹ /s -14 ² /s 025.	0		(crop for oil, food or feed purpose 0 (crop for seeding purposes on crickets, house crickets, mole crickets, rs, sowbugs and ants)		
Asneragus, strewberries	40 lbs.	14 ² /s 02\$.	1	Annual plants—herbaceous plants	T		
Broccoll, Brussels sprouts, cabbags, cauliflower, cowpeas, garden beets (routs), head lettuce, horseradish, kohlrabi, parsnipa, radishes, rutabagas, salsify (roots), tobacco (field), turnips (roots)	20-40 lbs.	7 /2 -14 ⁹ /2 ozs.	3	and shrubs, trees, woody plants (ornamental), grapes Apples, cherries, peaches, pears, plums Nectarines Citrus	40-48 lbs. 40-48 lbs. 40-48 lbs. 40-48 lbs.	7½ -14½ 028 7½ -14½ 028 7½ -14½ 028 7½ -14½ 029	. 0
Affaife, pees	JO lbs.	11 ozs.	3				1
Cotton	30 lbs.	11 029.	3	Apricots		71/2 -147/2 OZS	<u> </u>
Celary, Chinese cabbage, collards, dendefion, endive (secarole), garden beet tops, Henover saled, Itale, leaf lettuce, mustard greens, persley, rice, salely (tops), spinech, sugar beets, Swise chard, tumip tops. "Do not use on rice fields in which crayfish and/or carlish farming is included in the cultural practice.	20-40 lbs.	71/2 -14 ⁸ /3 0 ~3	14	NOTE: Do notmake more than two as and clovers emerge from boot. Do no crops.	pecations at	ier grain neads	ou wheat, all an

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directione, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURERS OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTACT, NEGLIGENCE, STRICT LIABILITY IN TORY OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.