

10140-2

SEBESTA'S PRAIRIE DOG, GROUND SQUIRREL, FIELD MICE, AND KANGAROO RAT BAIT

Active Ingredient		
Strychnine	0.44%	
Inert Ingredients	99.56%	
Total	100.00%	

Net Weight 2½ Lbs.

ALSO AVAILABLE IN 5 LB. & 10 LB. BAGS

DANGER - CONVULSIVE POISON: KEEP AWAY
from children and domestic animals. Avoid contamination of foodstuffs. Poisonous if
swallowed. Wash hands after using.

Directions For Use

- For use primarily on grass lands and harvested fields.
- For Prairie Dogs: Scatter tablespoonful quantities of bait near active mounds preferably on the side away from the pile of dirt which has been removed from the burrow.
- For Richardson and Franklin Ground Squirrels: Scatter teaspoonful quantities of bait on surface of ground near burrow entrance.
- For Thirteen-Lined Ground Squirrels: Scatter teaspoonful quantities of bait at the grass edge along the trails as close to the burrow opening as possible.
- For Kangaroo Rats: Scatter teaspoonful quantities of bait near kangaroo rat mounds. Use several bait placements for large mounds.
- For Directions for Use of this bait for Field Mice, Pine Mouse, Meadow mouse and White-footed Mouse see back panel.



POISON

ANTIDOTE: CALL A PHYSICIAN IMMEDIATELY.
If less than an hour has passed since the poison was taken, give a tablespoon of salt in a glass of water to remove unabsorbed strychnine. Have victim lie down in a quiet, dark room and keep him warm.

WARRANTY

The manufacturer being beyond our knowledge and control, we make no warranty expressed or implied, as to the effects of such use, when not used in accordance with directions.

**MANUFACTURED & DISTRIBUTED BY
SEBESTA BAIT MIXING PLANT
1220 E. HAVENS - LOT #23
MITCHELL, SOUTH DAKOTA 57301**

DESTROY CONTAINERS BY BURNING

40-00

40-00

DIRECTIONS FOR USE

For Field Mice, Pine Mice, Meadow Mice and White-Footed Mice.

1. SPOT OR TRAIL BAITING

- Place one teaspoon of bait in surface trails or at the mouth of holes leading to underground burrow systems.
- Two to four spots should be made near the base of each infested tree.
- Cover bait by pulling overhanging grass back into place, do not disturb the runway-system.
- Baiting should be conducted at the rate of 2 to 3 pounds per acre.

2. BROADCAST BAITING

- May be applied by cyclone sprayer or by hand. When applied by hand, throw 1/2 teaspoon amounts into heavy cover along tree trunks, creek edges and fence lines.
- Apply at the rate of 6 to 10 pounds per acre.
- Do not broadcast over growing crops.

3. MACHINE BAITING

- Use a modified corn planter with plates set to drop one kernel of bait along a line just inside the drip line on both sides of the tree.
- Bait should be applied at the rate of 5 to 6 pounds per acre.

4. SPECIAL CAUTIONS

- Store poisoned grains in locations not accessible to children, poultry and domestic animals.

**CAUTION: BIRDS FEEDING ON TREATED AREAS
MAY BE KILLED.**