

PM/16 10182-41/

181

Talon[®]-G

RODENTICIDE MINI-PELLETS

For Effective Control of Commensal Mice and Rats

ACCEPTED
SEP 01 1988
Under the Federal Insecticide, Fungicide and Rodenticide Act, as amended
Pesticide registered under EPA Reg. No. 10182-41/

EPA Reg. No. 10182-41
EPA Est. No. 2393 W-1
Net Contents 4.5 lb (20.4 kg)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

MAY BE HARMFUL OR FATAL IF SWALLOWED. KEEP AWAY FROM HUMANS, DOMESTIC ANIMALS AND PETS. WASH HANDS AFTER HANDLING BAIT.

IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE. For 24-hour emergency assistance call ICI Americas Inc. 1-800-327-8633.

IF BAIT IS EATEN BY ANIMALS OR PETS, CALL ICI Americas Inc. 1-800-327-8633 for 24-hour emergency assistance.

In case of significant spill call CHEMTREC 800/424-9300.

NOTE TO PHYSICIAN OR VETERINARIAN

This product may reduce the clotting ability of the blood and cause hemorrhaging. If poisoning occurs intramuscular and oral administration of vitamin K₁ are indicated, as in poisoning from overdose of bishydroxy coumarin. For human cases, vitamin K₁ is antidotal at doses of 10-20 mg (mg/kg). For animal cases, vitamin K₁ is antidotal at 2-5 mg/kg. Repeated doses may need to be given up to two weeks (based on monitoring of prothrombin time). In severe cases, blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard in birds of prey and mammals. Keep out of any body of water.

NOTICE TO BUYER AND USER: ICI Americas Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller and Buyer, and User assumes the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

Kills warfarin-resistant Norway rats and house mice. Effects may commence a lethal dose in one feeding with first dead rodents appearing later in the days after treatment begins.

ACTIVE INGREDIENT

Brodifacoum
3-[3-(4-bromo-1,1-biphenyl)-4-yl]-1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-benzopyran-2-one

INERT INGREDIENTS

TOTAL	100.000%
	0.000%
	0.000%
	100.000%

KEEP OUT OF REACH OF CHILDREN CAUTION

May Be Harmful or Fatal if Swallowed
Read Additional Precautionary Statements on Side Panel

This product is protected by U.S. patent numbers 3957824 and 4035505 which are the property of Ward Blenkinsop and Company Limited.



ICI Agricultural Products
Wilmington, Delaware 19897

A Business Unit of ICI Americas Inc.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container except for holding additional TALON rodenticide.

STORAGE: Store in original container only in a dry place inaccessible to children and pets.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Dispose of bait container in a sanitary landfill or by incineration if allowed by state and local authorities.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: For control of Norway rats, roof rats and house mice in and around homes, industrial, commercial, agricultural and public buildings. TALON may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. Do not use in sewers. This product must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

Authorized by the USDA for use in Federally inspected meat, poultry and egg processing plants.

SELECTION OF TREATMENT AREAS: Determine areas where rats and/or mice will most likely find and consume the bait. Generally these areas are along walls, by gnawed openings in or beside drywalls, in corners and concealed places between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

APPLICATION DIRECTIONS

Norway and Roof Rats

Apply 4 to 16 ounces of bait (usually at intervals of 15 to 30 feet) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease.

House Mice

Apply 1/2 to 1 ounce of bait at intervals of 8 to 12 feet per placement. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.

Rats & Mice

Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait according to disposal paragraph. To prevent reinfestation eliminate food, water and harborage as much as possible. If reinfestation does occur repeat treatment. Where a continuous source of infestation is present establish permanent bait stations and replenish bait as needed.

05-0780-265R2 AH