

PM 16  
10/82 -75

1901

**ACCEPTED**  
AUG 30 1983  
Under the Federal Insecticide, Fungicide, and  
Rodenticide Act, as amended, for this  
pesticide registered under  
EPA Reg. No. 10162-75

# HAVOC™

**RODENTICIDE  
MINI-PELLETS**

**KILLS RATS  
& MICE**



Kills warfarin-resistant  
Norway Rats and House Mice.

Rodents may consume a lethal dose in one feeding  
with first dead rodents appearing four or five days after  
treatment begins.

Active Ingredient:

Brodifacoum-	
3-[3-(4'-bromo-[1,1'-biphenyl]-4-yl)-	
1,2,3,4-tetrahydro-1-naphthalenyl]-4-	
hydroxy-2H-1-benzopyran-2-one	0.005%
Inert ingredients	99.995%
Total	100.000%

**KEEP OUT OF REACH  
OF CHILDREN**

**CAUTION**

May Be Harmful or Fatal if Swallowed.  
Read Additional Precautionary Statements  
on Side Panel.

ICI Americas Inc.  
Wilmington Delaware 19897

EPA Reg. No. 10162-75  
EPA Est. No. 523-IL-1  
Net Contents: 45 lb (20.4 kg)

**PRECAUTIONARY STATEMENTS  
DANGERS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

**IT MAY BE HARMFUL OR FATAL IF SWALLOWED. KEEP AWAY  
FROM HUMANS, DOMESTIC ANIMALS, AND PETS. WASH  
HANDS AFTER HANDLING BAIT.**

**IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE.  
24-hour emergency assistance call ICI Americas  
Medical Emergency Information-1-800-327-8633  
IF BAIT IS EATEN BY ANIMALS OR PETS,  
24-hour emergency assistance call ICI Americas  
Medical Emergency Information-1-800-327-8633**

**In case of significant spill, call CHEMTREC 800/424-9300.**

**WARNING TO PHYSICIAN OR VETERINARIAN**  
This product may reduce the clotting ability of the blood and  
cause hemorrhaging. If this occurs, intramuscular and oral  
administration of vitamin K, combined with blood transfusions  
indicated as in the case of hemorrhage caused by overdoses  
of hydroxycoumarin. Vitamin K<sub>1</sub> is antidotal at doses of 10 to  
mg (not mg/kg) and should be given and repeated as  
necessary (based on monitoring of prothrombin times).

**ENVIRONMENTAL HAZARDS**  
This product is toxic to fish, birds and wildlife. This product can  
be a secondary hazard to birds of prey and mammals. Keep  
away from any body of water.

**WARNING TO BUYER AND USER:** Seller warrants that this  
product conforms to the chemical description on the label and is  
suitably fit for the purposes stated on the label when used in  
accordance with directions under normal conditions of use. This  
warranty does not extend to the use of this product contrary to  
label instructions, or under abnormal use conditions, or under  
conditions not reasonably foreseeable to Seller and Buyer and  
Buyer assumes the risk of any such use. SELLER DISCLAIMS  
ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING  
ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUEN-  
TIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM  
USE OR HANDLING OF THIS PRODUCT AND SELLER'S  
LIABILITY AND BUYER'S AND USER'S EXCLUSIVE  
REMEDY SHALL BE LIMITED TO THE REFUND OF THE  
PURCHASE PRICE.

## STORAGE AND DISPOSAL

**STORAGE:** Do not contaminate water, food or feed by storage or  
disposal. Open dumping is prohibited. Do not reuse empty  
containers.

**DISPOSAL:** Bait that cannot be used according to label instruc-  
tions must be disposed of according to applicable Federal,  
State, or local procedures. Dispose of bait container in a sanitary  
landfill or by incineration if allowed by State and local authorities.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner  
inconsistent with its labeling.

**USE RESTRICTIONS:** For control of Norway Rats, Roof rats and  
House mice in and around homes, industrial, commercial, agr-  
cultural and public buildings. HAVOC may also be used in and  
around transport vehicles (ships, trains, aircraft) and related port  
or terminal buildings. Do not use in sewers. This product must be  
placed in temper-proof bait boxes or in locations not accessible  
to children, pets, domestic animals or wildlife. Do not place bait  
in areas where there is a possibility of contaminating food or  
surfaces that come in direct contact with food.

**SELECTION OF TREATMENT AREAS:** Determine areas where  
rats and/or mice will most likely find and consume the bait.  
Generally, these areas are along walls, by gnawed openings, in  
or beside burrows, in corners and concealed places, between  
floors and walls, or in locations where rodents or their signs have  
been observed. Remove as much food as possible.

## APPLICATION DIRECTIONS

**Norway and Roof Rats:**  
Apply 4 to 16 ounces of bait (usually at intervals of 15 to 30  
feet) per placement. Maintain an uninterrupted supply of fresh  
bait for 10 days or until signs of rat activity cease.

**House Mice:**  
Apply 1/2 to 1 ounce of bait at intervals of 8 to 12 feet per  
placement. Larger placements (up to 2 ounces) may be needed  
at points of very high mouse activity. Maintain an uninterrupted  
supply of fresh bait for 15 days or until signs of mouse activity  
cease.

**Rats & Mice:**  
Replace contaminated or spoiled bait immediately. Collect  
and dispose of all dead animals and unconsumed bait according  
to "Disposal" paragraph. To prevent reinfestation, eliminate  
food, water, and harborage as much as possible. If reinfestation  
does occur, repeat treatment. Where a continuous source of  
infestation is present, establish permanent bait stations and  
replenish bait as needed.

1901