

100 GALS. AER. CLO. 5

P1059

SUPPLEMENT - 1968

USE DESIGNATED AMOUNT IN 100 GALS. WATER AND SPRAY AS SPECIFIED.  
APPLY THOROUGH, FULL COVERAGE SPRAY. REPEAT AS NECESSARY UNLESS OTHERWISE SPECIFIED.

WHEN TO APPLY: Do not apply to Apples, Cherries, Grapes, Prunes, Plums, Broccoli, Turnips, Cranberries; Eggplant, Peppers, Strawberries and Green Onions within 3 days of harvest; Peaches, Brussels Sprouts, Cauliflower, Cabbage, Kale, Mustard Greens, Celery, Beets, Citrus and Spinach within 7 days; Beans, Cucumbers, Squash, Melons, Pears and Tomatoes within 1 day, Head Lettuce within 7 days; Leaf Lettuce within 14 days of harvest. Use up to harvest on Pecans, Walnuts, Potatoes and Cotton.

CRANBERRIES: Leafhoppers, Spittlebug Nymphs, Black-headed fireworms, Cranberry Fruitworm - 1 1/2 pts.

CUCUMBERS, SQUASH, MELONS: Aphids, Spider Mites - 1 1/2 pts.

COTTON: Aphids, Spider Mites - 1 1/2 pts. per acre in water to cover.

CLOVER: Aphids, Mites, Leafhoppers, Grasshoppers, use 1 1/2 to 2 1/4 pts. per acre in water to cover. Do not apply to Clover in bloom. May be grazed or harvested on day of application.

EGGPLANT: Spider Mites, Aphids - 1 pt. Lace Bugs - 1 1/2 pt.

ONIONS: Onion Thrips - 1 1/2 pts. in water to cover one acre.

PEAS: Pea Aphid - 1 1/2 pts. as needed. Do not apply within 3 days of harvest or 7 days if forage is to be fed to livestock.

PECANS: Spider Mites, Aphids - 1/2 to 2 pts.

PEPPERS: Aphids - 1 pt.

POTATOES: Aphids, Leafhoppers - 1 pint.

SEED ALFALFA, SEED CLOVER, SEED VETCH: Aphids, Spider Mites, Leafhoppers - 1 1/2 to 2 pts. in water to cover one acre.

STRAWBERRIES: Aphids, Spider Mites - 1 1/2 pints.

TOMATOES: Aphids - 1 pint. One application. Spider Mites - 1 1/2 pints.

WALNUTS: Aphids, Red Spider - 1 to 1 1/2 pints. Repeat at 7 to 10 days for Red Spiders.

FLIES: For use in and around buildings which house domestic animals, around yards and meat processing plants, and homes on out sites. Do not use when plant is in operation. Cover all equipment and processing surfaces or wash thoroughly before use. Use 2 gals. in 100 gals. water. Apply as a spray at the rate of 1 gal. per 1,000 sq. ft. on painted surfaces and 2 gals. per 1,000 sq. ft. on unpainted surfaces where flies alight or congregate, such as walls, ceilings, stanchions, windows in dairy barns, fences, around garbage cans, etc. As a floor treatment bait spray, use 5 ounces of MALATHION 5 EC with 1 cup corn syrup or sugar in 2 gals. of water. As a spot treatment this mixture can also be applied to windows, stanchions, support beams, doors, etc. For control of fly maggots, apply as a bait spray over the surface of manure or poultry droppings. In loafing sheds, spray the dry bedding within 18 inches of the walls and 3' and upright braces. For effective control in and around dairy barns and other agricultural premises, fly-breeding sites such as manure and other waste materials should be eliminated. Do not apply to freshly whitewashed surfaces. Wait 14 days before applying. Repeat applications as necessary. Avoid contamination of milk, milk equipment and water. AVOID contamination of feed and food products, also drinking fountains and feed troughs. Remove lactating dairy animals from buildings before treating, also remove animals under one month of age before treating.

ORNAMENTALS: Aphids, Spider Mites, Whitefly, Mealybugs, Thrip, Japanese Beetle adult, Four-lined Leaf bug, Tarnished plant bug, Rose leafhopper, Potato leafhopper, Scruffy scale, European pine shoot moth - 1 1/2 pts. Pine Needle Scale - 1 pt. Oak Kermes, Fletcher scale, Florida Red Scale, Boxwoods, Azalea Scale, Magalia Scale, Pine Leaf scale, Birch Leaf Miner, Boxwood leaf miner, Tent caterpillar - 2 pts. Evergreen scale - 1 to 1 1/2 pts. To control listed Scale - apply a thorough full coverage spray at time scale crawlers have settled on foliage. Do not use on Boston, Maidenhair and Pteris ferns, and some species of Casula.

LIVESTOCK: Do not treat calves under 1 month of age. Do not use on lactating or non-lactating dairy animals. Apply complete coverage spray. One treatment may be sufficient. Repeating, if needed. Avoid contamination of feed, food products, drinking fountains and feed troughs. Avoid contamination of milk, milk equipment and water.

BEEF CATTLE: Lice - 1 qt. per 100 gals. water. Ticks - 1 to 2 gals. per 100 gals. water. Back-rubbing devices; to reduce lice, use 9 1/6 tablespoonfuls or 4.58 fl. ozs. (7% aq. Malathion) in 1 qt. fuel oil. Mix thoroughly each time before use. There may also be tick infestation in horn ticks. These devices should be made conveniently a feasible, one to each 35 to 45 head of cattle and re-treated every 2 to 3 weeks. Do not apply on mice accessible to dairy animals.

HOGS: Lice - 1 gal. per 100 gals. water. Apply complete coverage also to pens and litter.

POULTRY (Direct application): Northern Fowl mite, Poultry Lice, and Chicken red mite (as a supplement to preens treatments). Use 6 teaspoonfuls per gallon of water per 100 to 150 birds. Repeat in 4 to 8 weeks, or when necessary.

POULTRY HOUSES: Northern Fowl mite, Chicken red mite, Poultry Lice, Flies - 4 1/2 tablespoonfuls per gallon water. Poultry: Ticks - 6 to 7 1/2 dunces per 1 gal. water. Repeat as necessary. Apply spray liberally to walls, ceilings and adjacent areas. Force spray into cracks and crevices. Use a high pressure sprayer for surface spray.

ROOST PAINTS: Chicken red mite, Poultry lice - 4 1/2 tablespoonfuls per 1 gal. water. Repeat as necessary. Brush on at rate of 1 pt. per 150 ft of roost.

1(f3)

**MILKOC GRANDE MALATHION**

**CAUTION:** Harmful if swallowed. Avoid prolonged breathing or spray mist. Avoid prolonged or repeated contact with skin. Wash thoroughly after using. Avoid contamination of feed and feedstuffs. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Do not apply under conditions involving possible drift to food, forage or other plantings, that might be damaged or the crop thereof rendered unfit for sale, use or consumption. Do not use in the edible product areas of meat or food processing plants.

This product is toxic to fish. Keep out of any body of water. Does exposed to direct treatment or residues on crops may be killed. For toxic information may be obtained from your Cooperative Agricultural Extension Service.

10% DESIGNATED AEROSOL IN AIR BALE. WATER UNLESS OTHERWISE SPECIFIED FOR GROUND APPLICATIONS. FOR AIR APPLICATION USE 3 TO 10% DESIGNATED BALE. CONSISTENT WITH DROPO GROWTH AND GOOD ILLUMINATION IN THE MISTBALE.

**NOTE** Before taking up your new assignment, you will receive a copy of the *Code of Ethics* and *Code of Professional Conduct* of the American Bar Association. You are expected to familiarize yourself with these codes and to observe them in all your professional conduct.

#### **ACTIVE INGREDIENT**

**Bis(4,6-dimethyl-Dinitiophosphate of D,L-  
succinate.)**

## **INERT INGREDIENTS:**

(Contains 5 pounds of Malathion per Gallon) - USE

DO NOT RE-USE EMPTY CONTAINERS OR DRUMS. DESTROY BY  
BURN OR DISCARD IN A SAFE PLACE. DRUMS MAY BE REUSED

10. WHICH TO APPLY: Serial, Classification, Grade, Priority, Category

Capri, Pepper, strawberries and green onions, Panached Brussels Sprouts, Chili, Cauliflower, Kale, Carrots, Onions, Tomatoes, Broccoli, Spinach.

*Lemons, Citrus, Apricots, Hot & cold fruits, & seeds, or nerves, Sausages, Melons, Peas, Potatoes, and vegetables, &c., & little if any of it can get lost, & there is not*

not within 7 days of harvest. Last return, the 8  
8% to be harvested on Dec 15, Rainy, few days no

APPLES: Codling moth, plum curculio, red scale, 10  
Bird cherry, Turner blight, woolly aphid, etc.

100% pts. shown above will be used  
injury to Retina and Cornea - 100%

