



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

October 24, 2017

Susan Watson  
Regulatory Manager  
Amrep, Inc.  
990 Industrial Park Drive  
Marietta, GA 30062

Subject: CSF Notification per PRN 98-10 –Revising basic and alt. CSF & minor label changes  
Product Name: CB-80-3 FOR INSECT CONTROL  
EPA Registration Number: 10807-450  
Application Date: 8/1/2017  
Decision Number: 532267

Dear Ms. Watson:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The CSFs submitted with your application have been stamped “Notification” and placed in our files.

The label submitted with the application has been stamped “Notification” and will be placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

Please note that the record for this product currently contains the following CSFs:

- Basic CSF dated 8/1/2017
- Alternate CSF 1 dated 8/1/2017

Page 2 of 2  
EPA Reg. No. 10807-450  
Decision No. 532267

Any CSFs other than those listed above are superseded/no longer valid. If you have any questions, please contact Angela Hollis at 703-347-0216 or by email at [hollis.angela@epa.gov](mailto:hollis.angela@epa.gov).

Sincerely

A handwritten signature in blue ink, appearing to read 'MW', is enclosed in a thin black rectangular box.

Michael Wagman, Acting Product Manager 4  
Invertebrate and Vertebrate Branch 1  
Registration Division (7505P)  
Office of Pesticide Programs

# CB-80-3 for Insect Control

## [Country Vet® FarmGard™ Farm & Dairy Fly Spray]

Pyrethrins are made from Chrysanthemum Flowers

- Provides Rapid Control of Residential, Industrial and Barn Pests
- Excellent Flushing
- 8 to 1 Synergist Ratio
- Quick Knockdown
- Nonaqueous
- 0.50% Pyrethrins for Quick Kill

### NOTIFICATION

10807-450

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

10/24/2017

**[KILLS and REPELS:** Flies, Mosquitoes, Wasps, Gnats, Cockroaches, Ants, Crickets, Spiders, Silverfish, Small Flying Moths, Indian meal Moths, Angoumois Grain Moths, Carpet Beetles, Confused Flour Beetles, Sawtoothed Grain Beetles, Granary Weevils, Rice Weevils, Red Flour Beetles, Grain Mites, Bedbugs, Fleas, Barn Flies, Deer Flies, Stable Files and Horn Flies.]

**[FOR USE IN:** Homes, Offices, Hotels, Motels, Schools, Hospitals, Nursing Homes, Day-care Centers, Factories, Warehouses, Stores, Supermarkets, Restaurants, Food Handling Establishments, Commercial or Institutional Kitchens, Food Processing Plants, Bottling Plants, Transportation Equipment (Automobiles, Buses, Trucks, Trailers, Boats, Ships, Trains, Aircraft), Barns, Stables, Livestock premises [quarters] on farms, Dairy Farms, Animal Quarters, Milkrooms, Poultry Houses, [as well as] Hog & Swine Operations and Beef Cattle Operations and Poultry Operations.]

*[For Reviewer: "Bulleted" items and items in [ ] above are optional and may/may not be included in part or in whole on final label.]*

#### ACTIVE INGREDIENTS:

Pyrethrins .....0.50%

\*Piperonyl Butoxide .....4.00%

**OTHER INGREDIENTS:** ..... 95.50%

**TOTAL** ..... 100.00%

\*Equivalent to 3.2% (butylcarbityl) (6-propylpiperonyl) ether and 0.8% of related compounds.

Contains Petroleum Distillates.

## KEEP OUT OF REACH OF CHILDREN

# CAUTION

See Back Panel for Additional Precautionary Statements.

**NET WEIGHT: 1 lb. (16 oz.) (453.6 g) [17 oz (482 g)]**

{Back Panel}

#### PRECAUTIONARY STATEMENTS

##### Hazards to Humans & Domestic Animals

**CAUTION.** Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

##### Personal Protective Equipment (PPE)

Mixers, loaders, applicators, and other handlers must wear the following: Long sleeve shirt, Long pants, Shoes and socks.

##### User Safety Requirements:

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

*[NOTE to Artist: The below paragraph must be enclosed in a box.]*

#### **USER SAFETY RECOMMENDATIONS:**

- User should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- User should remove clothing/PPE immediately if pesticide gets inside. Then **wash** thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

*[For Reviewer: Boxed & "Bulleted" format will be used if label space permits; otherwise a paragraph format will be used.]*

#### **FIRST AID**

**IF SWALLOWED:** Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-541-2016 for emergency medical treatment information. [For emergency information, call 1-800-222-1222, 24 hours, 7 days a week.]

**NOTE TO PHYSICIAN:** May pose an aspiration pneumonia hazard. Contains petroleum distillate.

#### **PHYSICAL OR CHEMICAL HAZARDS**

Extremely flammable. Contents under pressure. Keep away from fire, sparks and heated surfaces. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### **USE RESTRICTIONS**

For use as a spray: • Except when applied to livestock, do not enter or allow others to enter until vapors, mists, and aerosols have dispersed and the treated area has been thoroughly ventilated. • Do not remain in treated area. Exit area immediately and remain outside the treated area until aerosols, vapors, and/or mists have dispersed.

For cracks and crevice, surface or space spray uses: • Remove or cover exposed food and drinking water before application. • Remove or cover dishes, utensils, food processing equipment, and food preparation surfaces, or wash them before use.

For use in feed/food handling and processing facilities: • Do not make space spray applications when facility is in operation. • Prior to space spray applications, cover or remove food. • Prior to space spray applications, cover food processing surfaces or clean after treatment and before use.

Do not use in aircraft cabins.

When used in dairy barns or facilities: Close milk bulk lids to prevent contamination from spray and from dead or falling insects.

Remove or cover milking utensils before application. Wash teats of animals before milking.

Do not apply directly into sewers or drains, or to any area like a gutter where drainage to storm sewers, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

**FLYING INSECTS:** Flies, Mosquitoes, Small Flying Moths, Wasps, Gnats-Calculate cubic feet of space to be treated. Close all doors and windows. Spray in sweeping motion to all parts of the room. Apply for 2-3 seconds per 1,000 cubic feet of space. Sweep up fallen insects. Repeat treatment as necessary.

**CRAWLING INSECTS:** Cockroaches, Ants, Spiders, Silverfish-General Volumetric Treatment: Open cabinets and doors in area to be treated. Shut off air conditioners, fans, and close doors and windows before spraying. Spray for 20 seconds per 1,000 cubic feet of space. Direct spray toward areas suspected of harboring insects. Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Keep area closed for 15 minutes. Open and ventilate before reoccupying. Repeat treatment as necessary.

**FOR COCKROACHES:** For best results, spray directly on cockroaches and thoroughly spray all parts of the room suspected of harboring these pests. Special attention should be paid to cracks, hidden surfaces, under sinks, behind

stoves and refrigerators, food storage areas, and wherever these pests are suspected of hiding. Also, spray around door and window frames. Repeat as necessary.

**CRACK & CREVICE APPLICATION:** Food Areas and/or Food Handling Establishments-Spray in cracks and crevices, using the crack and crevice nozzle provided.

**FOR SPIDERS AND SILVERFISH:** For control of these insects, frequent and repeated spraying of baseboards, windows, doorframes, bookcases and storage areas should be made. For best results, spray directly on insects.

**FOR CRICKETS:** Spray thoroughly and frequently behind stoves, refrigerators, under cabinets, in food storage areas, and especially in closets, basements, and other damp areas. For best results, spray directly on insects.

**FOR ANTS:** Spray ant trails, nests and points of entry. For initial clean-up, apply for 10-15 seconds per 100 square feet of floor area. Then periodically spray as a space treatment for 3-5 seconds per 100 cubic feet of space.

**FOR INDIANMEAL MOTHS AND ANGOUMOIS GRAIN MOTHS:** Close doors and windows. Direct spray into all parts of room, especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1,000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed.

**FOR CONFUSED FLOUR BEETLES, SAWTOOTHED GRAIN BEETLES, GRANARY WEEVILS, RICE WEEVILS, RED FLOUR BEETLES, GRAIN MITES:** Spray into cracks and crevices, processing and handling equipment around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary. Infested stored food products in the home should be disposed of.

**FOR CARPET BEETLES:** Spray directly on beetles, baseboards, floors, shelves and carpets. Repeat treatment as necessary. ~~surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary. Infested stored food products in the home should be disposed of.~~

**FOR BEDBUGS:** Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Repeat treatment as necessary.

**FOR FLEAS:** Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. For best results, use with a product approved for application to pets.

**HOME USE:** To kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets-Spray the insect directly when possible and spray all probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves and under appliances. Repeat as necessary.

**Applications in Aircraft:** This product can be applied in cargo areas but not in aircraft cabins.

**FOOD PROCESSING PLANT USE:** To kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets-Spray the insect directly when possible. Spray all probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves, under appliances, and in food storage areas. To kill Flies, Mosquitoes, Gnats, Small Flying Moths-Close all windows and doors. Direct spray upward, spraying for 2-3 seconds for 1,000 cubic feet of space. Do not remain in treated areas.

Ventilate after treatment.

**VOLUMETRIC TREATMENT FOR INACCESSIBLE VOIDS (Attics, False Ceilings, Wall and Equipment Voids):** For crack and crevice application, calculate void volume in cubic feet. Treat 20 seconds per 1,000 cubic feet. Repeat as necessary.

**Volumetric Treatment of Food Areas and/or Food Handling Establishments:** Cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying food processing plants, thoroughly wash all equipment, benches, shelving, etc. where exposed food will be handled with an effective cleaning compound and rinse with potable water before use. **Volumetric Treatment in Homes:** In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

**NOTE,** when using this product in an enclosed area, such as milkroom, barn or stable, remove animals before spraying.

**DAIRY FARM USE:** Milking Parlor and Milkroom: Close all windows and doors. Direct spray upwards, spraying for 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Cover milking utensils and milk to prevent contamination from spray or dead and falling insects. **ANIMAL USE:** Thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders, and back where saliva accumulates from head tossing.

**BEEF CATTLE OPERATIONS:** In barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat application as necessary.

**ANIMAL USE:** For animal application, thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

**STANCHION BARN USE:** Walk behind animals and direct spray over backs, allowing 5 seconds per animal. For most effective results, apply each morning.

**HORSE BARN OR STABLES:** In horse and pony barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in the treated area. After 15 minutes, ventilate the area. Repeat the application as necessary. **ANIMAL USE:** For animal application, lightly mist over back of horses and ponies from a distance of 2 feet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

**HOG AND SWINE OPERATIONS:** In hog houses, close all doors and windows. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

**SWINE USE:** For animal application, direct spray over back of hogs and spray for 5 seconds per hog. Do not spray directly towards the animal's face or eyes. Repeat application daily or as necessary.

**POULTRY OPERATIONS:** In poultry houses, close all windows and doors. Spray roosts, walls, nests, and cages thoroughly. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated areas. After 15 minutes, ventilate the area. Treat daily or as necessary.

*[For Reviewer: Boxed & "Bulleted" format will be used if label space permits; otherwise a paragraph format will be used.]*

## STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Do not store near heat or open flame. Store in cool dry area away from children.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Do Not Puncture or Incinerate! **If empty:** Place in trash or offer for recycling, if available. **If partly filled:** Call your local solid waste agency for disposal instructions.

[Contains no CFCs or Other Ozone Depleting Substances.]

[Federal regulations prohibit CFC propellants in aerosols.]

[For Additional Products and Information, visit us at [www.Country-Vet.com](http://www.Country-Vet.com)]

*[For Reviewer: Optional-CFCs logo,*

*Optional-Premium Pyrethrum logo,*

*Steel graphics symbol]*



EPA Reg. No. 10807-450

EPA Est. No. 58996-MO-1 (A)

~~EPA Est. No. 10807-GA-1 (M)~~

~~EPA Est. No. 11623-GA-1 (O)~~

EPA Est. No. 11623-GA-2 (J)

~~EPA Est. No. 11623-GA-3 (P)~~

EPA Est. No. 33595-MO-4 (N)

Distributed by

**Amrep, Inc.**

990 Industrial Park Drive

Marietta, GA 30062

770.422.2071 • 800.241.7766