




sions of Sale, includ-  
sion, before using

reflect the opinion of  
laboratory animals  
conditions of use  
for which this product  
is used, it is impossible  
e of abnormal con-  
the Seller offer, and  
ding that the Buyer

use preparation that  
e and non-lactating  
e of an animal gives  
cation usually gives  
on at the rate of 12  
the correct amount

#### WILL TREAT:



tion uniformly along  
opped from any lean  
  
**DAIRY CATTLE FOR**  
treatment is important  
is soon as possible.  
such as bloat, should  
en cattle are treated  
is in the gutter or  
of the normal ranges  
of mat. feed, it  
decided to treat.

and the animal should  
the animal should  
not be treated  
and from less than 6  
least 6 weeks before



61556-32

# NEGUVON

Brand of  
**trichlorfon**

## CATTLE INSECTICIDE POUR-ON

**FOR CONTROL OF CATTLE GRUBS AND LICE**

#### GUARANTEE

ACTIVE INGREDIENT  
D-methyl-2,2-dichlorovinyl phosphate 2%

NETTING INGREDIENT

EPA Reg. No. 11556-32

PROTECT FROM EXTREME HEAT  
PROTECT FROM SUNLIGHT

**WARNING**  
KEEP OUT OF REACH OF CHILDREN

NET CONTENTS FIVE GALLONS

**NOTICE TO VETERINARIAN:** If the proper dosage of NEGUVON Pour-On has been applied and adverse reactions, such as opisthotonus, elevation and posterior paralysis occur, it is highly probable that a host-parasite reaction exists. Administer symptomatic treatment. Anti-inflammatory agents may be helpful. If necessary relieve bloat by the insertion of a stomach tube, may traumatize a severely swollen esophagus. Do not administer atropine, as it is contraindicated in host-parasite reactions. If toxicity should occur as a result of gross overdose, gastric lavage is indicated.

#### RESTRICTIONS

**TAKE TIME:** Do not treat lactating dairy cattle, animals less than 3 months old sick, convalescent, or stressed livestock.

Non lactating dairy cattle may be treated up to 7 days of freshening. If freshening should occur within 7 days after treatment, do not use milk for human food for the balance of the 7 day interval.

**OBSERVE LABEL DIRECTIONS:** Do not treat cattle within 21 days of slaughter.

Do not treat animals for 10 days before or after shipping or weaning or after exposure to contagious or infectious diseases.

**WARNING:** NEGUVON (brand of trichlorfon) is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals.

#### CONTAINER DISPOSAL

Do not reuse empty container. Destroy by perforating or crushing and bury in a safe place.

#### WARNING

May be harmful or fatal if swallowed. Harmful if absorbed through the skin. Do not get in eyes, on skin, or on clothing. When necessary, wash hands and wash with water after handling. Wash contaminated clothing with soap and hot water before reuse.

Do not contaminate food or feed, keep out of reach of children. If poisoning occurs, obtain prompt medical aid. Prolonged exposure to high concentrations of cholinesterase inhibitor.

#### WARRANTY AND LIMITATIONS OF DAMAGES

Chemagro warrants that the materials offered to the chemical industry, in the regular course of business, fit for the purpose intended to the satisfaction of the user and condition, if used according to the recommendations of the manufacturer. CHEMAGRO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, EXCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS FOR MERCHANTABILITY and except that Chemagro is authorized to do so, except as written with a specific reference to this warranty. Any damages arising from a breach of the warranty shall be limited to direct damages and shall not include consequential, commercial damages, or indirect or profit damages.

© 1970 Chemagro Corporation

61556-32

1556-32

JUL

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

1974

## USE ONLY AS DIRECTED

### DIRECTIONS FOR USE

**IMPORTANT:** Read these entire Directions and Conditions of Sale, including the Warranty and Limitation of Liability, enclosed in this package, before using NEGUVON Pour-On.

**CONDITIONS OF SALE** These Directions for use reflect the opinion of experts based on tests of effectiveness of toxicity to laboratory animals and of residues on grain fed and forage fed cattle and horses. However, the use of this wide range of conditions under which this product may be used, even though abnormal, does not allow it impossible to eliminate risks associated with its use because of abnormalities in the species being treated or the nature of the disease or infestation. The Buyer uses this product subject to the understanding that the Buyer assumes all risks.

NEGUVON Pour-On Cattle Insecticide is a ready-to-use preparation that requires no further dilution for use on beef cattle and non-lactating dairy cattle. A single application placed on the back of each animal gives effective control of cattle grubs. Also, a single application usually gives effective control of lice.

**AMOUNT TO USE** Apply NEGUVON Pour-On Cattle Insecticide at the rate of one gallon per acre.

THE CONTENTS OF THIS CONTAINER WILL TREAT

ANIMALS TO BE TREATED	NUMBER OF ANIMALS	NET CONTENTS
Beef cattle	1	1 GALLON

### HOW TO APPLY

### WHEN TO APPLY TO BEEF AND NON-LACTATING DAIRY CATTLE FOR GRUB AND LOUSE CONTROL

Apply NEGUVON Pour-On Cattle Insecticide to cattle after they have been shorn.

### NOTE

Do not apply to cattle during lactation.

Do not apply to cattle during pregnancy.

Do not apply to cattle during the last month of pregnancy.

Do not apply to cattle during the last month of lactation.

Do not apply to cattle during the last month of the breeding season.

Do not apply to cattle during the last month of the calving season.

Do not apply to cattle during the last month of the breeding season.

Do not apply to cattle during the last month of the calving season.

Do not apply to cattle during the last month of the breeding season.

Do not apply to cattle during the last month of the calving season.

Do not apply to cattle during the last month of the breeding season.

Do not apply to cattle during the last month of the calving season.

Do not apply to cattle during the last month of the breeding season.

Do not apply to cattle during the last month of the calving season.

Do not apply to cattle during the last month of the breeding season.

Do not apply to cattle during the last month of the calving season.

Chemagro

# NEGUVON

Brand of  
trichlorfon

## CATTLE INSECTICIDE POUR-ON

FOR CONTROL OF CATTLE GRUBS AND LICE

GUARANTEE

PROTECT FROM EXTREME HEAT PROTECT FROM SUNLIGHT

**WARNING** KEEP OUT OF REACH OF CHILDREN

NET CONTENTS ONE GALLON

NOTICE  
EXCESSIVE  
TOMATIC  
TRAUMATIC  
TOXICITY

TAK TIME  
OBSERVE LAW  
DIRECTIONS

DO NOT TI  
STICKERS  
WARNIN

RE

RE

RE

RE

RE

RE

RE

Chemagro

# NEGUVON

Brand of  
trichlorfon

## CATTLE INSECTICIDE POUR-ON

FOR CONTROL OF CATTLE GRUBS AND LICE

GUARANTEE



PROTECT FROM EXTREME HEAT PROTECT FROM SUNLIGHT

**WARNING** KEEP OUT OF REACH OF CHILDREN


NET CONTENTS ONE GALLON

**NOTICE TO VETERINARIAN:** If the proper dosage of NEGUVON Pour-On has been applied and adverse reactions, such as bloat, excessive salivation, and posterior paralysis occur, it is highly probable that a host parasite reaction exists. Administer symptomatic treatment. Anti-inflammatory agents may be helpful. If necessary, relieve bloat by trocarization; as a stomach tube may traumatize a severely swollen esophagus. Do not administer atropine, as it is contraindicated in host parasite reactions. If toxicity should occur as a result of gross overdosage, atropine is antidotal.

### RESTRICTIONS



Do not treat lactating dairy cattle, lambs less than 5 months old, sick, convalescent, or stressed livestock. Non-lactating dairy cattle may be treated up to 7 days of freshening. If freshening stock is sold within 7 days after treatment, do not use milk for human food for the balance of the 7 day interval.



Do not treat cattle within 14 days of slaughter. Do not treat animals > 10 mos. of age after stripping or weaning. Do not expose to cold air, sun, or wind during diseases.

**WARNING:** NEGUVON Pour-On treatment is a cholinesterase inhibitor. Do not use this product on mammals previously treated with or few days before or after treatment with or exposure to cholinesterase inhibiting drugs, such as organophosphates.

### CONTAINER DISPOSAL

Empty containers should be discarded in accordance with local, state, and federal regulations.

### WARNING

May irritate eyes. Avoid contact with eyes. If contact occurs, flush eyes immediately with water. If irritation persists, seek medical advice. Avoid contact with skin. If contact occurs, wash thoroughly with soap and water. If irritation persists, seek medical advice.

Do not drink from treated water troughs. Do not drink from treated tanks. Do not drink from treated streams. Do not drink from treated ponds.

### WARRANTY AND LIMITATIONS OF DAMAGES

Chemagro warrants that this product will be free from defects in material and workmanship for a period of one year from the date of manufacture. This warranty does not apply to damage resulting from misuse, abuse, or negligence. Chemagro's liability under this warranty is limited to the repair or replacement of the defective product. Chemagro shall not be liable for any incidental or consequential damages resulting from the use of this product. Chemagro makes no other warranties, express or implied, including the implied warranties of merchantability and fitness for a particular purpose.

**CHEMAGRO**