

BEST DOCUMENT AVAILABLE

ACCEPTED

11731-4

APR 24 1979

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO.

KEEP

OUT OF

REACH

OF

CHILDREN

CAUTION

No. 820 Industrial Microbiocide

FOR INDUSTRIAL USE ONLY. NOT FOR DOMESTIC USE. SEE DATA SHEET.

E.P.A. EST. NO.

E.P.A. REG. NO. 11731-4

ACTIVE INGREDIENTS:

Poly [oxyethylene (dimethylimino) - ethylene (dimethylimino) ethylene dichloride] 20%

INERT INGREDIENTS:

80%
This product contains 1.74 lbs. of active ingredient per gallon and weighs 8.69 lb./gal.

PRECAUTIONARY STATEMENTS:

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes or clothing.

FIRST AID: If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds or public water. For guidance, contact the regional office of the Environmental Protection Agency

Manufactured by:

AMERICAN *Water Treatment, Inc.*

7116 OAKLAND AVENUE
ST. LOUIS, MISSOURI 63117

DIRECTIONS FOR USE General Classification

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

No. 820 Microbiocide is used to control algae bacteria and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algae growth, microbiological slime and other deposits.

An initial slug addition of 2.9 to 7.3 fluid ounces of No. 820 Microbiocide per 1,000 gallons of water to provide a concentration of 24 to 60 parts per million of No. 820 Microbiocide based on the total weight of water in the system, is recommended. Repeat initial dosage until control is evident.

Subsequent slug additions of 0.73 to 8.3 fluid ounces of No. 820 Microbiocide per 1,000 gallons of water (6 to 60 parts per million) of No. 820 Microbiocide should be employed every 2 to 5 days or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

STORAGE AND DISPOSAL:

Keep container closed when not in use. Do not contaminate water, food or feed by storage, disposal, or cleaning of equipment. Triple rinse (or equivalent) and dispose of container in an incinerator or in an approved landfill or bury in a safe place. Rinse that cannot be used or reprocessed should be disposed of in a landfill approved for pesticides or bury in a safe place away from water supplied. Open dumping is prohibited.