

DIRECTIONS FOR USE:

INITIAL DOSE:

When the system is noticeably fouled, apply 2 1/2 to 5 pints of B-0035, Slime Control, per 1,000 gallons of water in the system. Repeat until control is achieved. Add this material to makeup water inlet for uniform mixing.

SUBSEQUENT DOSE:

When microbial control is evident, add 1 to 2 1/2 pints of B-0035, Slime Control, per 1,000 gallons of water in the system every 5 days or as needed to maintain control. Badly fouled systems must be cleaned before treatment is begun.

CAUTION!

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

ACCEPTED
MAR 2 1981
Fungicide
Registered
EPA Reg. No. 14886-2

B-0035

SLIME CONTROL

For Control of Bacterial and Algae Slime Growth in Recirculating Industrial Cooling Water Systems.

ACTIVE INGREDIENTS

Alkyl (50%), Cu (40%), Cu (10%), C (dimethyl benzyl amine) (10%) 9.88%

INERT INGREDIENTS

EPA Reg. No. 14886-2 EPA Establishment No. 14836-NC-1 90.12%

DANGER

KEEP OUT OF REACH OF CHILDREN.

See right side of panel for additional precautionary statement
Container Disposal: When empty, rinse container thoroughly with water and destroy it by crushing or burying it in a safe place away from water supplies.

MANUFACTURED BY

PARKER ENGINEERING & CHEMICALS, INC
320 Hodgin Street P.O. Box 869
High Point, North Carolina 27261

NET CONTENTS

31

DANGER!

Keep out of reach of children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

STORE IN AREA ABOVE 32°F