



ACCEPTED

DEC 16 1997

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 19713-34

Methoxychlor 2 E.C.

Emulsifiable Insecticide

For control of various insects on livestock, ornamentals and vegetable crops.

ACTIVE INGREDIENT:

Methoxychlor: 1,1,1-trichloro-2,2-bis (p-methoxyphenyl) ethane*	25.0%
OTHER INGREDIENTS:**	75.0%
TOTAL:	100.0%

*Equivalent to 22% - 1,1,1-trichloro-2,2 bis(p-methoxyphenyl) ethane, 3% other isomers and related compounds.
 **Contains Petroleum Distillates.
 This product contains 2 lb. technical methoxychlor per gallon.

KEEP OUT OF REACH OF CHILDREN
CAUTION
See First Aid Below

EPA Reg. No. 19713-34
EPA Est. No. 19713-GA-2 Net Contents: _____

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center. Vomiting may cause aspiration pneumonia. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove to fresh air.

IF IN EYES: Immediately flush eyes with plenty of water and get medical attention.

IF ON SKIN: Wash thoroughly with soap and warm water. Get medical attention if irritation persists.

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals.

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Do not breathe spray. Keep away from food, feedstuffs or water supplies.

PERSONAL PROTECTIVE EQUIPMENT (PPE)
Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear: Long-sleeved shirt and long pants, chemical-resistant gloves such as Barrier Laminate or Viton and shoes plus socks.

For exposure in enclosed areas: A respirator with either an organic vapor-removing cartridge with a pre-filter approved for pesticides (MSHA-NIOSH approval number prefix TC-14G).

For exposure outdoors: Dust/mist filtering respirator (MSHA / NIOSH approved prefix TC-21C).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should: 1) Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. 2) Remove clothing immediately if pesticide gets inside. 3) Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Aerial application over such sites is permissible only when they are not visible from above the tree canopy. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS
Do not use, pour, spill or store near heat or open flame.

GENERAL INFORMATION
METHOXYCHLOR 2 E.C. should be used only in accordance with recommendations on this label.

NOTE: PHI stands for Pre-Harvest Intervals.

METHOXYCHLOR 2 E.C. contains 2 lbs. technical methoxychlor per gallon. On mixing with water it forms a stable emulsion for the control of various insects. It is suitable for use in either high or low volume application equipment. When applying METHOXYCHLOR 2 E.C., use sufficient spray volume to thoroughly cover all surfaces to be treated.

Do not use this product with any spray material(s) that is not compatible with oil. Do not apply within 14 days of sulfur or sulfur containing products.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. **DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.** Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural pesticides. It contains the requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: Coveralls, chemical-resistant gloves such as Barrier Laminate or Viton and shoes plus socks.

Manufactured By:
Drexel Chemical Company
P.O. BOX 13327, MEMPHIS, TN 38113-0327
SINCE 1972

RECOMMENDATIONS FOR USE

Apply METHOXYCHLOR 2 E.C. at recommended rates with ground or aerial equipment to thoroughly cover foliage and fruit. Mix METHOXYCHLOR 2 E.C. in water using the appropriate volume of water for equipment to be used (for example, 3 to 20 gallons of water per acre for aircraft or ground concentrate sprayers and greater amounts for high volume sprayers). Begin application at first sign of infestation (for fruits, begin at petal fall) and repeat at 7 to 14 day intervals or as needed. Do not apply within the number of days to harvest as indicated in () immediately following each.

ASPARAGUS (3): *Asparagus beetles* - Use 2 to 4 quarts per acre. If applied within 3 days of harvest, remove residues by washing or blanching.

APPLES (7), PEARS (7), QUINCE (7): *Apple maggots, Codling moths, Japanese beetles, Plum curculios, Tent caterpillars* - Use 7½ to 15 quarts per acre or 2 to 3 quarts per 100 gallons water at 350 to 500 gallons per acre. Repeat at 7 to 10 day intervals.

APRICOTS (21), CHERRIES (7), NECTARINES (21), PEACHES (21), PLUMS (7), PRUNES (7): *Cherry fruitworms, Cherry fruitflies, Japanese beetles, Plum curculios, Rose chafers, Tent caterpillars, Cankerworms* - Use 7 to 15 quarts per acre, or 2 to 3 quarts per 100 gallons water at 350 to 500 gallons per acre. Repeat at 7 to 10 day intervals.

BEANS (3), BLACK-EYED PEAS (3): *Corn earworms, Rose chafers, Cowpea curculios, Imported cabbageworms, Cross-striped cabbageworms, Cucumber beetles, Fall armyworms, Alfalfa and Garden webworms, Flea beetles, Mexican bean beetles, Potato leafhoppers, Bean leaf beetles, Japanese beetles* - Apply 2 to 6 quarts per acre. Do not apply within 7 days of harvest if vines are to be used for feed or forage. Repeat at 7 to 10 day intervals.

BROCCOLI (14), BRUSSELS SPROUTS (14), BEETS (Roots-14, Tops-14), CABBAGE (3), CAULIFLOWER (7), CARROTS (14), COLLARDS (14), EGGPLANTS (7), KALE (14), KOHLRABI (7), LETTUCE (14), PEPPERS (7), RADISHES (7), RUTABAGAS (7), SPINACH (14), TURNIPS (Roots-7, Tops-14): *Blistar beetles, Flea beetles, Leafhoppers, Alfalfa loopers, Fall armyworms, Japanese beetles, Cross-striped cabbageworms, Imported cabbageworms* - Apply 2 to 4 quarts per acre.

BLACKBERRIES (14), LOGANBERRIES (14), RASPBERRIES (14), BOYSENBERRIES (14), DEWBERRIES (14), YOUNGBERRIES (14), STRAWBERRIES (14): *Rose chafers, Strawberry weevils, Flea beetles, Omnivorous leafhoppers, Spittlebugs, Japanese beetles* - Apply 2 to 3 quarts 100 gallons per acre.

BLUEBERRIES (14), CURRANTS (14), GOOSEBERRIES (14): *Japanese beetles, Cranberry fruitworms, Leafhoppers, San Jose scale (crawlers)*: 2 to 6 quarts per acre, or 2 to 3 quarts per 100 gallons of water at 100 gallons per acre.

CANTALOUPE (7), CUCUMBERS (7), MELONS (7), PUMPKINS (7), SQUASH (7): *Squash vine borer, Fall armyworm, Melon worm, Cucumber beetles, Flea beetles* - 2 to 6 quarts per acre. May be applied within one day of harvest if 3 quarts per acre or less is used.

GRAPES (14): *Berry moths, Grape leafhoppers, Japanese beetles, Leaf skeletonizers, Rose chafers* - Use 2 to 6 quarts per acre or 2 to 3 quarts per 100 gallons of water at 100 to 200 gallons per acre.

CORN - Field & Sweet (7): *Fall armyworms, Flea beetles, Japanese beetles, Rose chafers* - Use 2 to 4 quarts per acre. Do not feed treated corn to livestock.

CORN - Sweet (7): *Corn earworms* - Apply to silks as soon as they appear. Make 3 to 4 applications at 2 to 3 day intervals as infestation warrants. Re-apply immediately after a rain. Do not feed treated corn to livestock.

LIVESTOCK (Beef and Dairy Cattle, Goats, Sheep and Swine) - Apply the following rates of METHOXYCHLOR 2 E.C. for the control of hornflies, lice, ticks, keds and fleas.

Beef Cattle - Mix 2 quarts in 25 gallons of water. As a spray, use maximum of 2 quarts of the mixture per head for larger animals and proportionally less for smaller ones. Thoroughly wet animals. Do not use in dipping vats. Wait 3 weeks and re-apply if necessary.

Spot Treatment for Tail Lice - Mix 4 to 6 quarts in 25 gallons of water and apply as a spray on the infested tail only. Do not apply to animals being finished for slaughter. Do not contaminate feed or water.

Dairy Cattle (Non-Lactating) - Mix 2 quarts in 25 gallons of water. As a spray use 2 quarts of the mixture per head for larger animals and proportionally less for smaller ones. Thoroughly wet animals. Do not use as a dip. Do not apply during lactation or within 2 weeks of freshening.

Goats, Sheep and Swine - Mix 2 quarts in 25 gallons of water. As a spray thoroughly wet animals. Do not apply to lactating goats.

NOTE: This product is toxic to fish. Keep out of lakes, streams and ponds. Rates recommended on this label are toxic to shrimp and crabs. Do not apply where these species are an important resource. Do not apply to water drainage areas where runoff or flooding will contaminate ponds.

FOREST AND SHADE TREES: For control of the following insects, use a 6% methoxychlor solution with a mist blower at rates recommended in the following table.

Mix 1 gallon METHOXYCHLOR 2 E.C. per 3 gallons of water for mist blower solution, or 1 to 2 quarts of METHOXYCHLOR 2 E.C. per 100 gallons of water for conventional sprayer.

Insects	Pints of 6% Solution Per Tree Tree Height				Gals. 6% Solution / A
	35-50'	50-65'	65-80'	80-120'	
Cankerworm	2	2	3	4	2
Eastern Tent Caterpillar	2	2	3	4	2
Elm Leaf Beetle	2	4	6	8	
Fall Webworm	½	1	2	2	1
Forest Tent Caterpillar	1	1½	2	3	2
Gypsy Moth	½	1	2	2	2
Japanese Beetle	2	4	4	4	3
Lace Bugs (Oaks and Sycamore)	1½	2	3	4	2
May Beetle	1½	2	3	4	2
Tussock Moth	2	2	3	4	2

Control of Elm Bark Beetle (Vectors of Dutch Elm Disease) - Apply first application before elm flowers or leaves sprout using a 12% spray solution of METHOXYCHLOR 2 E.C. with a mist blower or a 2% spray solution in a conventional sprayer. This application will usually be applied in March in the Southern States and in April in the Northern States. Use 2 to 3 gallons of spray solution, wetting all surfaces thoroughly. Repeat at 7 to 14 day intervals. Second application should be made 2 to 3 months after the first treatment. Reduce the active ingredient to 6% for mist blower and 1% for conventional sprayer. Cover all leaf and bark surfaces.

Control of Elm Leafhopper (Vectors of Elm Phloem Necrosis) - Apply first application when elm leaves are fully grown. Use a 6% spray solution in a conventional sprayer. This application will be made in May in the Southern States and in June in the Northern States. Thoroughly cover all leaf surfaces. When second flush of growth appears, usually 1 to 2 months after first treatment, repeat the above application covering all leaf surfaces thoroughly. In those states where both the elm leafhopper and elm leaf beetle are known to be present a three spray schedule will provide effective control. Apply protective spray before elm flowers or leaves sprout, follow with second protective spray 2 to 3 months after the first. Then apply the third protective spray about 1 to 3 months after the second protective spray. Be sure to thoroughly cover all bark and foliar surfaces with each protective spray.

Preparation of Spray Solutions

12% spray - dilute 1 part METHOXYCHLOR 2 E.C. with 1 part water.
6% spray - dilute 1 part METHOXYCHLOR 2 E.C. with 3 parts water.
2% spray - dilute 1 part METHOXYCHLOR 2 E.C. with 12 parts water.
1% spray - dilute 1 part METHOXYCHLOR 2 E.C. with 24 parts water.

SHRUBS: *Tent caterpillars, Cankerworms, Japanese beetles* - Use 2 to 3 quarts per 100 gallons of water. Spray thoroughly to runoff. Begin applications when insects first appear and repeat at 6 to 14 day intervals as needed.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

STORAGE INSTRUCTIONS

Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool dry area away from any heat or ignition source. Avoid storage at high temperatures. Do not move containers from one area to another unless they are securely sealed. Keep container tightly sealed when not in use. Keep away from any puncture source. Avoid storage near water supplies, food, feed and fertilizer to avoid contamination. Avoid contamination with acids or alkalis. Store in original containers only. If the contents are leaking or material is spilled, follow these steps:

1. Contain spill, absorb with a material such as saw dust, clay granules or dirt.
2. Collect and place in suitable containers for disposal.
3. Wash area with water and soap to remove remaining pesticide.
4. Follow washing with clean water rinse.
5. Place a leaking container in a plastic tub and transfer contents as soon as possible to an empty original container.
6. Do not allow runoff to enter sewer or contaminate water supplies.
7. Dispose of waste by burying in a noncrop area away from water supplies or by incineration.

(Continued on page 3)

STORAGE AND DISPOSAL

(Continued from page 2)

PESTICIDE DISPOSAL

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration, or, if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

WARRANTY - CONDITION OF SALE

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the Seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

In no case shall Drexel or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by Drexel Chemical Company and is accepted as such by the Buyer.

3 of 3